

FINITE ELEMENT HETEROGENEOUS MULTISCALE METHODS WITH NEAR OPTIMAL COMPUTATIONAL COMPLEXITY

ASSYR ABDULLE * AND BJORN ENGQUIST †

Abstract. This paper is concerned with a numerical method for multiscale elliptic problems. Using the framework of the Heterogeneous Multiscale Methods (HMM), we propose a micro-macro approach which combines finite element method (FEM) for the macroscopic solver and the pseudo-spectral method for the micro solver. Unlike the micro-macro methods based on standard FEM proposed so far in HMM we obtain, for periodic homogenization problems, a method that has *almost-linear* complexity in the number of degrees of freedom of the discretization of the macro (slow) variable.

Key words. multiscale method, heterogeneous FEM, spectral method, elliptic homogenization

AMS subject classifications. 65N30,74Q05,74Q15,74Q20,39A12

1. Introduction. The numerical solution of problems encompassing a variety of strongly coupled scales pose major computational challenges in terms of analysis modeling and simulation. The direct numerical simulation of problems for which significant physical phenomena occur on length scales which differ by several orders of magnitude is often impossible, due to the computational cost for resolving the smallest scale.

For problems with scale separation, mathematical tools, as homogenization theory, have been developed to derive “macro” or “effective” or “homogenized” models (see [7], [21], [11] and the references therein). The numerical simulation of these macro models can be done by standard methods. However, beside restrictive assumptions on the media, the simulation techniques based on this macro models suffer from several drawbacks. The parameters of the effective models have usually to be computed numerically so that it is difficult to obtain an error control of the discretized macro model. Furthermore, the small scale information is lost in these approaches. It can be recovered by solving corrector problems, but this procedure is as expensive as solving the full fine scale problem.

Introduced in [13], the Heterogeneous Multiscale Method (HMM) has been proved to be a useful framework for the design and analysis of multiscale methods. Such methods based on finite elements, the so-called finite element heterogeneous multiscale methods (FE-HMM) have been developed in [2],[3],[4],[5],[14].

In these approaches, the unknown effective problem is solved directly with a coarse mesh by a macro FE solver. The unknown data of the macro model are extracted on the fly by testing the microstructure on sampling domains with a micro finite element method (FEM). Recently, a fully discrete error analysis has been derived for these type of methods [3],[4],[5]. The analysis in the aforementioned paper shows that for N “macro” degrees of freedom, the overall complexity, taking into account the micro FE discretization, is superlinear in N . More precisely, consider piecewise linear FE space for the micro and macro problem and denote by N_{mac} and N_{mic} the degrees of freedom of the *macro* FEM and the *micro* FEM, respectively. Assume further that the costs (floating point operations) of the methods are proportional to their degrees of freedom (as for example when using multigrid linear solver). Since the macro FEM

*School of Mathematics, University of Edinburgh EH9 3JZ, Scotland, a.abdulle@ed.ac.uk

†Mathematics Department, The University of Texas at Austin, University Station C1200, Austin, TX 78712-0257,engquist@

has input data coming from the micro problems, the numerical error when solving these latter problems has an impact on the global error. For the approximation of the homogenized (upscaled) problem, the global error for solving numerically a multiscale elliptic problem with an HMM type method is, as shown in [3] given by $err_{mac} + err_{mic} + err_b$, where err_{mac} is the error of the macro FEM, err_{mic} is the error contribution from the micro FEM and err_b comes from the (possibly inexact) boundary conditions and boundary layer terms [14]. We note that $err_b = 0$ with an heterogenous multiscale type method for the numerical approximation of periodic homogenization problems [2],[3]. When using linear FEM for the micro problem, it has been shown in [3] that $err_{mic} = \mathcal{O}(N_{mic}^{-\frac{2}{d}})$. Thus, for solving elliptic homogenization problems in the L^2 norm with the usual quadratic convergence rate we should use $N_{mic} \simeq N_{mac}$ and the total cost is $\mathcal{O}(N_{mac}^2)$, while in the H^1 norm (with a linear convergence rate) we can use $N_{mic} \simeq N_{mac}^{1/2}$ and the total cost is $\mathcal{O}(N_{mac}^{3/2})$. Finally if we want to approximate the fine scale solution, a reconstruction procedure has been proposed in [13],[1] and the analysis in [3] shows that the total cost (for linear convergence rate in the H^1 norm) is $\mathcal{O}(N_{mac}^2)$. Notice that in this latter case, $err_b = \sqrt{\varepsilon}$, where ε is the length of the small scale oscillation.

In this paper, we propose and analyze a numerical method for elliptic multiscale problems based on a coupling of a macro FEM with a micro pseudo-spectral method. Provided sufficient regularity of the conductivity tensor, we show that the micro solution has *spectral accuracy*. Furthermore, if the the conductivity tensor is analytic, we show that the micro solution has *exponential convergence*. In this latter situation, the overall complexity is quasi optimal, i.e. *almost-linear* in the number of degrees of freedom N_{mac} . More precisely, using the above notation, we show that err_{mic} decreases with a spectral or an exponential rate (these statements about spectral and exponential convergence rates will be made precise in the proofs).

For important classes of problems, including problems with periodic coefficients, with random stationary coefficients as well as for some nonlinear problems, the micro problems can be defined in periodic function spaces [13],[14],[2],[3],[4],[5]. The use of pseudo-spectral methods is thus well-suited for these micro problems. Furthermore, using spectral method on the micro domains (which can be chosen as squares/cubes) does not prevent to apply the numerical method to domains with complicated geometry. It is the macro triangulation which meshes the domain of the physical problem. Beside the optimal complexity for two scale problems, let us mention several issues in HMM that can be addressed by the method proposed in this paper. An important issue in multiscale computation is high order methods. Theoretically, FE-HMM can be easily constructed using several sampling domains within each macroelements [14]. However, the fully discrete analysis [3] shows that higher order micro solvers should also be implemented, for otherwise, the computational complexity will be governed by err_{mic} , the contribution of the low order micro solvers. By using spectral methods for the micro solver, it is possible to construct efficient high order FE heterogeneous multiscale methods. Another issue is the numerical solution of problems with more than two (separated) scales. For such problems, the FE-HMM consists in a hierarchy of micro-macro methods. Refining *simultaneously* the whole hierarchy of meshes (based on the rate derived in [3],[4],[5]) can be computationally expensive. In this situation, almost-linear complexity with respect to the macro degrees of freedom N could also be achieved by coupling a macro FEM for the physical domain with pseudo-spectral methods for the hierarchy of micro scales.

The paper is organized as follows. In Section 2 we introduce the multiscale method

based on the coupling of a macro FEM and micro pseudo-spectral Fourier methods (FES-HMM), we state the main results and put them in perspective with previously obtained results. Section 3 is devoted to the analysis of the method and the proof of convergence results. In Section 4 we extend our results for higher order macro FEM. Finally, in Section 5, we present numerical examples which illustrate the convergence rates of our method.

Notations. In what follows, $C > 0$ denotes a generic constant, independent of ε , whose value can change at any occurrence but depends only on the quantities which are indicated explicitly. For $r = (r_1, \dots, r_d) \in \mathbb{N}^d$, we denote $|r| = r_1 + \dots + r_d$, $D^r = \partial_1^{r_1} \dots \partial_d^{r_d}$. We will consider the usual Sobolev space $H^s(\Omega) = \{u \in L^2(\Omega); D^r u \in L^2(\Omega), |r| \leq s\}$, with norm $\|u\|_{H^s(\Omega)} = (\sum_{|r| \leq s} \|D^r u\|_{L^2(\Omega)}^2)^{1/2}$. We will also consider $H_0^1(\Omega)$ the closure of $C_0^\infty(\Omega)$ for the $\|\cdot\|_{H^1(\Omega)}$ norm and the spaces $W^{l,\infty}(\Omega) = \{u \in L^\infty(\Omega); D^r u \in L^\infty(\Omega), |r| \leq l\}$. We will also consider $W_{per}^s(Y) = \{v \in H_{per}^s(Y); \int_Y v dx = 0\}$, where $H_{per}^s(Y)$ is defined as the closure of $C_{per}^\infty(Y)$ (the subset of $C^\infty(\mathbb{R}^d)$ of periodic functions in the unit cube $Y = (0, 1)^d$) for the H^s norm. For $s = 0$, we will denote $H_{per}^0(Y) = L_{per}^2(Y)$. Finally we consider $C_{per}^0(Y)$, the set of continuous periodic complex-valued functions in Y .

2. Spectral Heterogeneous Multiscale Finite Element Method. In this section we first briefly recall homogenization theory and pseudo-spectral methods and introduce the multiscale FEM with micro problems solved by the pseudo-spectral Fourier method (FES-HMM). At the end of the section we state the main convergence results. Although the analysis of our method will be performed in the periodic homogenization framework, we emphasize that the numerical method itself *is not restricted* to periodic homogenization problems.

2.1. Homogenization Problems. We consider the following elliptic model problem in the domain $\Omega \subset \mathbb{R}^d$

$$(2.1) \quad -\nabla \cdot (a^\varepsilon \nabla u^\varepsilon) = f \text{ in } \Omega, \quad u^\varepsilon = 0 \text{ on } \partial\Omega,$$

where ε is a small parameter that represents explicitly the multiscale nature (eg. small scale) of the problem. Throughout, we will assume that the tensor $a^\varepsilon(x)$ is uniformly elliptic and bounded, i.e.,

$$(2.2) \quad \gamma_1 |\xi|^2 \leq \xi^T a^\varepsilon(x) \xi \leq \gamma_2 |\xi|^2 \quad \forall \xi \in \mathbb{R}^d \text{ and a.e. } x \in \Omega.$$

For several classes of such multiscale problems, it is known from homogenization theory (see e.g. [7, Chap.1],[26]) that u^ε converges (usually in a weak sense) to a ‘‘homogenized solution’’ u^0 , solving an elliptic problem where the small scale have been averaged out.

The analysis of our methods will be presented for the case when the tensor $a^\varepsilon(x) = a(x, \frac{x}{\varepsilon}) = a(x, y)$ is symmetric, coercive and periodic with respect to each component of y in the cube $Y = (0, 1)^d$, $f \in L^2(\Omega)$, $a_{ij}(x, \cdot) \in L^\infty(\mathbb{R}^d)$ and $x \rightarrow a_{ij}(x, \cdot)$ is smooth from $\bar{\Omega} \rightarrow L^\infty(\mathbb{R}^d)$. In this situation, u^ε converges weakly in $H_0^1(\Omega)$ to u^0 , solution of the homogenized problem

$$(2.3) \quad -\nabla \cdot (a^0(x) \nabla u^0) = f(x) \in \Omega, \quad u^0 = 0 \quad \text{on } \partial\Omega,$$

where the homogenized diffusion coefficient a^0 is a smooth matrix with coefficients given by $a_{ij}^0(x) = \int_Y \left(a_{ij}(x, y) + \sum_{k=1}^n a_{ik}(x, y) \frac{\partial \chi^j}{\partial y_k}(x, y) \right) dy$. Here, $\chi^j(x, \cdot)$ denote

the solutions of the so called cell problems

$$(2.4) \quad \int_Y \nabla \chi^j a(x, y) \nabla v dy = \int_Y (a(x, y) e_j)^T \nabla v dy, \quad \forall v \in W_{per}^1(Y), \quad j = 1, \dots, d,$$

where $(e_j)_{j=1}^d$ is the canonical basis of \mathbb{R}^d . Classical homogenization theory [7],[26],[21] gives

$$(2.5) \quad \|u^\varepsilon - u^0\|_{L^2(\Omega)} \leq C\varepsilon.$$

Some regularity on $\chi^j(x, \cdot)$ is needed for this estimate. The assumption $\chi^j(x, \cdot) \in W^{1,\infty}(Y)$ together with the assumption that the homogenized solution u^0 belongs to $H^2(\Omega)$ are sufficient for estimation (2.5) to hold (see [21, Section 1.4],[18, Remark 3.3],[17, Remark 7]). In the nonperiodic case, there still exists a homogenized problem, but the homogenized matrix $a^0(x)$ is usually unknown [21]. We emphasize that our numerical method is not restricted to the periodic case (see Remark on page 6 and Section 5.3).

2.2. Spectral and pseudo-spectral Fourier approximation. We consider the Hilbert space $L^2(Y)$, the space of Lebesgue measurable square integrable functions $u : Y \rightarrow \mathbb{C}$, where $Y = (0, 1)^d$ with the scalar product $(u, v) = \int_Y u \bar{v} dy$ and the norm $\|u\|_{L^2(Y)}^2 = (u, u)$.

Let us first suppose that $d = 1$ and we set $I = (0, 1)$. We recall that

$$(2.6) \quad u \in L^2(I) \iff u = \sum_{k=-\infty}^{k=\infty} \hat{u}_k e^{2ik\pi y} \quad \text{with} \quad \sum_{k=-\infty}^{k=\infty} |\hat{u}_k|^2 < \infty.$$

The coefficients \hat{u}_k are given by $\hat{u}_k = (u, e^{2ik\pi y})$. We further consider for an integer $M > 0$ the subspace spanned by $\text{span}\{\psi_k = e^{2ik\pi y}; |k| \leq M\}$ and the L^2 -orthogonal projection

$$(2.7) \quad P_M(u) = \sum_{k=-M}^{k=M-1} \hat{u}_k \psi_k.$$

In general it is not possible to calculate explicitly the Fourier coefficients \hat{u}_k of the orthogonal projection. Let us define a mesh on I with *pseudospectral points* given by

$$(2.8) \quad I_M := \{y_l = lh, l = 0, 1, \dots, 2M-1, h = 1/2M\}.$$

We next consider the so-called *pseudo-spectral method* which is a collocation procedure at the pseudospectral points y_l of defined above. The discrete Fourier coefficients of a function $u \in C_{per}^0(I)$ with respect to the pseudospectral points I_M are given by

$$(2.9) \quad \tilde{u}_k = \frac{1}{2M} \sum_{l=0}^{2M-1} u(y_l) e^{-2ik\pi y_l},$$

and the trigonometric interpolant of u at the pseudospectral points I_M is defined by

$$(2.10) \quad u_M := Q_M(u) = \sum_{k=-M}^{k=M-1} \tilde{u}_k \psi_k,$$

Notice that $Q_M(u(y_l)) = u(y_l)$, $\forall l = 0, 1, \dots, 2M - 1$. In higher dimension $d > 1$, we use the tensor product interpolant

$$(2.11) \quad \mathbf{Q}_M(u) = (Q_M^1 \otimes \dots \otimes Q_M^d)(u),$$

where Q_M^i are given by (2.10). The d -dimensional pseudospectral mesh on $Y = (0, 1)^d$ is defined as tensor product of the 1-dimensional pseudospectral points

$$(2.12) \quad Y_M := \{(y_{l_1}, \dots, y_{l_d}); y_{l_i} \in I_M\}.$$

2.3. Spectral Heterogeneous Multiscale Finite Element Method. We construct now the multiscale FEM with micro problems solved by the pseudo-spectral Fourier method. We concentrate here for simplicity on piecewise linear continuous FEM in the macro spaces. We will consider higher order FEM in Section 4. We assume in the sequel that the domain Ω where the multiscale problem is defined is a convex polygon. Let therefore the macro finite element space be defined by

$$(2.13) \quad S_0^1(\Omega, \mathcal{T}_H) = \{u^H \in H_0^1(\Omega); u^H|_K \in \mathcal{P}^1(K), \forall K \in \mathcal{T}_H\},$$

where $\mathcal{P}^1(K)$ is the space of linear polynomials on the triangle K , and \mathcal{T}_H is a quasi-uniform triangulation of $\Omega \subset \mathbb{R}^d$ of shape regular triangles K . By ‘‘macro finite elements’’ we mean that H , the size of the triangulation can be larger than the micro length scale ε .

REMARK 2.1. *Standard a-priori estimates for an H^2 -regular solution u^ε of problem (2.1) give $\|u^\varepsilon - u^H\|_{H^1(\Omega)} \leq C(H/\varepsilon)\|f\|_{L^2(\Omega)}$, (the factor $1/\varepsilon$ is due to the small oscillations in u^ε , see [23]). The goal is thus to define a numerical methods with convergence rate independent of ε .*

We consider for a macro triangle $K \in \mathcal{T}_h$ a sampling sub-domain centered at the barycenter x_K of K defined by $K_\delta = x_K + \delta[-1/2, 1/2]^d$, where $\delta \geq \varepsilon$ (see the last paragraph of this section). We define a pseudospectral mesh on K_δ based on the mesh defined in (2.12)

$$(2.14) \quad \begin{aligned} K_{\delta M} &:= (x_K + \delta(-1/2, 1/2)^d) + \delta Y_M \\ &= \{\xi_l = (\xi_{l_1}, \dots, \xi_{l_d}), 0 \leq l_i \leq 2M - 1, i = 1, \dots, d\}. \end{aligned}$$

We also define

$$(2.15) \quad S_M(K_\delta) := \text{span}\{e^{2i\pi kx/\varepsilon}; x \in K_\delta, k \in \mathbb{Z}^d, -M \leq k_i \leq M - 1\}/\mathbb{R},$$

where the quotient denotes the equivalence relation $u \simeq v \iff u - v$ is a constant. We will also use $S_M(Y)$ which is defined as (2.15) for the domain $Y = (0, 1)^d$ instead of K_δ . We define a bilinear form on $S_M(K_\delta)$ built on the pseudospectral mesh $K_{\delta M}$ given by

$$(2.16) \quad (u, v)_M := \frac{|K_\delta|}{(2M)^d} \sum_{l_1, \dots, l_d=0}^{2M-1} u(\xi_l) \bar{v}(\xi_l).$$

For functions $\mathbf{u}, \mathbf{v} \in S_M(K_\delta)$ we define

$$(2.17) \quad (\mathbf{u}, \mathbf{v})_M := \sum_{i=1}^d (u_i, v_i)_M.$$

We will use the same notation when working in the space $S_M(Y)$ with a pseudospectral mesh defined in (2.12) and a weight factor given by $\frac{|Y|}{(2M)^d} = \frac{1}{(2M)^d}$.

REMARK 2.2. *Notice that*

$$(2.18) \quad (u_M, v_M)_M = (u_M, v_M) \quad \forall u_M, v_M \in S_M(K_\varepsilon).$$

This follows from the fact that the integration formula $\int_0^1 v(x)dx \simeq \frac{1}{2M} \sum_{j=0}^{2M-1} v(x_j)$ is exact for $v \in S_{2M}(I)$ (see formula (A.2)).

The FES-HMM for the elliptic homogenization problems, based on the macro space $S_0^1(\Omega, \mathcal{T}_H)$, is defined by a *modified macro bilinear form*

$$(2.19) \quad B(u^H, v^H) = \sum_{K \in \mathcal{T}_H} \frac{|K|}{|K_\delta|} \int_{K_\delta} \nabla u_M a(x_K, x/\varepsilon) (\nabla v_M)^T dx,$$

where $K_\delta = x_K + \delta[-1/2, 1/2]^d$, is a sampling sub-domain centered at the barycenter x_K of K and $|K|, |K_\delta|$ denote the measure of K and K_δ , respectively. The (unknown) micro function u_M is the solution of the following *micro problem*: for $u^H \in S_0^1(\Omega, \mathcal{T}_H)$, find u_M such that $(u_M - u^H) = w_M \in S_M(K_\delta)$ and

$$(2.20) \quad (a(x_K, x/\varepsilon) \nabla w_M, \nabla z_M)_M = (a(x_K, x/\varepsilon) \nabla u^H, \nabla z_M)_M, \quad \forall z_M \in S_M(K_\delta).$$

The macro FES-HMM solution is defined by the following variational problem: find $u^H \in S_0^1(\Omega, \mathcal{T}_H)$ such that

$$(2.21) \quad B(u^H, v^H) = \int_\Omega f v^H dx := \langle f, v^H \rangle, \quad \forall v^H \in S_0^1(\Omega, \mathcal{T}_H).$$

REMARK 2.3. *Previous FE-HMM methods are based on FEM at the macro and the micro level [13],[14],[2],[3]. In these methods, the macro bilinear form is similar as (2.19), and the micro functions are such that $(u^h - u^H) \in S_{per}^1(K_\delta, \mathcal{T}_h)$ and*

$$(2.22) \quad \int_{K_\delta} \nabla u^h a(x_K, x/\varepsilon) (\nabla z^h)^T dx = 0 \quad \forall z^h \in S_{per}^1(K_\delta, \mathcal{T}_h),$$

where

$$(2.23) \quad S_{per}^1(K_\delta, \mathcal{T}_h) = \{z^h \in W_{per}^1(K_\delta); z^h|_T \in \mathcal{P}^1(T), T \in \mathcal{T}_h\},$$

and $\mathcal{P}^1(T)$ is the space of linear polynomials on the triangle T . Notice that setting $u^h - u^H = w^h \in S_{per}^1(K_\delta, \mathcal{T}_h)$, we can reformulate (2.22) as a periodic problem with right hand side involving u^H similarly as in (2.20).

Remark regarding the non-periodic case If the problem (2.1) is periodic with period ε one can chose $\delta = \varepsilon$ in the above method. There are many practical multiscale problems with periodic structures but even more without periodicity or where the period is not known. Semi-discrete convergence but not computational complexity of FE-HMM for these problems was studied in [14]. These problems typically require a computational micro-scale domain with a diameter δ that is larger than the period ε .

For problems with a periodic fast scale but with unknown period there are two error terms with implications for complexity: $\mathcal{O}(\delta) + \mathcal{O}(\varepsilon/\delta)$, see [14]. The first occurs if the coefficient is given as a^ε and not with explicitly defined dependence on the periodic oscillatory component and the second originates from a mismatch of the

boundary conditions. The $\mathcal{O}(\delta)$ term can actually be improved to $\mathcal{O}(\delta^q)$ for any given integer q by replacing the simple average used to approximate point wise values in [14] by a weighted average. The weight or kernel should satisfy certain moment and regularity conditions, see theorem 2.7 in [15]. The overall complexity can anyway not be bounded as well as with our periodic assumption. There are, however, cases for which only a narrow boundary layer is affected by the mismatch at the boundary of the microscale domain, see [19]. This is the reason why only a limited over sampling is often effective in the multiscale finite element method. We can then have $\delta = \mathcal{O}(\varepsilon)$ and the overall complexity will be of the same order as in the known periodic case. Notice that in the case that the explicit form of the tensor a^ε is not known or in the nonperiodic case, one should replace $S_M(K_\delta)$ defined in (2.15) by a micro space based on Chebyshev pseudospectral points. One still retains the fast convergence of the micro solutions, provided sufficient regularity of the problem.

2.4. Main results. After showing that (2.21) is well-posed we obtain, for periodic homogenization problems, the following convergence results, proved in Section 3 and extended in Section 4 for higher order macro FE spaces. Let u^0 be the solution of the homogenized problem (2.3), and assume u^0 is H^2 regular. Let u^H be the solution of problem (2.21). Provided sufficient regularity of the conductivity tensor $a^\varepsilon(x, x/\varepsilon)$, we have

Theorem (3.9).

$$(2.24) \quad \|u^0 - u^H\|_{H^1(\Omega)} \leq (C_1 H + C_2 M^{2(1-s)}),$$

where C_2 depends on $\|a_{ij}^\varepsilon\|_{L^\infty(\Omega; H_{per}^s(Y))}$, $\|\chi^j\|_{L^\infty(\Omega; H_{per}^s(Y))}$, where H is the size of the triangulation of the macro finite element space (2.13) and where M^d is the number of pseudospectral points of the micro space $S_M(K_\varepsilon)$ given in (2.15).

Corollary.

$$(2.25) \quad \|u^0 - u^H\|_{L^2(\Omega)} \leq (C_1 H^2 + C_2 M^{2(1-s)}) \|f\|_{L^2(\Omega)},$$

$$(2.26) \quad \|u^\varepsilon - u^H\|_{L^2(\Omega)} \leq (C_1 H^2 + C_2 M^{2(1-s)} + C_3 \varepsilon) \|f\|_{L^2(\Omega)}.$$

Provided analyticity of the conductivity tensor $a^\varepsilon(x, x/\varepsilon)$, we have

Theorem (3.11).

$$(2.27) \quad \|u^0 - u^H\|_{H^1(\Omega)} \leq \left(C_1 H + C_2 (M(\log M)^{(d-1)} e^{-\alpha M})^2 \right),$$

where C_2 depends on \mathcal{B} (see Definition 3.4), where H is the size of the triangulation of the macro finite element space (2.13) and where M^d is the number of pseudospectral points of the micro space $S_M(K_\varepsilon)$ given in (2.15).

Corollary.

$$(2.28) \quad \|u^0 - u^H\|_{L^2(\Omega)} \leq \left(C_1 H^2 + C_2 (M(\log M)^{(d-1)} e^{-\alpha M})^2 \right) \|f\|_{L^2(\Omega)},$$

$$(2.29) \quad \|u^\varepsilon - u^H\|_{L^2(\Omega)} \leq \left(C_1 H^2 + C_2 (M(\log M)^{(d-1)} e^{-\alpha M})^2 + C_3 \varepsilon \right) \|f\|_{L^2(\Omega)}.$$

Defining a numerical corrector u_p^ε (see 3.31) constrained by the known computed macro solution u^H , we obtain an approximation of the fine scale solution u^ε . Provided sufficient regularity of the conductivity tensor $a^\varepsilon(x, x/\varepsilon)$, we have

Theorem (3.13).

$$(2.30) \quad \|u^\varepsilon - u_p^\varepsilon\|_{H^1(\Omega)} \leq (C_1 H + C_2 M^{1-s} + C_3 \sqrt{\varepsilon}),$$

where C_2 depends on $\|a_{ij}^\varepsilon\|_{L^\infty(\Omega; H_{per}^s(Y))}$, $\|\chi^j\|_{L^\infty(\Omega; H_{per}^s(Y))}$, where H is the size of the triangulation of the macro finite element space (2.13) and where M^d is the number of pseudospectral points of the micro space $S_M(K_\varepsilon)$ given in (2.15).

The norm $\bar{H}^1(\Omega)$ (defined in 3.33) is a broken Sobolev norm which is needed since the reconstructed solution u_p^ε can be discontinuous across the macro elements K .

Provided analyticity of the conductivity tensor $a^\varepsilon(x, x/\varepsilon)$, we have

Theorem (3.14).

$$(2.31) \quad \|u^\varepsilon - u_p^\varepsilon\|_{\bar{H}^1(\Omega)} \leq (C_1 H + C_2 M (\log M)^{d-1} e^{-\alpha M} + C_3 \sqrt{\varepsilon}),$$

C_2 depends on \mathcal{B} (see Definition 3.4), where H is the size of the triangulation of the macro finite element space (2.13) and where M^d is the number of pseudospectral points of the micro space $S_M(K_\varepsilon)$ given in (2.15).

Under the stated assumptions, we see in Theorems and Corollaries 3.11 and 3.14 that the contribution of the error of the micro solution, $M(\log M)^{d-1} e^{-\alpha M}$, decreases exponentially fast. Going back to the notation of the introduction and denoting by N_{mac} the degrees of freedom of the macro FEM, we see that for solving elliptic homogenization problems in the L^2 norm with the usual quadratic convergence rate we have a total cost of $\mathcal{O}(N_{mac})$ (up to exponential convergence of the micro problem). The same is true for the H^1 norm and for the reconstructed solution. With the ‘‘standard’’ FE-HMM (see Remark 2.3) these costs are $\mathcal{O}(N_{mac}^2)$, $\mathcal{O}(N_{mac}^{3/2})$, $\mathcal{O}(N_{mac}^2)$, respectively [3],[5]. Thus, the proposed method achieves an *almost-linear* complexity (independent of ε) with respect to the number of macro degrees of freedom.

Higher order macro FEM. In Section 4 we will extend the numerical method for conforming higher order macro FE spaces of polynomials of degree $p \geq 1$ on quasiuniform meshes, provided u^0 and f , the right hand side of (2.3), are smooth enough ($u^0 \in H^{l+1}$ and $f \in H^l$ will suffice in what follows). For the original FE-HMM, using conforming FE spaces of polynomials of degree $p \geq 1$ for the macro solver while keeping piecewise linear FE space for the micro-solver one would obtain following [3] and [14]

$$(2.32) \quad \|u^0 - u^H\|_{H^1(\Omega)} \leq C(H^{\min(p,l)} + (h/\varepsilon)^2).$$

In the above estimate, $(h/\varepsilon)^2 = \mathcal{O}(N_{mic}^{-2/d})$ and $H^s = \mathcal{O}(N_{mac}^{-s/d})$, where $s = \min(p, l)$. Thus, not only the complexity is superlinear in the number of macro macro degrees of freedom N_{mac} , but for any $p, l \geq 1$ the global convergence rate cannot be faster than $\mathcal{O}(N_{mic}^{-2/d})$, (quadratic rate) unless higher order FEM are used for the micro problems. For the proposed approach, the same pseudo-spectral method, as defined in Section 2.3 can be used for the micro problems with higher order macro FEM. Provided sufficient regularity of the conductivity tensor $a^\varepsilon(x, x/\varepsilon)$, we obtain in Section 4

$$\|u^0 - u^H\|_{H^1(\Omega)} \leq C_1 H^{\min(p,l)} + C_2 M^{2(1-s)}.$$

Provided analytic conductivity tensor we obtain in Section 4 results of the form

$$\|u^0 - u^H\|_{H^1(\Omega)} \leq C_1 H^{\min(p,l)} + C_2 (M(\log M)^{d-1} e^{-\alpha M})^2,$$

i.e., up to exponentially decreasing micro error, an overall *almost linear* complexity of $\mathcal{O}(N_{mac})$ for a macro convergence rate of $H^{\min(p,l)}$. Convergence results for the L^2 norm will also be given for higher order macro FEM in Section 4.

3. Error analysis. We start by showing that the method is well posed and then prove the main convergence results. In the sequel, we set $\delta = \varepsilon$ in the FES-HMM defined in Section 2.3.

3.1. Well-posedness. We show in the following proposition that the bilinear form (2.19) is elliptic and bounded, thus (2.21) has a unique solution.

PROPOSITION 3.1. *The problem (2.21) has a unique solution which satisfies*

$$(3.1) \quad \|u^H\|_{H^1(\Omega)} \leq C\|f\|_{L^2(\Omega)}.$$

Proof. We first show that the problem (2.20) has a unique solution. The coercivity of the bilinear form defined in (2.20) follows from

$$\begin{aligned} (a(x_K, x/\varepsilon)\nabla w_M, \nabla w_M^T)_M &= \frac{|K_\varepsilon|}{(2M)^d} \sum_{j=0}^{2M-1} \nabla w_M(x_j) a(x_K, x_j/\varepsilon) \nabla w_M(x_j)^T \\ &\geq \gamma_1 \frac{|K_\varepsilon|}{(2M)^d} \sum_{j=0}^{2M-1} |\nabla w_M(x_j)|^2 = \gamma_1 \int_{K_\varepsilon} |\nabla w_M(x)|^2 dx = C\|w_M(x)\|_{H^1(K_\varepsilon)}^2, \end{aligned}$$

where we used the coercivity of $a(x, x/\varepsilon)$ (see (2.2)), the equivalence of discrete and continuous scalar products for functions in $S_M(K_\varepsilon)$ (see Remark 2.2) and the norm equivalence $\|\nabla w_M(x)\|_{L^2(K_\varepsilon)} \simeq \|w_M(x)\|_{H^1(K_\varepsilon)}$. We next show that the bilinear form defined in (2.20) is bounded

$$\begin{aligned} (a(x_K, x/\varepsilon)\nabla v_M, \nabla w_M^T)_M &\leq \gamma_2 \left(\frac{|K_\varepsilon|}{(2M)^d} \sum_{j=0}^{2M-1} \nabla v_M(x_j)^2 \right)^{1/2} \left(\frac{|K_\varepsilon|}{(2M)^d} \sum_{j=0}^{2M-1} \nabla w_M(x_j)^2 \right)^{1/2} \\ &\leq \gamma_2 \|\nabla v_M(x)\|_{L^2(K_\varepsilon)} \|\nabla w_M(x)\|_{L^2(K_\varepsilon)} \leq C\|v_M(x)\|_{H^1(K_\varepsilon)} \|w_M(x)\|_{H^1(K_\varepsilon)}, \end{aligned}$$

where we used the boundedness of $a(x, x/\varepsilon)$, the Cauchy-Schwarz inequality and as above, the equivalence of discrete and continuous scalar products and norms. Thus, the existence and uniqueness of a solution of problem (2.20) follows from the Lax-Milgram theorem.

It remains to show that (2.21) has a unique solution. Since $v_M - v^H \in S_M(K_\varepsilon) \subset W_{per}^1(K_\varepsilon)$ and ∇v^H is constant over a macro triangle K , we have

$$(3.2) \quad \int_{K_\varepsilon} |\nabla v_M|^2 dx = \int_{K_\varepsilon} |\nabla v_M - \nabla v^H|^2 dx + \int_{K_\varepsilon} |\nabla v^H|^2 dx,$$

$$(3.3) \quad \int_{K_\varepsilon} \nabla v_M a(x_K, x/\varepsilon) (\nabla v_M - \nabla v^H)^T dx = 0.$$

Equality (3.2) implies that $B(v^H, v^H) \geq C\|v^H\|_{H^1(K)}^2$ and thus the bilinear form B is coercive. Equality (3.3) implies that $\|\nabla v_M\|_{L^2(K_\varepsilon)} \leq C\|\nabla v^H\|_{L^2(K_\varepsilon)}$ and it follows that B is bounded. The existence and uniqueness of a solution u^H of problem (2.21) as well as (3.1) follow from the Lax-Milgram theorem. \square

The following representation for the solution w_M of problem (2.20) will be useful in the sequel.

LEMMA 3.2. *Let $w_M \in S_M(K_\varepsilon)$ be the solution of problem (2.20). Then*

$$(3.4) \quad w_M = \varepsilon \sum_{j=1}^d \chi_M^j(x_K, x/\varepsilon) \frac{\partial u^H}{\partial x_j},$$

where $\chi_M^j(x_K, y) \in S_M(Y)$, $j = 1, \dots, d$ are the solutions of the problem

$$(3.5) \quad (a(x_K, y) \nabla \chi_M^j, \nabla z_M)_M = (a(x_K, y) e_j, \nabla z_M)_M, \quad \forall z_M \in S_M(Y),$$

where $\{e_j\}_{j=1}^d$ is the standard basis of \mathbb{R}^d and where we set $y = x/\varepsilon$.

Proof. We know from Proposition 3.1 that Problems (2.20) and (3.5) have a unique solution. The lemma follows simply by inserting (3.4) in (2.20). \square

3.2. Convergence results. In this section, we estimate the convergence rate for the FES-HMM defined in section 2. In order to obtain convergence results for $\|u^0 - u^H\|$, where u^0 is the solution of the homogenized problem (2.3) and u^H is the solution of the FES-HMM defined in (2.21), we have to introduce the following semi-discrete bilinear form

$$(3.6) \quad \tilde{B}(u^H, v^H) = \sum_{K \in \mathcal{T}_H} \frac{|K|}{|K_\varepsilon|} \int_{K_\varepsilon} \nabla u \cdot a(x_K, x/\varepsilon) (\nabla v)^T dx,$$

where u (respectively v) is the solution of the micro problem (2.22) such that $(u - u^H) = w \in W_{per}^1(K_\varepsilon)$. It can be shown similarly as in Lemma 3.2 that (see also [3] for details)

$$(3.7) \quad w = \varepsilon \sum_{j=1}^d \chi^j(x_K, x/\varepsilon) \frac{\partial u^H}{\partial x_j},$$

where $\chi^j(x_K, y) \in W_{per}^1(Y)$ $j = 1, \dots, d$ are the solutions of the problem

$$(3.8) \quad \int_Y \nabla \chi^j a(x_K, y) (\nabla z)^T dx = \int_{K_\varepsilon} e_j^T a(x_K, y) (\nabla z)^T dx, \quad \forall z \in W_{per}^1(Y),$$

where $\{e_j\}_{j=1}^d$ is the standard basis of \mathbb{R}^d and where we set as usual $y = x/\varepsilon$. We define a semi-discrete macro solution by \tilde{u}^H , solution of the problem: find $\tilde{u}^H \in S_0^1(\Omega, \mathcal{T}_H)$ such that

$$(3.9) \quad \tilde{B}(\tilde{u}^H, v^H) = \langle f, v^H \rangle, \quad \forall v^H \in S_0^1(\Omega, \mathcal{T}_H).$$

Similarly as for Problem 2.21, it can be shown that this problem has a unique solution which satisfies $\|\tilde{u}^H\|_{H^1(\Omega)} \leq C \|f\|_{L^2(\Omega)}$.

Following [3], the error estimates for the FES-HMM can be obtain as sketched below. Let u^H, \tilde{u}^H be the solution of problems 2.21 and 3.9, respectively. Let u^0 be the solution of the homogenized problem (2.3). Then,

$$(3.10) \quad \|u^0 - u^H\|_{H^1(\Omega)} \leq CH \|f\|_{L^2(\Omega)} + \|\tilde{u}^H - u^H\|_{H^1(\Omega)},$$

where we used the triangle inequality and the estimation $\|u^0 - \tilde{u}^H\|_{H^1(\Omega)} \leq CH \|f\|_{L^2(\Omega)}$ (see [3]). The second term of the right-hand side of the inequality can be estimated in the following way. Denoting by $v^H = \tilde{u}^H - u^H$, we have

$$(3.11) \quad \alpha \|\tilde{u}^H - u^H\|_{H^1(\Omega)}^2 \leq \tilde{B}(\tilde{u}^H - u^H, v^H) = B(u^H, v^H) - \tilde{B}(u^H, v^H)$$

and thus

$$(3.12) \quad \|\tilde{u}^H - u^H\|_{H^1(\Omega)} \leq \frac{1}{\alpha} \frac{|B(u^H, v^H) - \tilde{B}(u^H, v^H)|}{\|v^H\|_{H^1(\Omega)}}.$$

This is the basis of our convergence analysis. For sufficiently smooth tensor $a(x, x/\varepsilon)$, we prove in the sequel that the right-hand side of (3.12) is spectrally decaying, i.e., decays faster than any fixed degree polynomial rate. Provided analyticity of $a^\varepsilon(x)$, we prove that the right-hand side of (3.12) is exponentially decaying. L^2 convergence rates will be derived as well.

3.3. Spectral and exponential convergence. We start with a lemma needed to derive the spectral accuracy of the method (see Appendix A for a proof).

LEMMA 3.3. *Assume $w(y) \in H_{per}^s(Y)$, with $s \geq 2$, then*

$$(3.13) \quad \|w(y) - \mathbf{Q}_M w(y)\|_{L^2(Y)} \leq CM^{-s} \|w\|_{H^s(Y)}$$

$$(3.14) \quad \|\nabla w(y) - \nabla \mathbf{Q}_M(w(y))\|_{L^2(Y)} \leq CM^{1-s} \|w\|_{H^s(Y)}$$

where for a function $w \in L_{per}^2(Y)$, the spectral interpolant $\mathbf{Q}_M w$ is defined in (2.11) and M^d is the number of pseudospectral points of the micro space $S_M(Y)$ (see (2.15)).

For the proof of the exponential decay estimate, we first need some notations.

DEFINITION 3.4. *Let $I = (0, 1)$. The set $\mathcal{A}_{B, \alpha}(I)$ consists of functions $g(y) \in C^\infty(\mathbb{R})$ I -periodic and having an analytic extension in the strip*

$$(3.15) \quad \mathcal{E}_\alpha := \{z \in \mathbb{C}; |Im(z)| \leq \alpha\}.$$

We define $B := \max_{z \in \mathcal{E}_\alpha} |g(z)|$. For a multivariate function $g(y) \in C^\infty(\mathbb{R})$ Y -periodic, where $Y = (0, 1)^d$, we say that $g(y)$ belongs to $\mathcal{A}_{B, \alpha}(Y)$ if for all $i = 1, \dots, d$ and each $Y_{-i} = (y_1, \dots, y_{i-1}, y_{i+1}, \dots, y_d) \in (0, 1)^{d-1}$, $g_i(y_i) := g(Y_{-i}, y_i)$ has an analytic extension in the strip \mathcal{E}_α . We denote by $\mathcal{B} := \max_{i \leq d} \{\max_{Y_{-i}} \{\max_{z \in \mathcal{E}_\alpha} |g(Y_{-i}, z)|\}\}$.

REMARK 3.5. *It is known that if the coefficients of an elliptic operator are real analytic in the closure of the domain of consideration \bar{D} and if the right-hand side is real analytic, then the solution of the corresponding elliptic equation is a real analytic function in \bar{D} . We refer the reader to [27],[9] for a precise discussion and proof of the above statement. For the problem (3.8), we note that the coefficients of the elliptic operator and the right-hand side are given by the same functions $a_{ij}(x_K, \cdot)$. Assuming $a_{ij}(x_K, \cdot) \in \mathcal{A}_{B, \alpha}(Y) \quad \forall i, j = 1, \dots, d$ it follows from the above consideration that the solutions $\chi^j(x_K, y)$ of (3.8) are also analytic for all $j = 1, \dots, d$.*

The following lemma is proved in the Appendix A.

LEMMA 3.6. *Let $w \in \mathcal{A}_{B, \alpha}(Y)$ Then*

$$(3.16) \quad \|w(y) - \mathbf{Q}_M w(y)\|_{L^2(Y)} \leq CB(\log M)^{d-1} e^{-\alpha M}$$

$$(3.17) \quad \|\nabla w(y) - \nabla \mathbf{Q}_M(w(y))\|_{L^2(Y)} \leq CBM(\log M)^{d-1} e^{-\alpha M}$$

where for a function $w \in L_{per}^2(Y)$, the spectral interpolant $\mathbf{Q}_M w$ is defined in (2.11) and M^d is the number of pseudospectral points of the micro space $S_M(Y)$ (see (2.15)).

Let $a^\varepsilon = (a_{ij}(x_K, x/\varepsilon))_{i,j=1}^d$ be the bounded coercive tensor of problem (2.1). We set $a_{ij}(x_K, x/\varepsilon) = a_{ij}(x_K, y)$. In view of (3.10) and (3.12), we prove the main results of this section, leading to spectral and exponential convergence of the micro solver.

LEMMA 3.7. *Assume that $a_{ij}(x, y), X^j(x, y) \in L^\infty(\Omega; H_{per}^s(Y)) \quad \forall i, j = 1, \dots, d$ with $s > 2$. Then, for $u^H, v^H \in S_0^1(\Omega, \mathcal{T}_H)$*

$$(3.18) \quad \left| B(u^H, v^H) - \tilde{B}(u^H, v^H) \right| \leq C(M^{1-s})^2 \|\nabla u^H\|_{L^2(\Omega)} \|\nabla v^H\|_{L^2(\Omega)},$$

where C depends on $\|a_{ij}\|_{L^\infty(\Omega; H_{per}^s(Y))}, \|\chi^j\|_{L^\infty(\Omega; H_{per}^s(Y))}$, where the bilinear forms $B(\cdot, \cdot), \tilde{B}(\cdot, \cdot)$ are defined in (2.19) and (3.6), respectively, and where M^d is the number of pseudospectral points of the micro space $S_M(K_\varepsilon)$ given in (2.15).

LEMMA 3.8. *Assume that $\chi^j(x, \cdot), a_{i,j}(x, \cdot) \in \mathcal{A}_{\mathcal{B}, \alpha}(Y) \forall i, j = 1, \dots, d$ and a.e. $x \in \Omega$. Then, for $u^H, v^H \in S_0^1(\Omega, \mathcal{T}_H)$*

$$(3.19) \quad \left| B(u^H, v^H) - \tilde{B}(u^H, v^H) \right| \leq C (M(\log M)^{(d-1)} e^{-\alpha M})^2 \|\nabla u^H\|_{L^2(\Omega)} \|\nabla v^H\|_{L^2(\Omega)},$$

where the constant C depends on \mathcal{B} , where the bilinear forms $B(\cdot, \cdot), \tilde{B}(\cdot, \cdot)$ are defined in (2.19) and (3.6), respectively, and where M^d is the number of pseudospectral points of the micro space $S_M(K_\varepsilon)$ given in (2.15).

The proof of both Lemmas follows the same lines. We prove Lemma 3.8 and indicate after the proof the modifications which lead to the proof of Lemma 3.7.

Proof. [Lemma 3.8] Using the definitions of (2.19) and (3.6) following Lemma 3.3 of [3] we have

$$(3.20) \quad \begin{aligned} & |B(u^H, v^H) - \tilde{B}(u^H, v^H)| = \\ & \left| \sum_{K \in \mathcal{T}} \frac{|K|}{|K_\varepsilon|} \left(\int_{K_\varepsilon} \nabla u a(x_K, x/\varepsilon) (\nabla v)^T dx - \int_{K_\varepsilon} \nabla u_M a(x_K, x/\varepsilon) (\nabla v_M)^T dx \right) \right| = \\ & \left| \sum_{K \in \mathcal{T}} \frac{|K|}{|K_\varepsilon|} \left(\int_{K_\varepsilon} \nabla(u - u_M) a(x_K, x/\varepsilon) (\nabla v)^T dx - \right. \right. \\ & \left. \left. \int_{K_\varepsilon} \nabla u_M a(x_K, x/\varepsilon) (\nabla(v_M - v))^T dx \right) \right|. \end{aligned}$$

We observe that the first member of the last line of (3.20) is zero since $(u - v_M) \in W_{per}^1(K_\varepsilon)$. Using the same argument and replacing u_M by $u_M - u$ in the second expression we have

$$(3.21) \quad \left| \sum_{K \in \mathcal{T}} \frac{|K|}{|K_\varepsilon|} \int_{K_\varepsilon} \nabla(u_M - u) a(x_K, x/\varepsilon) (\nabla(v_M - v))^T dx \right| \leq$$

$$(3.22) \quad C \sum_{K \in \mathcal{T}} \frac{|K|}{|K_\varepsilon|} \|\nabla u_M - \nabla u\|_{L^2(K_\varepsilon)} \|\nabla v_M - \nabla v\|_{L^2(K_\varepsilon)},$$

where we used that the bilinear form is bounded. We have next to estimate

$$(3.23) \quad \begin{aligned} \|\nabla u_M - \nabla u\|_{L^2(K_\varepsilon)} &= \left\| \varepsilon \sum_{j=1}^d \nabla(\chi_M^j(x_K, x/\varepsilon) - \chi^j(x_K, x/\varepsilon)) \frac{\partial u^H}{\partial x_j} \right\|_{L^2(K_\varepsilon)} \\ &= \sqrt{|K_\varepsilon|} \left\| \sum_{j=1}^d \nabla(\chi_M^j(x_K, y) - \chi^j(x_K, y)) \frac{\partial u^H}{\partial x_j} \right\|_{L^2(Y)} \\ &\leq C \sqrt{|K_\varepsilon|} \|\nabla u^H\|_{L^2(Y)} \max_j \|\nabla(\chi_M^j - \chi^j)\|_{L^2(Y)}. \end{aligned}$$

It remains to estimate $\|\nabla(\chi_M^{j_m} - \chi^{j_m})\|_{L^2(Y)}$, where j_m is the index corresponding to the maximum in the above expression. In view of (3.5) and (3.8) using Strang lemma we get

$$(3.24) \quad \begin{aligned} \|\nabla(\chi_M^{j_m} - \chi^{j_m})\|_{L^2(Y)} &\leq C \left(\inf_{z_M \in S_M(Y)} \left(\|\nabla(\chi^{j_m} - z_M)\|_{L^2(Y)} \right. \right. \\ & \left. \left. + \sup_{w_M \in S_M(Y)} \frac{|(a(x_K, y)(\nabla z_M)^T, \nabla w_M) - (a(x_K, y)(\nabla z_M)^T, \nabla w_M)_M|}{\|\nabla w_M\|_{L^2(Y)}} \right) \right) \end{aligned}$$

$$+ \sup_{w_M \in S_M(Y)} \frac{|(a(x_K, y)e_{j_m}, \nabla w_M) - (a(x_K, y)e_{j_m}, \nabla w_M)_M|}{\|\nabla w_M\|_{L^2(Y)}}.$$

We chose $z_M = \chi_M^{j_m}$ for the infimum in (3.24) and using Lemma 3.6, we obtain

$$\|\nabla(\chi^{j_m} - \chi_M^{j_m})\|_{L^2(Y)} \leq C\mathcal{B}M(\log M)^{d-1}e^{-\alpha M}.$$

For the second term of the right-hand side of (3.24) (with $z_M = \chi_M^{j_m}$) we have

$$\begin{aligned} & \left| (a(x_K, y)(\nabla \chi_M^{j_m})^T, \nabla w_M) - (a(x_K, y)(\nabla \chi_M^{j_m})^T, \nabla w_M)_M \right| \leq \\ & \left| \left((a(x_K, y) - \mathbf{Q}_M(a(x_K, y))) (\nabla \chi_M^{j_m})^T, \nabla w_M \right) \right| + \left| \left((a(x_K, y) - \mathbf{Q}_M(a(x_K, y))) (\nabla \chi_M^{j_m})^T, \nabla w_M \right)_M \right| \end{aligned}$$

where we used Remark 2.2. The second term is zero and the first term is bounded by

$$\begin{aligned} & \left| \left((a(x_K, y) - \mathbf{Q}_M(a(x_K, y))) (\nabla \chi_M^{j_m})^T, \nabla w_M \right) \right| \\ & \leq \|\nabla \chi_M^{j_m}\|_{L^\infty(Y)} \int_Y \sum_{l,m=1}^d \left| (a_{l,m}(x_K, y) - \mathbf{Q}_M(a_{l,m}(x_K, y))) \partial_{y_l} w_M \right| dy \\ & \leq \|\nabla \chi_M^{j_m}\|_{L^\infty(Y)} \max_{l,m} \|a_{l,m}(x_K, y) - \mathbf{Q}_M(a_{l,m}(x_K, y))\|_{L^2(Y)} \|\nabla w_M\|_{L^2(Y)} \\ & \leq C\mathcal{B}(\log M)^{d-1}e^{-\alpha M} \|\nabla w_M\|_{L^2(Y)} \end{aligned}$$

using the assumptions and Lemma 3.6. For the third term of the right-hand side of (3.24) we write $g_{j_m} = a(x_K, y)e_{j_m}$ and $\mathbf{Q}_M g_{j_m}$ the spectral interpolant of g_{j_m} in $S_M(Y)$. Using Remark 2.2 and Lemma 3.6 we obtain

$$\begin{aligned} & |(g_{j_m}, \nabla w_M) - (g_{j_m}, \nabla w_M)_M| = |(g_{j_m} - \mathbf{Q}_M(g_{j_m}), \nabla w_M)| \\ & \leq \|g_{j_m} - \mathbf{Q}_M(g_{j_m})\|_{L^2(Y)} \|\nabla w_M\|_{L^2(Y)} \leq C\mathcal{B}e^{-\alpha M} \|\nabla w_M\|_{L^2(Y)}. \end{aligned}$$

using Lemma 3.6. Summing all the terms together and using that ∇u^H is constant, we obtain

$$\begin{aligned} & \sqrt{|K|/|K_\varepsilon|} \|\nabla u_M - \nabla u\|_{L^2(K_\varepsilon)} \leq C\mathcal{B}M(\log M)^{d-1}e^{-\alpha M} \sqrt{|K|} \|\nabla u^H\|_{L^2(Y)} \\ & \leq C\mathcal{B}M(\log M)^{d-1}e^{-\alpha M} \|\nabla u^H\|_{L^2(K)}. \end{aligned}$$

The second term of (3.22), $\|\nabla v_M - \nabla v\|_{L^2(K_\varepsilon)}$, is treated similarly. Finally summing up over $K \in \mathcal{T}$ we find that (3.22) is bounded by

$$C (\mathcal{B}M(\log M)^{d-1}e^{-\alpha M})^2 \|\nabla v^H\|_{L^2(\Omega)} \|\nabla w^H\|_{L^2(\Omega)},$$

and the proof is complete. \square

Proof. [Lemma 3.7] The first part of the proof until inequality (3.24) is identical as for Lemma 3.8. We next discuss the estimation of the three terms on the right hand side of the inequality (3.24). We chose $z_M = \chi_M^{j_m}$ for the infimum in (3.24) and the first term can be bounded using Lemma 3.3 by

$$\|\nabla(\chi^{j_m} - \chi_M^{j_m})\|_{L^2(Y)} \leq CM^{1-s} \|\chi^{j_m}(x_K, y)\|_{H^s(Y)}.$$

For the second term on the right-hand side of (3.24) (with $z_M = \chi_M^{j_m}$) Lemma 3.3 leads to

$$\begin{aligned} & \left| (a(x_K, y)(\nabla \chi_M^{j_m})^T, \nabla w_M) - (a(x_K, y)(\nabla \chi_M^{j_m})^T, \nabla w_M)_M \right| \\ & \leq CM^{1-s} \max_{l,m} \|a_{l,m}(x_K, y)\|_{H^s(Y)} \|\nabla \chi_M^{j_m}(x_K, y)\|_{L^\infty(Y)} \|\nabla w_M\|_{L^2(Y)}, \end{aligned}$$

which is obtained similarly as in Lemma 3.8. Notice that since $\chi^j(x, y) \in L^\infty(\Omega; H_{per}^s(Y))$ with $s > 2$, we have that $\chi^j(x, y) \in L^\infty(\Omega; W^{1,\infty}(Y))$ (Sobolev's inequality). For the third term of the right-hand side of (3.24) we write $g_{j_m} = a(x_K, y)e_{j_m}$ and $\mathbf{Q}_M g_{j_m}$ the interpolant of g_{j_m} in $S_M(Y)$. Similarly as in Lemma 3.8 and using Lemma 3.3 we obtain

$$\begin{aligned} & |(g_{j_m}, \nabla w_M) - (g_{j_m}, \nabla w_M)_M| = |(g_{j_m} - \mathbf{Q}_M(g_{j_m}), \nabla w_M)| \\ & \leq CM^{-s} \max_{l,m} \|a_{l,m}(x_K, y)\|_{H^s(Y)} \|\nabla w_M\|_{L^2(Y)}. \end{aligned}$$

Summing all the terms together and using that ∇u^H is constant, we obtain

$$\sqrt{|K|/|K_\varepsilon|} \|\nabla u_M - \nabla u\|_{L^2(K_\varepsilon)} \leq M^{1-s} \|\nabla u^H\|_{L^2(Y)}$$

Using similar argument for the second term of (3.22), summing over $K \in \mathcal{T}$ and using that $a_{ij}(x, y), \chi^j(x, y) \in L^\infty(\Omega; H_{per}^s(Y))$ we find that (3.22) is bounded by

$$C (M^{1-s})^2 \|\nabla u^H\|_{L^2(\Omega)} \|\nabla v^H\|_{L^2(\Omega)},$$

and the proof is complete. \square

3.4. Error estimates for the FES-HMM. We can now give the error estimate between the homogenized solution and the solution of the FES-HMM.

THEOREM 3.9. *Let u^0 be the solution of the homogenized problem (2.3), and assume u^0 is H^2 -regular. Let u^H be the solution of problem (2.21) and suppose that the assumptions of Lemma 3.3 hold. Then*

$$(3.25) \quad \|u^0 - u^H\|_{H^1(\Omega)} \leq (C_1 H + C_2 M^{2(1-s)}) \|f\|_{L^2(\Omega)},$$

where C_2 depends on $\|a_{ij}\|_{L^\infty(\Omega; H_{per}^s(Y))}, \|\chi^j\|_{L^\infty(\Omega; H_{per}^s(Y))}$, where H is the size of the triangulation of the macro finite element space (2.13) and where M^d is the number of pseudospectral points of the micro space $S_M(K_\varepsilon)$ given in (2.15).

Proof. Using (3.10), (3.12) and Lemma 3.3 gives the result. \square

COROLLARY 3.10. *Suppose that the assumptions of Theorem 3.9 hold. Then we have the following estimates*

$$(3.26) \quad \|u^0 - u^H\|_{L^2(\Omega)} \leq (C_1 H^2 + C_2 M^{2(1-s)}) \|f\|_{L^2(\Omega)},$$

$$(3.27) \quad \|u^\varepsilon - u^H\|_{L^2(\Omega)} \leq (C_1 H^2 + C_2 M^{2(1-s)} + C_3 \varepsilon) \|f\|_{L^2(\Omega)}.$$

Proof. Estimation (3.26) follows from the estimation $\|u^0 - \tilde{u}^H\|_{L^2(\Omega)} \leq CH^2 \|f\|_{L^2(\Omega)}$ for the solution of (3.9) (see [2, App. A]), the estimation for (3.12) obtained in Lemma 3.3 and the triangle inequality. Estimation (3.27) follows from (2.5), (3.26) and the triangle inequality. \square

Using Lemma 3.6 instead of Lemma 3.3 we obtain

THEOREM 3.11. *Let u^0 be the solution of the homogenized problem (2.3), and assume u^0 is H^2 -regular. Let u^H be the solution of problem (2.21) and suppose that the assumptions of Lemma 3.6 hold. Then*

$$(3.28) \quad \|u^0 - u^H\|_{H^1(\Omega)} \leq \left(C_1 H + C_2 (M(\log M)^{(d-1)} e^{-\alpha M})^2 \right) \|f\|_{L^2(\Omega)},$$

where C_2 depends on \mathcal{B} , where H is the size of the triangulation of the macro finite element space (2.13) and where M^d is the number of pseudospectral points of the micro space $S_M(K_\varepsilon)$ given in (2.15).

COROLLARY 3.12. *Suppose that the assumptions of Theorem 3.11 hold. Then we have the following estimates*

$$(3.29) \quad \|u^0 - u^H\|_{L^2(\Omega)} \leq \left(C_1 H^2 + C_2 (M(\log M)^{(d-1)} e^{-\alpha M})^2 \right) \|f\|_{L^2(\Omega)},$$

$$(3.30) \quad \|u^\varepsilon - u^H\|_{L^2(\Omega)} \leq \left(C_1 H^2 + C_2 (M(\log M)^{(d-1)} e^{-\alpha M})^2 + C_3 \varepsilon \right) \|f\|_{L^2(\Omega)}.$$

3.5. Retrieving microscopic information. So far we gave a numerical procedure to approximate the macro (homogenized) solution of the problem (2.3). Following [28] (see also [14],[3]) we consider a procedure to retrieve the microscopic information in order to approximate the microscopic solution of problem (2.1). We define u_p^ε by

$$(3.31) \quad u_p^\varepsilon(x)|_K = u^H(x) + (u_M(x) - u^H(x))|_K^P \text{ for } x \in K \in \mathcal{T}_H,$$

where $|_K^P$ denotes the periodic extension of the fine scale solution $(u_M - u^H)$, available in K_ε , on each element K . This extension is defined for a function $w \in H^1(K_\varepsilon)$ or in $\bar{S}_M(K_\varepsilon)$ by

$$w_p(x + \varepsilon j) = w(x) \quad \forall j = (j_1, \dots, j_d) \in \mathbb{Z}^d, \quad \forall x \in K_\varepsilon \quad \text{such that } x + \varepsilon j \in K.$$

The function u_p^ε which can be expressed by

$$(3.32) \quad u_p^\varepsilon = u^H(x) + \varepsilon \sum_{j=1}^d \chi_M^j(x_K, x/\varepsilon)|_K^P \frac{\partial u^H}{\partial x_j}$$

using (3.4) for the whole macro element K , defines a fully discrete fine scale approximation of the solution u^ε of problem (2.1). Since u_p^ε can be discontinuous across the macro elements K , we define a broken H^1 norm by

$$(3.33) \quad \|u\|_{\bar{H}^1(\Omega)} := \left(\sum_{K \in \mathcal{T}_H} \|\nabla u\|_{L^2(K)}^2 \right)^{1/2}.$$

Similarly as in (3.32), we define

$$(3.34) \quad \tilde{u}_p^\varepsilon = \tilde{u}^H(x) + \varepsilon \sum_{j=1}^d \chi^j(x_K, x/\varepsilon)|_K^P \frac{\partial \tilde{u}^H}{\partial x_j},$$

the reconstructed semi-discrete function based on the formula (3.7), where \tilde{u}^H is the solution of problem (3.9). In the sequel we will skip the notation $|_K^P$ for the functions

defined in (3.32) and (3.34). We obtain a error estimate of the fully discrete fine scale solution (3.31) in the following way

$$(3.35) \quad \|u^\varepsilon - u_p^\varepsilon\|_{\bar{H}^1(\Omega)} \leq \|u^\varepsilon - \tilde{u}_p^\varepsilon\|_{\bar{H}^1(\Omega)} + \|\tilde{u}_p^\varepsilon - u_p^\varepsilon\|_{\bar{H}^1(\Omega)}.$$

The first term of the right-hand side of the above inequality can be bounded by $\|u^\varepsilon - \tilde{u}_p^\varepsilon\|_{\bar{H}^1(\Omega)} \leq C(\sqrt{\varepsilon} + H)\|f\|_{L^2(\Omega)}$ (see [14],[3]) and it remains to estimate the second term. We have

$$\begin{aligned} \sum_{K \in \mathcal{T}_H} \|\nabla(\tilde{u}_p^\varepsilon - u_p^\varepsilon)\|_{L^2(K)}^2 &\leq \sum_{K \in \mathcal{T}_H} \|\nabla(\tilde{u}^H - u^H)\|_{L^2(K)}^2 + \\ \sum_{K \in \mathcal{T}_H} \left\| \sum_{j=1}^n \nabla(\varepsilon \chi^j(x_K, x/\varepsilon)) \left(\frac{\partial \tilde{u}^H}{\partial x_j} - \frac{\partial u^H}{\partial x_j} \right) \right\|_{L^2(K)}^2 &+ \\ \sum_{K \in \mathcal{T}_H} \left\| \sum_{j=1}^n \nabla(\varepsilon(\chi^j - \chi_M^j)(x_K, x/\varepsilon)) \frac{\partial u^H}{\partial x_j} \right\|_{L^2(K)}^2. & \end{aligned}$$

The first two terms are bounded by $C M^2(\log M)^{2(d-1)} e^{-2\alpha M} \|f\|_{L^2(\Omega)}^2$ if the assumptions of Lemma 3.8 hold and by $C M^{2(1-s)} \|f\|_{L^2(\Omega)}^2$ if the assumptions of Lemma 3.7 hold. For the last term, we take a finite covering of $K \subset \bigcup_{x_l \in K} K_\varepsilon(x_l)$, where $K_\varepsilon(x_l) = x_l + \varepsilon[-1/2, 1/2]^d$ and bound it by $C M^2(\log M)^{2(d-1)} e^{-2\alpha M} \|\nabla u^H\|_{L^2(\Omega)}^2$ using Lemma 3.6 or by $C M^{2(1-s)} \|\nabla u^H\|_{L^2(\Omega)}^2$ with Lemma 3.3.

Using the norm defined in (3.33) and (3.1) we obtain

$$\|\tilde{u}_p^\varepsilon - u_p^\varepsilon\|_{\bar{H}^1(\Omega)} \leq C M(\log M)^{d-1} e^{-\alpha M} \|f\|_{L^2(\Omega)} \text{ or } \|u_p^\varepsilon - u_p^\varepsilon\|_{\bar{H}^1(\Omega)} \leq C M^{1-s} \|f\|_{L^2(\Omega)},$$

if the assumptions of Lemma 3.6 or Lemma 3.3 hold, respectively. We have proved

THEOREM 3.13. *Let u_p^ε be defined by (3.31) and u^ε be the solution of (2.1). Suppose that the assumptions of Lemma 3.3 hold. Then*

$$(3.36) \quad \|u^\varepsilon - u_p^\varepsilon\|_{\bar{H}^1(\Omega)} \leq (C_1 H + C_2 M^{1-s} + C_3 \sqrt{\varepsilon}) \|f\|_{L^2(\Omega)},$$

where C_2 depends on $\|a_{ij}\|_{L^\infty(\Omega; H_{per}^s(Y))}$, $\|\chi^j\|_{L^\infty(\Omega; H_{per}^s(Y))}$, where H is the size of the triangulation of the macro finite element space (2.13), and where M^d is the number of pseudospectral points of the micro space $S_M(K_\varepsilon)$ given in (2.15).

THEOREM 3.14. *Let u_p^ε be defined by (3.31) and u^ε be the solution of (2.1). Suppose that the assumptions of Lemma 3.6 hold. Then*

$$(3.37) \quad \|u^\varepsilon - u_p^\varepsilon\|_{\bar{H}^1(\Omega)} \leq (C_1 H + C_2 M(\log M)^{d-1} e^{-\alpha M} + C_3 \sqrt{\varepsilon}) \|f\|_{L^2(\Omega)},$$

where C_2 depends on \mathcal{B} , where H is the size of the triangulation of the macro finite element space (2.13) and where M^d is the number of pseudospectral points of the micro space $S_M(K_\varepsilon)$ given in (2.15).

4. Higher order macroscopic solver. In this section we discuss higher order multiscale methods. We consider a macro finite element space defined by

$$(4.1) \quad S_0^p(\Omega, \mathcal{T}_H) = \{u^H \in H_0^1(\Omega); u^H|_K \in \mathcal{P}^p(K), \forall K \in \mathcal{T}_H\},$$

where $\mathcal{P}^p(K)$ is the space of polynomials of degree $p \geq 1$ on the triangle K , and \mathcal{T}_H is a quasi-uniform triangulation of $\Omega \subset \mathbb{R}^d$ of shape regular triangles K . We further

assume that the solution of the homogenized problem (2.3) satisfies $u^0 \in H^{l+1}(\Omega)$ and that the right hand side satisfies $f \in H^l(\Omega)$.

We also consider a p th-order numerical quadrature scheme (ω_i, x_i) $i = 1, \dots, L$ on K , with $\omega_i > 0 \forall i$ such that ¹

$$(4.2) \quad \frac{1}{|K|} \int_K p(x) dx = \sum_{i=1}^L \omega_i p(x_i) \text{ for all } p(x) \in \mathcal{P}^{2p-2}(K).$$

For $u^H \in S_0^p(\Omega, \mathcal{T}_H)$ we define its linear approximation at a quadrature point x_i by

$$(4.3) \quad u_{lin}^H := u^H(x_i) + \nabla u^H(x_i) \cdot (x - x_i).$$

Following [14], we define a bilinear form with help of the quadrature formula by

$$(4.4) \quad B(u^H, v^H) = \sum_{K \in \mathcal{T}_H} \sum_{i=1}^L \omega_i \int_{K_{\varepsilon,i}} \nabla u_{lin,M} a(x_i, x/\varepsilon) (\nabla v_{lin,M})^T dx,$$

where $K_{\varepsilon,i} = x_i + \varepsilon[-1/2, 1/2]^d \subset K$ is a sampling sub-domain centered at the quadrature point x_i and $u_{lin,M}$ is such that $(u_{lin,M} - u_{lin}^H) = w_M \in S_M(K_\varepsilon)$ and w_M the solution of the micro problem defined in (2.20).

The macro solution of the FES-HMM is then given by the following variational problem: find $u^H \in S_0^p(\Omega, \mathcal{T}_H)$ such that

$$(4.5) \quad B(u^H, v^H) = \langle f, v^H \rangle, \quad \forall v^H \in S_0^p(\Omega, \mathcal{T}_H).$$

Following the line of Proposition 3.1 shows that the problem (4.5) is well-posed. Notice that the assumption on the quadrature formula (4.2) are needed to ensure the coercivity of the bilinear form (4.4) (see [12, Theorem 4.1.2]).

As in Section 3, we consider the semi-discrete bilinear form $\tilde{B}(\cdot, \cdot)$ similar to (4.4) but with micro solution $(u - u_{lin}^H) = w \in W_{per}^1(K_\varepsilon)$ and the semi-discrete solution $\tilde{u}^H \in S_0^l(\Omega, \mathcal{T}_H)$ of the corresponding problem (3.9). Let u^0 be the solution of the homogenized problem (2.3). Then,

$$(4.6) \quad \|u^0 - u^H\|_{H^1(\Omega)} \leq \|u^0 - \tilde{u}\|_{H^1(\Omega)} + \|\tilde{u} - u^H\|_{H^1(\Omega)}.$$

It has been shown in [14] that, provided $u^0 \in H^{l+1}(\Omega)$ and $f \in H^{l+1}(\Omega)$ and provided $2l > d$ that $\|u^0 - \tilde{u}\|_{H^1(\Omega)} \leq C(H^{\min(p,l)} + \varepsilon)$ (see also [12, Thm. 4.1.6] for a discussion on the regularity assumptions and the need of the assumption $2l > d$).

REMARK 4.1. *The above result has been shown for a modified bilinear form similar to (4.4), but where the tensor $a(x, x/\varepsilon)$ is not collocated at the integration points. For the bilinear form (4.4), following [2, App. A] and the proof in [14] in can be shown that*

$$(4.7) \quad \|u^0 - \tilde{u}\|_{H^1(\Omega)} \leq CH^{\min(p,l)}.$$

The remaining term $\|\tilde{u} - u^H\|_{H^1(\Omega)}$ can be estimated following the proof of Lemmata 3.8 and 3.7. Furthermore, we can define a reconstructed solution similarly as in (3.31). We summarize our discussion.

¹We assume of course, that the quadrature points are of type ‘‘PI’’ (positive interior), i.e. that $\omega_i > 0$, $x_i \in K$, $\forall i$.

THEOREM 4.2. *Let u^0 be the solution of the homogenized problem (2.3), and assume $u^0 \in H^{l+1}(\Omega)$ and $f \in H^l(\Omega)$, $2l > d$. Let u^H be the solution of problem (4.5) and suppose that the assumptions of Lemma 3.3 hold. Then*

$$(4.8) \quad \|u^0 - u^H\|_{H^1(\Omega)} \leq C_1 H^{\min(p,l)} + C_2 M^{2(1-s)},$$

$$(4.9) \quad \|u^0 - u^H\|_{L^2(\Omega)} \leq C_1 H^{\min(p+1,l)} + C_2 M^{2(1-s)},$$

$$(4.10) \quad \|u^\varepsilon - u^H\|_{L^2(\Omega)} \leq C_1 H^{\min(p+1,l)} + C_2 M^{2(1-s)} + C_3 \varepsilon,$$

$$(4.11) \quad \|u^\varepsilon - u_p^\varepsilon\|_{\bar{H}^1(\Omega)} \leq C_1 H^{\min(p,l)} + C_2 M^{1-s} + C_3 \sqrt{\varepsilon},$$

where C_2 depends on $\|a_{ij}\|_{L^\infty(\Omega; H_{per}^s(Y))}$, $\|\chi^j\|_{L^\infty(\Omega; H_{per}^s(Y))}$, where H is the size of the triangulation of the macro finite element space (2.13), and where M^d is the number of pseudospectral points of the micro space M^d is the number of meshpoints of the micro space $S_M(K_\varepsilon)$ given in (2.15).

THEOREM 4.3. *Let u^0 be the solution of the homogenized problem (2.3), and assume $u^0 \in H^{l+1}(\Omega)$ and $f \in H^l(\Omega)$, $2l > 2$. Let u^H be the solution of problem (4.5) and suppose that the assumptions of Lemma 3.6 hold. Then*

$$(4.12) \quad \|u^0 - u^H\|_{H^1(\Omega)} \leq C_1 H^{\min(p,l)} + C_2 (M(\log M)^{(d-1)} e^{-\alpha M})^2,$$

$$(4.13) \quad \|u^0 - u^H\|_{L^2(\Omega)} \leq C_1 H^{\min(p+1,l)} + C_2 (M(\log M)^{(d-1)} e^{-\alpha M})^2,$$

$$(4.14) \quad \|u^\varepsilon - u^H\|_{L^2(\Omega)} \leq C_1 H^{\min(p+1,l)} + C_2 (M(\log M)^{(d-1)} e^{-\alpha M})^2 + C_3 \varepsilon,$$

$$(4.15) \quad \|u^\varepsilon - u_p^\varepsilon\|_{\bar{H}^1(\Omega)} \leq C_1 H^{\min(p,l)} + C_2 M(\log M)^{d-1} e^{-\alpha M} + C_3 \sqrt{\varepsilon},$$

where C_2 depends on \mathcal{B} , where H is the size of the triangulation of the macro finite element space (2.13) and where M^d is the number of pseudospectral points of the micro space $S_M(K_\varepsilon)$ given in (2.15).

5. Numerical experiments. We end this paper with a few numerical experiments illustrating our theory.

5.1. Example 1: Uniformly periodic problem. In order to see the effect of the spectral micro solver and to compare it with a FEM-type micro solver we consider the (quasi 1-d) model problem (see [3])

$$(5.1) \quad -\nabla \cdot \left(a \left(\frac{x}{\varepsilon} \right) \nabla u^\varepsilon \right) = f(x) \quad \text{in } \Omega = (0, 1)^2$$

$$(5.2) \quad u^\varepsilon|_{\Gamma_D} = 0 \quad \text{on } \Gamma_D := \{x_1 = 0\} \cup \{x_1 = 1\}$$

$$(5.3) \quad n \cdot \left(a \left(\frac{x}{\varepsilon} \right) \nabla u^\varepsilon \right)|_{\Gamma_N} = 0 \quad \text{on } \Gamma_N := \partial\Omega \setminus \Gamma_D,$$

where $a(y) = (\cos 2\pi y_1 + 2)$, $y = (y_1, y_2) \in Y = (0, 1)^2$, and $f(x) \equiv 1$. The homogenized solution is a quadratic polynomial depending on the homogenized tensor which can be easily computed for this simple problem (see [3] for details). The purpose of the following numerical experiments is to illustrate the different impact of the errors in the micro solver at a macroscale, when solving the micro problem with either a FEM or a spectral method. We therefore consider only the homogenized problem in order to be free of the boundary layer term when comparing the fine scale solution with the reconstructed heterogeneous multiscale solution (see Section 3.5).

In the Figures below, we study the macro convergence of the FE-HMM (Finite Element Heterogeneous Multiscale Method with macro and micro piecewise linear

FE spaces) and the FES-HMM (Finite Element Spectral Heterogeneous Multiscale Method with macro piecewise linear FE space and micro pseudospectral method). In both cases, the macro mesh is denoted by H and we denote by $h_L^\varepsilon = h_L/\varepsilon$ the meshsize of the micro FEM, where $h_L = 1/L$. When using pseudo-spectral methods $h_L^\varepsilon = h_L/\varepsilon$ will denote the equidistant spacing of the pseudospectral points and L the number of pseudospectral points (in one dimension).

We first study in Figures 5.1 and 5.2 for a given macro mesh H the influence of the micro mesh refinement on the macro error (for H chosen, $h_L = 1/2, 1/4, 1/10, 1/22, 1/46$). An horizontal line indicates that the micro calculation are precise enough so that the global error is given by the macromesh size (which is kept constant for each experiment). We see for the FES-HMM in Figure 5.1, both for the L^2 error (left picture) and the H^1 error (right picture), that after a short transient $h_L < 1/8$ we have a perfect horizontal line, indicating that the influence of micromesh refinement can no longer be seen. For the FE-HMM, we see in Figure 5.2 (left) that for the L^2 norm, the micro mesh has to be refined, until $h_L \simeq H$ in order to stabilize the global error. We see in Figure 5.2 (right) that for the H^1 norm, the influence of the micro error is less severe and the micro mesh has to be refined, until $h_L \simeq \sqrt{H}$. These behaviours for the FE-HMM are in accordance with the fully discrete error analysis of [3] (see (2.32) for $l = 1$) which indicates that both micro and macro meshes have to be refined simultaneously for the L^2 norm and that micro and macro meshes should be refined according to $h_L \simeq \sqrt{H}$ for the H^1 norm.




FIG. 5.1. FES-HMM: micromesh refinement for fixed macro mesh $H = 2^{-\alpha}$, $\alpha = 1, \dots, 8$.

We next study macro mesh refinements in Figures 5.3 and 5.4. Here we fix the micromesh h_L and refine the macromesh $H = 2^{-\alpha}$, $\alpha = 1, \dots, 8$. We see for the FES-HMM in Figure 5.1, for both the L^2 error (left picture) and the H^1 error (right picture) and for all $h_L \geq 1/8$ we have perfect quadratic and linear convergence. This again confirms our theoretical results: up to exponential convergence, the error of the FES-HMM is given by the usual error of the macro FEM, i.e. *independent* of the micro solver. For the FE-HMM, we see in Figure 5.3 (left) that for the L^2 norm, the quadratic macro convergence rate can be observed until $H \simeq h_L$. Refining further the macromesh does not decrease the global error any further. For the H^1 error, the refinement rate $h_L \simeq \sqrt{H}$ can be observed. Again, these behaviours for the FE-HMM




FIG. 5.2. *FE-HMM: micromesh refinement for fixed macro mesh $H = 2^{-\alpha}$, $\alpha = 1, \dots, 8$.*

are in accordance with the fully discrete error analysis of [3] (see (2.32) for $l = p = 1$)².




FIG. 5.3. *FES-HMM: macromesh refinement for fixed micro mesh $h_L = 1/2, 1/4, 1/10, 1/22, 1/46$.*

5.2. Example 2: Two-scale problem. We next consider a truly two-scale problem (also considered in [25]) given by

$$(5.4) \quad -\nabla \cdot \left(a(x, \frac{x}{\varepsilon}) \nabla u^\varepsilon \right) = f(x) \quad \text{in } \Omega = (0, 1)^2$$

$$(5.5) \quad u^\varepsilon(x) = 0 \quad \text{on } \partial\Omega,$$

²Observe that the convergence plots for the H^1 norm are slightly different than the similar one in [3]. The numerical computation in [3, Figure 4.1, right picture] show a more severe micro error influence than $h_L \simeq \sqrt{H}$ as predicted by the theory given in [3]. This is due to a small error in the code which has been corrected in the present paper. Here, the micro-macro refinement $h_L \simeq \sqrt{H}$ can be observed




FIG. 5.4. *FE-HMM*: : macromesh refinement for fixed micro mesh $h_L = 1/2, 1/4, 1/10, 1/22, 1/46$.

where

$$(5.6) \quad a(x, \frac{x}{\varepsilon}) = \frac{1.5 + \sin(2\pi x_1/\varepsilon)}{1.5 + \sin(2\pi x_2/\varepsilon)} + \frac{1.5 + \sin(2\pi x_2/\varepsilon)}{1.5 + \cos(2\pi x_1/\varepsilon)} + \sin(4x_1x_2) + 1,$$

where $f = 10$ and $x = (x_1, x_2)$. This multiscale problem does not have periodic coefficients but the fast scale is periodic (see Figure 5.5, right picture). In contrast to the previous example, we do not have an analytical solution for the homogenized problem. We will compute a reference solution of the fine scale problem on a very fine mesh of 10^6 degrees of freedom with a classical FE method. As a measure of convergence (see Corollaries 3.10,3.12) study

$$(5.7) \quad \frac{\|u^\varepsilon - u^H\|_{L^2(\Omega)}}{\|u^\varepsilon\|_{L^2(\Omega)}}.$$

We monitor in Figure 5.5 the convergence results for macromesh refinement ($H = 1/2, 1/4, 1/8, 1/16, 1/24, 1/32$). We study the case when δ the cell size is equal to the periodicity ε and the case when the ratio $\delta/\varepsilon = 5/3$ is non integer. The number of pseudospectral points are fixed ($M = 10$ if $\delta = \varepsilon$ and $M = 17$ if $\delta/\varepsilon = 5/3$). We see that we obtain the expected convergence results independent of M if $\delta = \varepsilon$. For noninteger cell size, the convergence deteriorates at the finest macromesh, but is much better than the a-priori estimates one expect for this case, which should involve a boundary layer term $C(\varepsilon/\delta)$ [14]. In general the boundary layer influence for the case of a non-integer ratio between period and cellsize might be larger.

5.3. Example 3: Random porous media problem. For the last example, we consider the multiscale elliptic problem (5.4) with random coefficients $a^\varepsilon(x) = a(x/\varepsilon, \omega)$. This is the typical pressure equation for porous media problems. In such a modeling, the natural media is seen as a statistically homogeneous realization of a random field and the permeability $a^\varepsilon(x)$ varies on a ε length scale much smaller than the characteristic macroscopic length scale of observation. We chose a^ε to be a log-normal stochastic field with mean-zero, variance $\sigma = 1$ and correlation length $\varepsilon_1 = 0.02, \varepsilon_2 = 0.03$. We generate a realization of this stochastic field by the moving ellipse




FIG. 5.5. Convergence of the FES-HMM for the example 2 (left picture), snapshot of the conductivity field (right picture).

averaging method [30]. We then compute a reference solution on a fine 1024×1024 grid and compare the solution with the FES-HMM (the reference solution can be seen as the exact solution of a finite difference version of (5.4)) [25]. We compute a solution on a coarse 32×32 grid for the FES-HMM with a sampling domain of size 0.06×0.06 . The error estimates for random coefficients are much weaker (see [14]) and a realistic complexity estimates will require improved analysis.




FIG. 5.6. Comparison of the fine scale solution (pressure profile) of problem (5.4) with random coefficients on a 1024×1024 grid and the FES-HMM solution on a 32 grid.

(5.4)

It can be seen in Figure 5.6 that the solution obtained from the FES-HMM on a coarse grid (32×32) is in good qualitative agreement with the solution of the standard FEM on the fine grid (1024×1024 points).

Acknowledgments. The authors are grateful to Christoph Schwab for helpful discussions.

Appendix A. Spectral estimates. Spectral and exponential convergence results for Fourier pseudo-spectral methods are usually given for the 1–dimensional case [16],[22],[29],[10]. We briefly discuss in this appendix these approximation results for trigonometric polynomials and extend them, with a tensor product argument, to the multi-dimensional cases needed for our convergence results.

We consider the notation introduced in Section 2.2. Let $u = \sum_{k=-\infty}^{k=\infty} \hat{u}_k e^{2ik\pi y} \in L^2(I)$ and $Q_M(u) = \sum_{k=-M}^{k=M} \hat{u}_k \psi_k$, $|k| \leq M$ the trigonometric interpolant of u at the pseudospectral points I_M defined in (2.8). In order to estimate the difference $\|u - Q_M(u)\|$ in the H^1 or in the L^2 norm, the following lemma, so-called aliasing lemma, is useful.

LEMMA A.1. *Assume $u \in C_{per}^0(I)$, then*

$$(A.1) \quad \hat{u}_k - \tilde{u}_k = \sum_{l \neq 0, l=-\infty}^{l=\infty} \hat{u}_{k+2Ml}, \quad -M \leq k \leq M.$$

This lemma can be proved simply by inserting the Fourier expansion (2.6) in (2.9), and using the orthogonality property

$$(A.2) \quad \frac{1}{2M} \sum_{l=0}^{2M-1} e^{-2i\pi k y_l} = 1 \text{ if } k = 0 \pmod{2M} \text{ and } 0 \text{ otherwise.}$$

Following [29], using (2.6), (2.9) and (A.1) we can write the difference $u - Q_M(u)$ as

$$(A.3) \quad u - Q_M(u) = - \sum_{k=-M}^{k=M} \left(\sum_{l \neq 0, l=-\infty}^{l=\infty} \hat{u}_{k+2Ml} \right) e^{2i\pi k x} + \sum_{|k| \geq M} \hat{u}_k e^{2i\pi k x}.$$

The next lemma gives the so-called spectral accuracy estimates for $\|u - Q_M(u)\|$ and is obtained by estimating both terms in the right-hand side of (A.3) (see for example [22] or [29] for details). [10, p.279].

LEMMA A.2. *Assume $u \in H_{per}^s(I)$ with $s > 1/2$, then for any real σ $0 \leq \sigma \leq s$*

$$(A.4) \quad \|u - Q_M(u)\|_{H^\sigma(I)} \leq CM^{\sigma-s} \|u\|_{H^s(I)}$$

Using (A.4) with $\sigma = s = 1$ gives the following stability result

$$(A.5) \quad \|\nabla Q_M(u)\|_{L^2(I)} \leq C \|u\|_{H^1(I)}.$$

For the estimates used in Section 3 we need the following lemma, proved in [29], which gives an exponential decay estimate, provided u is analytic. (see Definition 3.4).

LEMMA A.3. *Suppose $u \in \mathcal{A}_{B,\alpha}(I)$, then*

$$(A.6) \quad \|u - Q_M(u)\|_{H^\sigma(I)} \leq CB(\alpha) M^\sigma e^{-\alpha M},$$

where $B(\alpha) = \max_{|\Im z| \leq \alpha} |u(z)|$ and the constant C depends on α and σ . We will also need the following L^∞ –stability estimate

$$(A.7) \quad \|Q_M(u)\|_{L^\infty(I)} \leq C \log M \|u\|_{L^\infty(I)},$$

which is proved in [20][p.119-121] and the result of Bernstein [6] (see also [24][pp.56-57])

$$(A.8) \quad \|\nabla Q_M(u)\|_{L^\infty(I)} \leq M \|Q_M(u)\|_{L^\infty(I)}.$$

In Section 3 we used d -dimensional version of the above approximation results in $Y = I^d = (0, 1)^d$. We recall the tensor product construction given in (2.11) and the notation $\mathbf{Q}_M(u) = Q_M^1 \cdots Q_M^d(u)$ introduced in Section 2.2. For simplicity of notation we present the proof for the case $d = 2$ and note $\mathbf{Q}_M(u) = Q_M^x Q_M^y(u)$. We emphasize that the same proofs work in higher dimension. We will also use the isomorphism $L^2(Y) = L^2(I; L^2(I))$, with the usual definition of the latter space.

LEMMA A.4. *Let $Y = (0, 1)^d$, $d = 2, 3$ and assume $u \in H_{per}^s(Y)$ with $s \geq 2$, then*

$$(A.9) \quad \|u - \mathbf{Q}_M(u)\|_{L^2(Y)} \leq CM^{-s} \|u\|_{H^s(Y)}$$

$$(A.10) \quad \|\nabla u - \nabla \mathbf{Q}_M(u)\|_{L^2(Y)} \leq CM^{1-s} \|u\|_{H^s(Y)}$$

Proof. Following [8], we write (for $d = 2$) $\|u - Q_M^x Q_M^y(u)\|_{L^2(Y)}$
 $\leq \|u - Q_M^x(u)\|_{L^2(I; L^2(I))} + \|u - Q_M^y(u)\|_{L^2(I; L^2(I))} + \|(Id - Q_M^x)(Id - Q_M^y)u\|_{L^2(I; L^2(I))}$
 $\leq CM^{-s} \|u\|_{H^s(I; L^2(I))} + CM^{-s} \|u\|_{L^2(I; H^s(I))} + CM^{-1} M^{-(s-1)} \|u\|_{H^1(I, H^{s-1}(I))},$

where we used (A.4) for the x and y variables in the first and second terms of the right-hand side of the inequality, and successively (A.4) for the x variable and for the y variable. Using the continuous embedding $H^s(Y) \subset H^1(I, H^{s-1}(I))$ concludes the proof of (A.9). To prove (A.10) we first estimate $\|\partial_x u - \partial_x Q_M^x Q_M^y(u)\|_{L^2(Y)}^2$

$$\leq \|\partial_x u - \partial_x Q_M^x(u)\|_{L^2(I; L^2(I))}^2 + \|\partial_x Q_M^x(u - Q_M^y(u))\|_{L^2(I; L^2(I))}^2$$

$$\leq CM^{2(1-s)} \|u\|_{H^s(I; L^2(I))}^2 + C(\|\partial_x(u - Q_M^y(u))\|_{L^2(I; L^2(I))}^2 + \|(u - Q_M^y(u))\|_{L^2(I; L^2(I))}^2),$$

where we used (A.4) and the stability result (A.5). In the second term of the last inequality, we observe that ∂_x and Q_M^y commute. We can then use (A.4) for both the second and the last term and bound them by $CM^{2(1-s)} \|u\|_{H^s(Y)}^2$. Using a similar argument for $\|\partial_y u - \partial_y Q_M^x Q_M^y(u)\|_{L^2(Y)}^2$, summing up and taking the square root yields (A.10). \square

LEMMA A.5. *Let $Y = (0, 1)^d$ and suppose $u \in \mathcal{A}_{\mathcal{B}, \alpha}(Y)$, then*

$$(A.11) \quad \|u - \mathbf{Q}_M(u)\|_{L^2(Y)} \leq C\mathcal{B} \log M^{d-1} e^{-\alpha M},$$

$$(A.12) \quad \|\nabla u - \nabla \mathbf{Q}_M(u)\|_{L^2(Y)} \leq C\mathcal{B}M \log M^{d-1} e^{-\alpha M},$$

where $\mathcal{B} = \max_{i \leq d} \{\max_{Y_{-i}} \max_{z \in \mathcal{E}_\alpha} |u(Y_{-i}, z)|\}$ (see 3.4) and the constant C depends on α .

Proof. For (A.11), we have (for $d = 2$) $\|u - Q_M^x Q_M^y(u)\|_{L^2(Y)}$
 $\leq \|u - Q_M^x(u)\|_{L^2(I; L^2(I))} + \|u - Q_M^y(u)\|_{L^2(I; L^2(I))} + \|(Id - Q_M^x)(Id - Q_M^y)u\|_{L^2(I; L^2(I))}$

For the first term of the right-hand side of the inequality, we use (A.6) and (A.7) and bound it by

$$\|u - Q_M^x(u)\|_{L^2(I; L^2(I))}^2 = \int_I \int_I |u(x, y) - Q_M^x(u)(x, y)|^2 dx dy$$

$$\leq Ce^{-2\alpha M} \max_{z \in \mathcal{E}_\alpha} \int_I |u(z, y)|^2 dy \leq Ce^{-2\alpha M} \max_{z \in \mathcal{E}_\alpha} \max_{y \in I} |u(z, y)|^2 \leq C\mathcal{B}^2 e^{-2\alpha M}.$$

The second term is treated similarly. For the last term we use (A.6) and (A.7) and bound it by

$$\begin{aligned} \|(Id - Q_M^x)(Id - Q_M^y)u\|_{L^2(I;L^2(I))}^2 &\leq C e^{-2\alpha M} \max_{z \in \mathcal{E}_\alpha} \int_I |u(z, y) - Q_M^y(u)(z, y)|^2 dy \\ &\leq C e^{-2\alpha M} \max_{z \in \mathcal{E}_\alpha} \max_{y \in I} |u(z, y) - Q_M^y(u)(z, y)|^2 \leq C \mathcal{B}^2 (\log M)^2 e^{-2\alpha M}. \end{aligned}$$

Summing and taking the square root yields (A.11).

To prove (A.12) we first estimate $\|\partial_x u - \partial_x Q_M^x Q_M^y(u)\|_{L^2(Y)}$

$$\leq \|\partial_x u - \partial_x Q_M^x(u)\|_{L^2(I;L^2(I))}^2 + \|\partial_x Q_M^x(u - Q_M^y(u))\|_{L^2(I;L^2(I))}^2.$$

For the first term of the right-hand side, we use (A.6) and bound it by $C \mathcal{B}^2 M^2 e^{-2\alpha M}$. For the second term of the right-hand side, we first use (A.8) and then (A.7) and we obtain the bound $C \mathcal{B}^2 M^2 \log M^2 e^{-2\alpha M}$. Using a similar argument for $\|\partial_y u - \partial_y Q_M^x Q_M^y(u)\|_{L^2(Y)}$ summing up and taking the square root yields (A.12). \square

REFERENCES

- [1] A. Abdulle and W. E, *Finite difference HMM for homogenization problems*, J. Comput. Phys. 191, no. 1, pp. 18–39, 2003.
- [2] A. Abdulle and C. Schwab, *Heterogeneous Multiscale FEM for Diffusion Problem on Rough Surfaces*, SIAM Multiscale Model. Simul., Vol. 3, No. 1, pp. 195–220, 2005.
- [3] A. Abdulle, *On a-priori error analysis of Fully Discrete Heterogeneous Multiscale FEM*, SIAM Multiscale Model. Simul., Vol. 4, No. 2, pp. 447–459, 2005.
- [4] A. Abdulle, *Multiscale methods for advection-diffusion problems*, Discrete Contin. Dyn. Syst. B, Suppl. Vol., pp. 11–21, 2005
- [5] A. Abdulle, *Analysis of a Heterogeneous Multiscale FEM for Problems in Elasticity*, Math. Mod. Meth. Appl. Sci. (M3AS), Vol. 16, No 2, pp.1–21, 2006.
- [6] S.N. Bernstein, *Sur l'ordre de la meilleure approximation des fonctions continues par les polynôme de degré donné*, Mémoire de l'Acad. Royale de Belgique 4, pp.1–104, 1912.
- [7] A. Bensoussan, J.-L. Lions and G. Papanicolaou, *Asymptotic Analysis for Periodic Structures*, North Holland, Amsterdam, 1978.
- [8] C. Bernardi, Y. Maday and F. Rapetti, *Discrétisations variationnelles de problèmes aux limites elliptiques*, Mathématiques et Applications, Vol. 45, Springer 2004.
- [9] L. Bers, F. John and M. Schechter, *Partial Differential Equation*, Lectures in Applied Mathematics, Proceedings of the Summer Seminar, Boulder Colorado, AMS, Providence, Rhode Island, 1957.
- [10] C. Canuto, M.Y. Hussaini, A. Quarteroni and T.A. Zang, *Spectral methods in fluid dynamics*, Springer, 1987.
- [11] D. Cioranescu and P. Donato, *An introduction to Homogenization*, Oxford University Press, 1999.
- [12] P.G. Ciarlet. *The Finite Element Method for Elliptic Problems*, North-Holland, Amsterdam, 1978.
- [13] W. E and B. Engquist, *The Heterogeneous Multi-Scale Methods*, Commun. Math. Sci., Vol. 1, No. 1, pp. 87–132, 2003.
- [14] W. E, Pingbing Ming and Pingwen Zhang, *Analysis of the heterogeneous multi-scale method for elliptic homogenization problems*, J. Amer. Math. Soc. Vol. 18, pp. 121–156, 2004.
- [15] B. Engquist and Y.-H. Tsai, *Heterogeneous multiscale methods for stiff ordinary differential equations*, Math. Comp., 74, pp. 1707–1742, 2005.
- [16] D. Gottlieb and S. Orszag, *Numerical Analysis of Spectral Methods: Theory and applications*, CBMS Regional Conference Series in Applied Mathematics, SIAM, Philadelphia, 1977.
- [17] V. H. Hoang and S. Schwab, *High-dimensional finite elements for elliptic problems with multiple scales*, SIAM Multiscale Model. Simul., Vol. 3, No. 1, pp. 168–194, 2005.
- [18] T-Y. Hou, X-H. Wu and Z. Cai, *Convergence of a multi-scale finite element method for elliptic problems with rapidly oscillating coefficients*, Math. of Comput., V. 68, N. 227, 1999, pp. 913–943.

- [19] T. Hou, X.H. Wu, *A multiscale finite element method for elliptic problems in composite materials and porous media*, J. Comp. Phys. 134, pp. 169-189, 1997.
- [20] D. Jackson, *The theory of approximation*, Vol.11, AMS Colloquium Publications, New York, 1930.
- [21] V.V. Jikov, S.M. Kozlov and O.A. Oleinik, *Homogenization of differential Operators and Integral Functionals*, Springer-Verlag, Berlin, Heidelberg, 1994.
- [22] H.-O. Kreiss and J. Oliger, *Stability of the Fourier Method*, SIAM J. Numerical Analysis, Vol. 16, No. 3, pp.421-433, 1979.
- [23] O. A. Ladyzhenskaya and N.N. Ural'tseva, *Linear and quasilinear elliptic equations*, Academic Press, New York, 39, 1968.
- [24] G. Meinardus, *Approximation von Funktionen und ihre numerische Behandlung*, Springer Tracts in Natural Philosophy, V. 4, 1964.
- [25] P. Ming and Y. Yue, *Numerical methods for multiscale elliptic problems*, J. Comput. Physics, V. 214, No 1, p. 421-445.
- [26] S. Moskow and M. Vogelius, *First order corrections to the homogenized eigenvalues of a periodic composite medium*, Proc. Roy Soc. Edinburgh. Sec. A., 127, 1997, pp.1263-1299.
- [27] L. Nirenberg and C.B.Jr. Morrey, *On the analyticity of the solution of linear elliptic systems of partial differential equations*, Comm. Pure Appl. Math.,**10**, 1957, pp.271-290.
- [28] J.T. Oden and K.S. Vemaganti, *Estimation of local modeling error and global-oriented adaptive modeling of heterogeneous materials: error estimates and adaptive algorithms*, J. Comput. Phys. 164, No. 1, 2000, pp. 22-47.
- [29] E. Tadmor, *The exponential accuracy of Fourier and Chebyshev differencing methods*, SIAM Journal on Numerical Analysis, V.23 No.1, pp. 1-10, 1986.
- [30] T.C. Wallstrom, S. Hou, M.A. Christie, L.J. Durlofsky, D.H. Sharp, *Accurate scale up of two phase flow using renormalization and nonuniform coarsening*, Computational Geosciences, Vol. 3, 1, 1999, pp. 69-87.
- [31] X.H. Wu, Y. Efendiev and T.Y. Hou, *Analysis of upscaling absolute permeability*, Discrete Contin. Dyn. Syst. B, Vol. 2, No. 2, 2002, pp. 186-204.
- [32] X. Yue and W. E, *The local microscale problem in the multiscale modeling of strongly heterogeneous media: effect of boundary conditions and cell size*, J. Comput. Phys., Vol. 222, 2, 2007, pp. 556-572.