

ECOLE POLYTECHNIQUE

CENTRE DE MATHÉMATIQUES APPLIQUÉES

UMR CNRS 7641

91128 PALAISEAU CEDEX (FRANCE). Tél: 01 69 33 41 50. Fax: 01 69 33 30 11

<http://www.cmap.polytechnique.fr/>

**LAMN property
for integrated diffusions**

Arnaud GLOTER, Emmanuel GOBET

R.I. 570

March 2005

LAMN property for integrated diffusions

Arnaud GLOTER* and Emmanuel GOBET†

March 25, 2005

Abstract

In this paper we prove the Local Asymptotic Mixed Normality (LAMN) property for the statistical model given by the observation of local means of a diffusion process X . Our data are given by $\int_0^1 X_{\frac{s+i}{n}} d\mu(s)$ for $i = 0, \dots, n-1$ and the unknown parameter appears in the diffusion coefficient of the process X only. Although the data are not Markovian neither Gaussian we can write down, with help of Malliavin calculus, an explicit expression for the log-likelihood of the model, and then study the asymptotic expansion. We actually find that the asymptotic information of this model is the same one as for a usual discrete sampling of X .

KEYWORDS: Diffusion processes, parametric estimation, LAMN property, Malliavin calculus, non-Markovian data

AMS 2000 SUBJECT CLASSIFICATION: 60Fxx; 60Hxx; 62Fxx; 62Mxx

1 Statement of the problem and main results

1.1 Introduction

Let us consider the family of strong solutions, X^θ , of the stochastic differential equation,

$$dX_t^\theta = a(X_t^\theta, \theta)dB_t + b(X_t^\theta)dt, \quad (1)$$

$$X_0^\theta = \xi_0, \quad (2)$$

where $(B_t)_{t \geq 0}$ is a one dimensional standard Brownian motion on some probability space (Ω, \mathcal{A}, P) , and where the initial value ξ_0 is some real constant. Suitable regularity assumptions on the coefficients b and a will be stated

*Université Bordeaux 4, GRAPE UMR 5113, av. Léon Duguit, 33608 Pessac Cedex, FRANCE - email: gloter@u-bordeaux4.fr

†Ecole Polytechnique, Centre de Mathématiques Appliquées, 91128 Palaiseau Cedex, FRANCE - email:emmanuel.gobet@polytechnique.fr - Corresponding author.

later (see assumption **(R)**). The unknown parameter θ is assumed to lie on some compact interval Θ of \mathbb{R} and for simplicity we have assumed that the initial value ξ_0 does not depend on θ and thus is known to the statistician.

Now, let μ be some probability measure on $[0, 1]$, and we assume that our observation of the process is given by the local means of X associated with this measure, with sampling of size $1/n$:

$$\textbf{(observations)} \quad \bar{X}_j = \bar{X}_{j,n} := \int_0^1 X_{\frac{s+j}{n}} d\mu(s), \text{ for } j = 0, \dots, n-1.$$

We also assume that μ does not depend on θ and is known by the statistician.

Clearly, the usual case of pointwise observation of X is obtained if μ is some Dirac measure. However we will exclude that the measure has mass only on the end points of the interval and hence make the assumption:

$$\mu((0, 1)) > 0. \tag{3}$$

This paper is concerned with the Local Asymptotic Mixed Normality property of this statistical model.

Motivations. Taking as the observation the integrated process instead of the process itself is actually quite natural. For instance, it arises when the realization of the process has been observed after passage through an electronic filter. Also, in random mechanics (see Krée and Soize [18]), X models the velocity of the system and in general, we observe its position, i.e. the integral of X . The modeling of ice-core data can be made through an integrated diffusion process (see Ditlevsen, Ditlevsen and Andersen [2]). Integrated processes also play an important role in finance, when modelling the stochastic volatility (see for instance Barndorff-Nielsen and Shepard [1] and references therein).

Literature background. Despite of these numerous motivations, a few statistical studies deal with this situation. The inherent difficulty of this situation is that the observation is no longer a Markov process, which eliminates many statistical and probabilistic tools. Gloter [6] [7] provides an estimator in the multiplicative case $a(x, \theta) = \theta a(x)$ and proves its consistency and asymptotic normality in the high frequency case $t_i = i/n$. The case of low frequency (say $t_{i+1} - t_i = 1$) is studied by Ditlevsen and Sørensen [3], using prediction-based estimating functions. For a direct observation of the diffusion, see Genon-Catalot and Jacod [5], Prakasa-Rao [23] and references therein.

Here, we directly address the problem of the LAMN property, which fundamental consequence is to provide information on the minimal dispersion for an estimator of the parameter θ (see Ibragimov and Has'minskii [13], Jeganathan [15] [16], Prakasa-Rao [23], Le Cam and Lo Yang [20]). Such properties in the diffusion case have been established in the one-dimensional setting by Dohndal [4], and then extended by the Gobet [9] [10] to the multi-dimensional setting, both in the high frequency and ergodic framework. For

this, Malliavin calculus techniques were used and paved the way to possibly handle more general situations than Markovian observations. This is exactly this way we follow in this work, to tackle the case of integrated diffusion.

1.2 Main results

Before going into the details of our results, we present a very simple example which gives an intuition on the type of results that one can expect.

Example 1 (Multiplicative Brownian case). *Assume that the model is*

$$X_t^\theta = \theta B_t$$

(corresponding to $b \equiv 0$ and $a(\cdot, \theta) = \theta, \xi_0 = 0$).

1. Consider a first situation where one observes the diffusion at discrete times. Hence, the observation is $(X_{t_i})_{1 \leq i \leq n}$, or equivalently $(Z_i = \theta(B_{t_i} - B_{t_{i-1}}) = \theta G_i)_{1 \leq i \leq n}$, where G_i are independent centered Gaussian variables, with a known variance. Thus, the estimation of θ^2 is achieved at rate \sqrt{n} , with a minimal variance equal to $2\theta^4$.
2. Now consider a second situation where one observes only the integrated diffusion at discrete times. Hence, the observation is $(\bar{X}_i = \theta \int_0^1 B_{\frac{(s+i)}{n}} \mu(ds) = \theta G'_i)_{1 \leq i \leq n}$, where $(G'_i)_i$ is a centered Gaussian vector, with a known covariance matrix. In addition, this matrix is invertible and thus, θ^2 can be estimated in the same conditions than before.

This means that observing the process at discrete times or its integrated version is equivalent for the inference problem. The results of this paper state that this is true, even for the more general models (1-2), which is far to be intuitive.

Before stating our main results, we define the working assumptions of this paper. The coefficients $a : \mathbb{R} \times \Theta \rightarrow \mathbb{R}$ and $b : \mathbb{R} \rightarrow \mathbb{R}$, are assumed to satisfy the following set of conditions (as usual, derivatives w.r.t. θ are denoted with a dot: for instance, $\partial_\theta a = \dot{a}$).

Assumption (R)

- 1) The function a is $\mathcal{C}^{1+\gamma}$ for some $\gamma \in (0, 1)$ and the functions $x \mapsto a(x, \theta), x \mapsto \dot{a}(x, \theta), x \mapsto b(x)$ are assumed to be $\mathcal{C}^3(\mathbb{R})$.
- 2) The functions a, \dot{a} and b and all their derivatives with respect to x are bounded uniformly in θ .
- 3) We have the non degeneracy condition, for some $\underline{a}: a(x, \theta) > \underline{a} > 0$ for all x, θ .

Actually, the uniform controls in **(R)** can be weakened to local ones, using extra techniques of space localization (see Lemma 4.1 in [9]). We omit further details. An extension of our results to a multidimensional parameter θ and to time dependent coefficients is straightforward, in the same way that it is done in [9] and [10].

We denote by \mathbb{P}^θ the law on $\mathcal{C}([0, 1])$ of the process X^θ , and then simply denote X the canonical process on $\mathcal{C}([0, 1])$. We let $\mathbf{p}^{\mathbf{n}, \theta}$ denote the law on \mathbb{R}^n of the observation $\mathcal{O}^n := (\bar{X}_j)_{j=0, \dots, n-1}$, when the true value of the parameter is θ . And for θ_0, θ_1 two values of the parameter we introduce the likelihood ratio,

$$Z_{\theta_0, \theta_1}^n = \frac{d\mathbf{p}^{\mathbf{n}, \theta_1}}{d\mathbf{p}^{\mathbf{n}, \theta_0}}(\mathcal{O}^n). \quad (4)$$

The main result, is that this statistical model satisfy the so called LAMN property. For this denote the sequence $u_n := n^{-1/2}$, and let $\theta_0 \in \Theta$ and $h \in \mathbb{R}$ such that $\theta_0 + u_n h \in \Theta, \forall n$. Then, by the following theorem, the model has the LAMN property for the likelihood at point θ_0 , with rate u_n and conditional information:

$$\mathcal{I}_{\theta_0} = 2 \int_0^1 \left(\frac{\dot{a}}{a} \right)^2 (X_s, \theta_0) ds.$$

Theorem 1. *Assume **(R)**, then we have the expansion,*

$$\log Z_{\theta_0, \theta_0 + u_n h}^n = h N_n + 1/2 h^2 I_n + R_n,$$

where $I_n \xrightarrow[n \rightarrow \infty]{\mathbb{P}^{\theta_0}} -\mathcal{I}_{\theta_0}$, $R_n \xrightarrow[n \rightarrow \infty]{\mathbb{P}^{\theta_0}} 0$ and there exists an extra random variable $N \sim \mathcal{N}(0, 1)$ independent of \mathcal{I}_{θ_0} such that, N_n converges in law under \mathbb{P}^{θ_0} to $N\sqrt{\mathcal{I}_{\theta_0}}$.

Moreover this convergence is stable, which in particular implies the joint convergence under \mathbb{P}^{θ_0} :

$$(I_n, N_n) \xrightarrow[n \rightarrow \infty]{law} (-\mathcal{I}_{\theta_0}, N\sqrt{\mathcal{I}_{\theta_0}}).$$

Remark 1. *Let us stress that the rate $u_n = n^{-1/2}$ and the information \mathcal{I}_{θ_0} are the same one as for the pointwise observation (see Genon-Catalot and Jacod [5]). This corroborates the intuition from Example 1.*

However, we will not be able to prove directly this result, but instead we shall consider first the easier problem where one can observe additionally the exact value of the diffusion at some sparse instants. This device was proved to be useful in Gloter and Jacod [8] for the study of a Gaussian diffusion process observed with noise that leads to non Markovian observations too.

Let $k = k_n$ be an integer in $\{1, \dots, n\}$ and define $L = L_n := \lfloor n/k \rfloor$, then we consider the set of random variables:

$$\mathcal{O}^{n, \text{aug}} = \mathcal{O}^n \cup \left\{ X_{\frac{kl}{n}}, l = 1, \dots, L \right\} \cup \{X_1\}.$$

Since this set of variables contains more data than the initial set, we call it the *augmented observation* set. Clearly, we can split the set of augmented observations into blocks, $\mathbb{B}_0, \dots, \mathbb{B}_L$, where for $l = 0, \dots, L - 1$

$$\mathbb{B}_l = \{\bar{X}_{kl}, \dots, \bar{X}_{kl+k-1}, X_{k(l+1)/n}\}$$

and $\mathbb{B}_L = \{\bar{X}_{kL}, \dots, \bar{X}_{n-1}, X_1\}$. Note that if $kL = n$ we consider that the last block is empty, and (immediate) modifications should take care of this in the sequel, however to have shorter notations we will not explicitly write these modifications.

The advantage of this set of augmented observation is that using the Markov property of X , the law the block \mathbb{B}_l conditional to the previous blocks $(\mathbb{B}_{l'})_{l' < l}$ only depends on the last variable, $X_{\frac{kl}{n}}$, of the block \mathbb{B}_{l-1} .

Denote by $\mathbf{p}^{\mathbf{n}, \text{aug}, \theta}$ the law of $\mathcal{O}^{\mathbf{n}, \text{aug}}$ on \mathbb{R}^{n+L+1} and introduce the likelihood ratio for the augmented observation:

$$Z_{\theta_0, \theta_1}^{\mathbf{n}, \text{aug}} = \frac{d\mathbf{p}^{\mathbf{n}, \text{aug}, \theta_1}}{d\mathbf{p}^{\mathbf{n}, \text{aug}, \theta_0}}(\mathcal{O}^{\mathbf{n}, \text{aug}}). \quad (5)$$

Theorem 2. *There exists a sequence $k_n \rightarrow \infty$, such that the augmented model satisfies the LAMN property:*

$$\log Z_{\theta_0, \theta_0 + u_n h}^{\mathbf{n}, \text{aug}} = hN_n^{\text{aug}} + 1/2h^2 I_n^{\text{aug}} + R_n^{\text{aug}},$$

where $I_n^{\text{aug}} \xrightarrow[n \rightarrow \infty]{\mathbb{P}^{\theta_0}} -\mathcal{I}_{\theta_0}$, $R_n^{\text{aug}} \xrightarrow[n \rightarrow \infty]{\mathbb{P}^{\theta_0}} 0$ and there exists an extra random variable $N \sim \mathcal{N}(0, 1)$ independent of \mathcal{I}_{θ_0} such that, N_n^{aug} converges in law under \mathbb{P}^{θ_0} to $N\sqrt{\mathcal{I}_{\theta_0}}$.

Moreover this convergence is stable, which implies in particular the joint convergence under \mathbb{P}^{θ_0} :

$$(I_n^{\text{aug}}, N_n^{\text{aug}}) \xrightarrow[n \rightarrow \infty]{\text{law}} (-\mathcal{I}_{\theta_0}, N\sqrt{\mathcal{I}_{\theta_0}}).$$

Remark 2.

- 1) *The fact that $k_n \rightarrow \infty$ means that the data added in the observation are sparse compared to the initial data. Actually, the Theorem 2 holds for any sequence k_n which growth to ∞ is slow enough.*
- 2) *If one let the size of the block $k_n = k$ fixed as $n \rightarrow \infty$ then the augmented model satisfy the LAMN property too but with an increased asymptotic information (see Section 3.3).*

From Theorem 2 and from the consequences of the LAMN property, an *asymptotically optimal* estimator θ_n in the augmented model should satisfy that $\sqrt{n}(\theta_n - \theta_0)$ is asymptotically distributed under \mathbb{P}_{θ_0} as $\frac{1}{\sqrt{\mathcal{I}_{\theta_0}}}N$. However any estimator in the initial model of observation \mathcal{O}^n can be seen as an estimator in the augmented model, hence the Theorem 2 would be sufficient by itself to imply a lower bound for estimation in the initial model.

1.3 Outline of the paper

In Section 2 we study the score function given by the observation of only one block of data (\mathbb{B}_0 for instance). We first focus on the existence of a density for a block of data; and in the case of a block of size 2, $(n^{1/2} \int_0^1 (X_{s/n} - X_0) d\mu(s), n^{1/2}(X_{\frac{1}{n}} - X_0))$ we give original lower and upper bounds of Gaussian type for the density. It is useful for our proof of the LAMN property, but it is also interesting for itself.

In Section 2.2 we present an exact expression for the score function of a block of data \mathbb{B}_0 (see Theorem 4). This result is the key point in the proof of the LAMN property, it extends a former result of Gobet [9] [10] which gave the score function for the observation of $X_{\frac{\theta}{n}}$. In Section 2.3 we study an explicit approximation for the score function when the sampling interval $1/n$ tends to zero and the length of the block k/n remains moderate so that one can consider the coefficients of the diffusion X almost constant on the interval $[0, k/n]$. The key point is the Gaussian approximation for the diffusion given in Section 2.3.1.

In Section 3 we deduce from the previous results a proof of Theorem 2 and Section 4 deals on how to deduce Theorem 1 from Theorem 2.

Finally the Appendix contains the proof of some results of Section 2.1 together with some useful lemmas.

Notations. In our proofs, we will keep the same notation for constants which may change from one line to another. In particular, the constants $c, c(k), c(p), c(p, k)$ will stand for all finite, non-negative and non-decreasing deterministic functions of an index p (arising from \mathbf{L}^p -norm) and of the block size k . These constants are independent on n, θ and depend on the process X^θ , only through the bounds on the coefficients a, b and their derivatives.

2 Score function for a block of data

In this section we shall study the law of the blocks of data \mathbb{B}_l ; recalling the Markov property of the process X it is sufficient to focus on $\mathbb{B}_0 = \{\bar{X}_0, \dots, \bar{X}_{k-1}, X_{k/n}\}$ assuming that the diffusion X now starts from some value x_0 . In this section it is convenient to transform the short time asymptotic $k/n \rightarrow 0$ into an almost stationarity property of the coefficients. To this end, we introduce the rescaled process $\mathcal{X}_t^{n,\theta} = n^{\frac{1}{2}}(X_{\frac{t}{n}}^\theta - x_0)$ (where X^θ solves (1) with $X_0^\theta = x_0$). It solves the equation

$$d\mathcal{X}_t^{n,\theta} = a_n(\mathcal{X}_t^{n,\theta}, \theta) dW_t + b_n(\mathcal{X}_t^{n,\theta}) dt, \quad \mathcal{X}_0^{n,\theta} = 0, \quad (6)$$

where W is a standard Brownian motion (arising from the rescaling of B), and

$$a_n(x, \theta) = a(x_0 + n^{-1/2}x, \theta), \quad b_n(x) = n^{-1/2}b(x_0 + n^{-1/2}x). \quad (7)$$

Since for the score we are only concerned with the law of $\mathcal{X}^{n,\theta}$, we can assume that W is independent of the rescaling coefficient n .

2.1 The density of an integrated diffusion

In this section, we will present preliminary results on the density of the law of the mean of a diffusion process. However the proofs are postponed to Section 5.1. To our knowledge, the lower and upper bounds for this density are new results.

2.1.1 Existence of the density

Our first result actually deals with the two dimensional variable given by solely one local mean and the exact value:

$$(U^{n,\theta}, V^{n,\theta}) := \left(\int_0^1 \mathcal{X}_s^{n,\theta} d\mu(s), \mathcal{X}_1^{n,\theta} \right) \quad (8)$$

$$\stackrel{\text{law}}{=} \left(n^{1/2} \int_0^1 (X_{\frac{s}{n}}^\theta - x_0) d\mu(s), n^{1/2} (X_{\frac{1}{n}}^\theta - x_0) \right).$$

Notice that, by the Markov property, the preliminary study of this bi-dimensional variable will be a key step to obtain results on the observation vector \mathcal{O}^n .

Theorem 3. *Assume (\mathbf{R}) , then the vector $(U^{n,\theta}, V^{n,\theta})$ admits a density $\mathbf{p}_{x_0}^n(\cdot, \cdot, \theta)$ on \mathbb{R}^2 , and there exist two constants $c_1 > c_2 > 0$, such that,*

$$c_1^{-1} e^{-c_1(u^2+v^2)} \leq \mathbf{p}_{x_0}^n(u, v, \theta) \leq c_2^{-1} e^{-c_2(u^2+v^2)}. \quad (9)$$

The constants c_1 and c_2 only depend on the bounds on the coefficients a, b and their derivatives.

The proof of this theorem is given in Section 5.1. The existence of the density is obtained by means of the Malliavin calculus. On the other hand, the upper and lower bounds rely on the direct study of $(U^{n,\theta}, V^{n,\theta})$ around its skeleton (see Hirsh and Song [11] [12] for related works; and Kohatsu-Higa [17] for different methods involving Malliavin calculus).

The following is a direct corollary of Theorem 3:

Corollary 1. *The vector $\mathbb{B}_0 = \{\bar{X}_0, \dots, \bar{X}_{k-1}, X_{k/n}\}$ admits a positive density.*

Proof. The bi-dimensional process $(\bar{X}_l, X_{\frac{l+1}{n}})_{l=0, \dots, k-1}$ is a Markov chain with transition density $p_{x_l}(\bar{x}_l, x_{l+1}, \theta) = n \mathbf{p}_{x_l}^n(n^{\frac{1}{2}}(\bar{x}_{l+1} - x_l), n^{\frac{1}{2}}(x_{l+1} - x_l), \theta)$. Then it is clear that the vector \mathbb{B}_0 admits a positive density. \square

2.1.2 Invertibility of the Malliavin covariance matrix of a block

Actually the existence of a density for the law of the random variable \mathbb{B}_0 will not be sufficient, and we need a non degeneracy condition for this variable.

Before this, let us precise briefly a few notations from the Malliavin calculus, used in the sequel (see Nualart [21] [22] for details). We let H be the Hilbert space $\mathbf{L}^2([0, 1])$ so that the Brownian motion $(W_t)_{t \in [0, 1]}$, appearing in (6), is canonically associated to this Hilbert space via the standard \mathbf{L}^2 isometry. In this setting, for any $p \geq 1$ and natural number q , recall that the set $\mathbb{D}^{q,p}$ denotes the space of real valued Wiener functionals with q derivatives and whose derivatives belong to $\mathbf{L}^p(\Omega)$. If we denote by D the derivative operator then the space $\mathbb{D}^{q,p}$ is endowed with the norm, $\|F\|_{q,p} = \left[E(|F|^p) + \sum_{j=1}^q E(\|D^j F\|_{\mathbf{L}^2([0,1]^j)}^p) \right]^{\frac{1}{p}}$. The space of variable with q derivatives in any $\mathbf{L}^p(\Omega)$ is denoted $\mathbb{D}^{q,\infty} = \cap_{p \geq 1} \mathbb{D}^{q,p}$. These definitions can be extended to random variables with values in any Hilbert space V and the corresponding spaces are denoted $\mathbb{D}^{q,p}(V)$, $\mathbb{D}^{q,\infty}(V)$ (see Section 1.5 in Nualart [21]). In particular the operator D is then well defined from $\mathbb{D}^{q,\infty}$ to $\mathbb{D}^{q-1,\infty}(H)$. Finally, the adjoint operator of D is the Skohorod integral δ , and the Malliavin covariance matrix of an element $F \in \mathbb{D}^{1,\infty}(\mathbb{R}^d)$ is defined as the matrix $\gamma_{F_1, \dots, F_d} = [\langle D.F_i, D.F_j \rangle_H]_{1 \leq i, j \leq d}$.

Now, we consider the variables,

$$U_0^{n,\theta} := \int_0^1 \mathcal{X}_s^{n,\theta} d\mu(s) \quad (10)$$

$$U_1^{n,\theta} := \int_0^1 (\mathcal{X}_{s+1}^{n,\theta} - \mathcal{X}_s^{n,\theta}) d\mu(s) \quad (11)$$

⋮

$$U_{k-1}^{n,\theta} := \int_0^1 (\mathcal{X}_{s+k-1}^{n,\theta} - \mathcal{X}_{s+k-2}^{n,\theta}) d\mu(s) \quad (12)$$

$$U_k^{n,\theta} := \int_0^1 (\mathcal{X}_k^{n,\theta} - \mathcal{X}_{s+k-1}^{n,\theta}) d\mu(s). \quad (13)$$

Note that the joint law of these $k+1$ variables is, by rescaling, the same as the law of the vector composed with variables of the first block \mathbb{B}_0 : $n^{\frac{1}{2}}(\bar{X}_0^\theta - x_0, \bar{X}_1^\theta - \bar{X}_0^\theta, \dots, \bar{X}_{k-1}^\theta - \bar{X}_{k-2}^\theta, X_{\frac{k}{n}}^\theta - \bar{X}_{k-1}^\theta)$. These variables satisfy the following non degeneracy property whose proof is postponed to Section 5.1.3.

Proposition 1. *Under (\mathbf{R}) , $(U_0^{n,\theta}, \dots, U_k^{n,\theta}) \in \mathbb{D}^{3,\infty}$. Denote by $K(\theta)$ the Malliavin covariance matrix of $(U_0^{n,\theta}, \dots, U_k^{n,\theta})$. It is a.s. a invertible matrix and for all $p \geq 1$, we have*

$$E(|\det(K(\theta))|^{-p}) \leq c(p, k).$$

2.2 An exact expression using Malliavin calculus

In this Section we intend to give an exact expression for the score function of the observation of \mathbb{B}_0 or equivalently for the vector $(U_0^{n,\theta}, \dots, U_k^{n,\theta})$ given by (10)–(13).

Under the condition **(R)**, we know that there exists a version of the solution of (6) such that P -almost surely the function $\theta \rightarrow \mathcal{X}_t^\theta$ is continuously differentiable for all t and $\tau_t^{n,\theta} := \frac{\partial \mathcal{X}_t^{n,\theta}}{\partial \theta}$ is solution of the stochastic equation (see Kunita [19]):

$$\begin{aligned} d\tau_t^{n,\theta} &= \frac{\partial a_n}{\partial x}(\mathcal{X}_t^{n,\theta}, \theta) \tau_t^{n,\theta} dW_t + \frac{\partial a_n}{\partial \theta}(\mathcal{X}_t^{n,\theta}, \theta) dW_t + \frac{\partial b_n}{\partial x}(\mathcal{X}_t^{n,\theta}) \tau_t^{n,\theta} dt \quad (14) \\ \tau_0^{n,\theta} &= 0. \end{aligned}$$

The main result of this section is an explicit representation for the derivative of the log-likelihood of one block. This extends a former result given by Gobet (see [9] [10]).

Theorem 4. *The random vector $(U_0^{n,\theta}, \dots, U_k^{n,\theta})$ admits a positive density on \mathbb{R}^{k+1} , denoted by $p_{x_0}(u_0, \dots, u_k, \theta)$. For a.e. (u_0, \dots, u_k) , this density is an absolutely continuous function with respect to the parameter θ and we have the formula:*

$$\frac{\dot{p}_{x_0}}{p_{x_0}}(u_0, \dots, u_k, \theta) = E \left[\delta \left(\sum_{0 \leq j, j' \leq k} \frac{\partial U_j^{n,\theta}}{\partial \theta} K(\theta)_{j,j'}^{-1} D.U_{j'}^{n,\theta} \right) \mid (U_j^{n,\theta} = u_j)_{j=0, \dots, k} \right],$$

where $K(\theta)^{-1}$ is the inverse of the Malliavin covariance matrix of $(U_0^{n,\theta}, \dots, U_k^{n,\theta})$.

Proof. Denote $U^{n,\theta}$ the Wiener functional, $U^{n,\theta} = (U_0^{n,\theta}, \dots, U_k^{n,\theta})$ and let $f : \mathbb{R}^{k+1} \rightarrow \mathbb{R}$ be a smooth function with compact support. Then the function $\theta \mapsto E[f(U^{n,\theta})]$ can be differentiated pointwise and:

$$\frac{\partial}{\partial \theta} E[f(U^{n,\theta})] = E \left[\sum_{j=0}^k \frac{\partial f}{\partial u_j}(U^{n,\theta}) \frac{\partial U_j^{n,\theta}}{\partial \theta} \right].$$

By Proposition 1 the Malliavin covariance matrix of $U^{n,\theta}$ is invertible and a standard computation on Wiener functionals (see formula (2.4) p.81 in Nualart [21]) shows that: $\frac{\partial f}{\partial u_j}(U^{n,\theta}) = \sum_{j'=0}^k \left\langle D(f(U^{n,\theta})), D.U_{j'}^{n,\theta} \right\rangle_H K(\theta)_{j,j'}^{-1}$. It follows that $\frac{\partial}{\partial \theta} E[f(U^{n,\theta})]$ is equal to

$$E \left[\sum_{j=0}^k \sum_{j'=0}^k \left\langle D(f(U^{n,\theta})), D.U_{j'}^{n,\theta} \right\rangle_H K(\theta)_{j,j'}^{-1} \frac{\partial U_j^{n,\theta}}{\partial \theta} \right] = E \left[\left\langle D(f(U^{n,\theta})), L^\theta \right\rangle_H \right]$$

where L^θ is the H -valued random variable:

$$L^\theta := \sum_{j=0}^k \sum_{j'=0}^k \frac{\partial U_j^{n,\theta}}{\partial \theta} K(\theta)_{j,j'}^{-1} D U_{j'}^{n,\theta}. \quad (15)$$

Introducing the δ is the adjoint operator of D , we get

$$\frac{\partial}{\partial \theta} E [f(U^{n,\theta})] = E [f(U^{n,\theta}) \delta(L^\theta)]. \quad (16)$$

Let g be any smooth function with compact support on \mathbb{R} . Using the integration by part formula and the equation (16) we have:

$$\begin{aligned} \int d\theta \dot{g}(\theta) E(f(U^{n,\theta})) &= - \int d\theta g(\theta) \frac{\partial}{\partial \theta} E [f(U^{n,\theta})] = - \int d\theta g(\theta) E [f(U^{n,\theta}) \delta(L^\theta)] \\ &= - \int d\theta g(\theta) E [f(U^{n,\theta}) E[\delta(L^\theta) | (U_0^{n,\theta}, \dots, U_k^{n,\theta})]]. \end{aligned}$$

Introducing the density of the random vector $U^{n,\theta}$ the equation above writes,

$$\begin{aligned} &\int \dot{g}(\theta) d\theta \int f(u_0, \dots, u_k) p_{x_0}(u_0, \dots, u_k, \theta) du_0 \dots du_k \\ &= - \int g(\theta) d\theta \int f(u_0, \dots, u_k) E[\delta(L^\theta) | (U_l^{n,\theta} = u_l)_l] p_{x_0}(u_0, \dots, u_k, \theta) du_0 \dots du_k. \end{aligned}$$

From Fubini's theorem this implies that for all smooth function g we have $du_0 \dots du_k$ -everywhere

$$\begin{aligned} &\int \dot{g}(\theta) p_{x_0}(u_0, \dots, u_k, \theta) d\theta \\ &= - \int g(\theta) E[\delta(L^\theta) | (U_l^{n,\theta} = u_l)_l] p_{x_0}(u_0, \dots, u_k, \theta) d\theta. \quad (17) \end{aligned}$$

Considering a countable sequence of functions dense among smooth compactly supported functions we may obtain that the equality above holds actually for all smooth function g on the same set of full Lebesgue measure. Eventually, observe that (17) is sufficient to imply that the function $\theta \rightarrow p_{x_0}(u_0, \dots, u_k, \theta)$ is absolutely continuous and that

$$\dot{p}_{x_0}(u_0, \dots, u_k, \theta) = E[\delta(L^\theta) | (U_l^{n,\theta} = u_l)_l] p_{x_0}(u_0, \dots, u_k, \theta).$$

Hence the theorem is proved. \square

2.3 A Gaussian approximation for the log-likelihood

In this section we intend to give a tractable approximation for the score function of $(U_0^{n,\theta}, \dots, U_k^{n,\theta})$.

2.3.1 Approximation for the diffusion

We introduce $\tilde{\mathcal{X}}_t^\theta = a(x_0, \theta)W_t$ and $\tilde{\tau}_t^\theta = \dot{a}(x_0, \theta)W_t$ which stand -by (6) and (14)- for the first order approximations of $\mathcal{X}_t^{n,\theta}$ and $\tau_t^{n,\theta} = \frac{\partial \mathcal{X}_t^{n,\theta}}{\partial \theta}$. Then, we consider the quantities obtained by replacing in (10–13) the process \mathcal{X} by this Gaussian approximation:

$$\tilde{U}_0^\theta := a(x_0, \theta) \int_0^1 W_s d\mu(s) = a(x_0, \theta) \int_0^1 \mu([s, 1]) dW_s, \quad (18)$$

$$\begin{aligned} \tilde{U}_j^\theta &:= a(x_0, \theta) \int_0^1 (W_{j+s} - W_{j-1+s}) d\mu(s), \quad \text{for } j = 1, \dots, k-1 \quad (19) \\ &= a(x_0, \theta) \int_{j-1}^j \mu([0, s - (j-1)]) dW_s + a(x_0, \theta) \int_j^{j+1} \mu([s - j, 1]) dW_s, \end{aligned}$$

$$\begin{aligned} \tilde{U}_k^\theta &:= a(x_0, \theta) \int_0^1 (W_k - W_{k-1+s}) d\mu(s), \quad (20) \\ &= a(x_0, \theta) \int_{k-1}^k \mu([0, s - (k-1)]) dW_s. \end{aligned}$$

In the next lemma we control the difference between the $U_j^{n,\theta}$ and their approximation in terms of Sobolev norm.

Lemma 1. *For all $k, p > 1$, there exist constants $c(k, p)$, $c(p)$ such that for all $j \in \{0, \dots, k\}$:*

$$\left\| U_j^{n,\theta} - \tilde{U}_j^\theta \right\|_{2,p} \leq c(k, p) n^{-1/2}, \quad \left\| \tilde{U}_j^\theta \right\|_{3,p} \leq c(p), \quad (21)$$

$$\left\| \frac{\partial U_j^{n,\theta}}{\partial \theta} - \frac{\partial \tilde{U}_j^{n,\theta}}{\partial \theta} \right\|_{2,p} \leq c(k, p) n^{-1/2}, \quad \left\| \frac{\partial \tilde{U}_j^\theta}{\partial \theta} \right\|_{3,p} \leq c(p), \quad (22)$$

$$\forall 0 \leq j, j' \leq k, \quad \left| E \left(U_j^{n,\theta} U_{j'}^{n,\theta} - \tilde{U}_j^\theta \tilde{U}_{j'}^\theta \right) \right| \leq c(k) n^{-1}. \quad (23)$$

Proof. The inequalities on the right hand side of (21)–(22) are immediate by the definition of \tilde{U}_j^θ .

Comparing expressions of (10–13) with (18–20), the two remaining bounds in (21)–(22) will be a consequence of the Minkowski inequality - for the Sobolev norm - and of the control on the diffusions:

$$\sup_{t \leq k} \left\| \mathcal{X}_t^{n,\theta} - \tilde{\mathcal{X}}_t^\theta \right\|_{2,p} + \sup_{t \leq k} \left\| \tau_t^{n,\theta} - \tilde{\tau}_t^\theta \right\|_{2,p} \leq n^{-1/2} c(k, p).$$

We only prove the control on $\mathcal{X}^{n,\theta}$ since the proof for $\tau^{n,\theta}$ is analogous. Recalling (6)–(7), we can write

$$\begin{aligned} \mathcal{X}_t^{n,\theta} - \tilde{\mathcal{X}}_t^\theta &= \int_0^t [a_n(\mathcal{X}_s^{n,\theta}, \theta) - a(x_0, \theta)] dW_s + \int_0^t b_n(\mathcal{X}_s^{n,\theta}) ds \\ &= \frac{1}{\sqrt{n}} \int_0^t \int_0^1 a'_x(x_0 + \frac{u\mathcal{X}_s^{n,\theta}}{\sqrt{n}}, \theta) \mathcal{X}_s^{n,\theta} du dW_s + \frac{1}{\sqrt{n}} \int_0^t b(x_0 + \frac{\mathcal{X}_s^{n,\theta}}{\sqrt{n}}) ds. \quad (24) \end{aligned}$$

But we know [21] that under **(R)** the variables $\mathcal{X}^{n,\theta}$ belongs to $\mathbb{D}^{3,\infty}$ with a control (independent of θ, n): $\sup_{u_1, u_2 \leq s \leq k} E(|D_{u_1, u_2}^2 \mathcal{X}_s^{n,\theta}|^p) \leq c(p, k)$. This is sufficient to deduce $\left\| \mathcal{X}_t^{n,\theta} - \tilde{\mathcal{X}}_t^\theta \right\|_{2,p} \leq n^{-1/2} c(p, k)$ after a few computations.

To obtain (23) note that by (21) it is sufficient to show $E \left((U_j^{n,\theta} - \tilde{U}_j^\theta) \tilde{U}_{j'}^\theta \right) \leq c(k)n^{-1}$. This property will follow again from an analogous relation on the diffusion,

$$\sup_{t, t' \leq k} \left| E \left((\mathcal{X}_t^{n,\theta} - \tilde{\mathcal{X}}_t^\theta) \tilde{\mathcal{X}}_{t'}^\theta \right) \right| \leq c(k)n^{-1}.$$

Indeed, from (24), the above expectation is equal to

$$\begin{aligned} n^{-1/2} \int_0^{t \wedge t'} \int_0^1 E \left[a'_x(x_0 + n^{-1/2} u \mathcal{X}_s^{n,\theta}, \theta) \mathcal{X}_s^{n,\theta} \right] du a(x_0, \theta) ds + \\ n^{-1/2} \int_0^t E \left[b(x_0 + n^{-1/2} \mathcal{X}_s^{n,\theta}) W_{t'} \right] a(x_0, \theta) ds. \end{aligned}$$

Using $\left| E[a'_x(x_0, \theta) \mathcal{X}_s^{n,\theta}] \right| = \left| \int_0^s a'_x(x_0, \theta) E[b_n(\mathcal{X}_u^{n,\theta})] du \right| \leq cn^{-1/2}$, $E[b(x_0)W_{t'}] = 0$ and the boundedness of a''_{xx} and b' , we get the required estimate. \square

2.3.2 Approximation for the log-likelihood

Let us denote the deterministic tridiagonal matrix \tilde{K} of size $(k+1) \times (k+1)$,

$$\tilde{K} = \begin{bmatrix} v_1 & c & 0 & 0 & 0 \\ c & v_1 + v_2 & \ddots & 0 & 0 \\ 0 & \ddots & \ddots & \ddots & 0 \\ 0 & 0 & \ddots & v_1 + v_2 & c \\ 0 & 0 & 0 & c & v_2 \end{bmatrix}, \quad (25)$$

where the entries of the matrix are:

$$v_1 = \int_0^1 \mu([s, 1])^2 ds, \quad v_2 = \int_0^1 \mu([0, s])^2 ds, \quad c = \int_0^1 \mu([0, s]) \mu([s, 1]) ds.$$

It can be easily checked that $a^2(x_0, \theta) \tilde{K}$ is the covariance matrix of the Gaussian vector $(\tilde{U}_0^\theta, \dots, \tilde{U}_k^\theta)$ and is invertible (see Lemma 10 in the Appendix). Now the idea is to introduce the score function that would be produced from the observation of this Gaussian vector. Hence we let:

$$\mathcal{L}_{x_0}(u_0, \dots, u_k, \theta) = \frac{\dot{a}}{a}(x_0, \theta) \left\{ a(x_0, \theta)^{-2} \sum_{0 \leq j, j' \leq k} u_j \tilde{K}_{j, j'}^{-1} u_{j'} - (k+1) \right\}. \quad (26)$$

In this section, we will show that this quantity is an approximation for the true score function $\frac{\dot{p}}{p}$:

Theorem 5. *Let us consider the difference,*

$$\frac{\dot{P}_{x_0}}{p_{x_0}}(u_0, \dots, u_k, \theta) - \mathcal{L}_{x_0}(u_0, \dots, u_k, \theta) := r_{x_0}(u_0, \dots, u_k, \theta). \quad (27)$$

Then we have the following bounds:

$$\left| E \left[r_{x_0}(U_0^{n,\theta}, \dots, U_k^{n,\theta}, \theta) \right] \right| \leq c(k)n^{-1}, \quad (28)$$

$$\forall p \geq 1, \quad E \left[\left| r_{x_0}(U_0^{n,\theta}, \dots, U_k^{n,\theta}, \theta) \right|^p \right]^{\frac{1}{p}} \leq c(k,p)n^{-1/2}. \quad (29)$$

Proof. Keeping in mind the definition of L^θ (see (15)), we introduce its approximation based on the Gaussian quantities defined above:

$$\tilde{L}^\theta := \sum_{j=0}^k \sum_{j'=0}^k \frac{\partial \tilde{U}_j^\theta}{\partial \theta} a(x_0, \theta)^{-2} \tilde{K}_{j,j'}^{-1} D \tilde{U}_{j'}^\theta.$$

The first step is to obtain the following control on the difference $r_1 := L^\theta - \tilde{L}^\theta$:

$$\forall p > 1, \quad \|r_1\|_{\mathbb{D}^{1,p}(H)} \leq c(k,p)n^{-1/2}. \quad (30)$$

Actually, it is a easy consequence of Lemma 1, Proposition 1 and the invertibility of \tilde{K} , noting that the Malliavin covariance matrix of \tilde{U}^θ coincides with the covariance matrix $a^2(x_0, \theta)\tilde{K}$ of the Gaussian vector \tilde{U}^θ . We omit further details.

The second step is to obtain a simple expression for $\delta(\tilde{L}^\theta)$. To see this, we first use the relation for $F \in \mathbb{D}^{1,\infty}$, $u \in \mathbb{D}^{1,\infty}(H)$, $\delta(Fu) = F\delta(u) - \langle D.F, u \rangle_H$ (see [21]):

$$\begin{aligned} \delta(\tilde{L}^\theta) &= \sum_{j=0}^k \sum_{j'=0}^k \frac{\partial \tilde{U}_j^\theta}{\partial \theta} a(x_0, \theta)^{-2} \tilde{K}_{j,j'}^{-1} \delta(D(\tilde{U}_{j'}^\theta)) \\ &\quad - \sum_{j=0}^k \sum_{j'=0}^k a(x_0, \theta)^{-2} \tilde{K}_{j,j'}^{-1} \left\langle D \cdot \frac{\partial \tilde{U}_j^\theta}{\partial \theta}, D \cdot \tilde{U}_{j'}^\theta \right\rangle_H. \end{aligned}$$

On the one hand, $\delta(D(\tilde{U}_{j'}^\theta)) = \tilde{U}_{j'}^\theta$ ($\delta \circ D$ is the identity operator on the first chaos space). On the other hand, one has $\frac{\partial \tilde{U}_j^\theta}{\partial \theta} = \frac{\dot{a}(x_0, \theta)}{a(x_0, \theta)} \tilde{U}_j^\theta$ by (18–20). Hence

$$\begin{aligned} \delta(\tilde{L}^\theta) &= \frac{\dot{a}(x_0, \theta)}{a(x_0, \theta)} \sum_{j=0}^k \sum_{j'=0}^k \tilde{U}_j^\theta a(x_0, \theta)^{-2} \tilde{K}_{j,j'}^{-1} \tilde{U}_{j'}^\theta \\ &\quad - \frac{\dot{a}(x_0, \theta)}{a(x_0, \theta)} \sum_{j=0}^k \sum_{j'=0}^k a(x_0, \theta)^{-2} \tilde{K}_{j,j'}^{-1} \left\langle D \cdot \tilde{U}_j^\theta, D \cdot \tilde{U}_{j'}^\theta \right\rangle_H \\ &= \frac{\dot{a}(x_0, \theta)}{a(x_0, \theta)} \sum_{j=0}^k \sum_{j'=0}^k \tilde{U}_j^\theta a(x_0, \theta)^{-2} \tilde{K}_{j,j'}^{-1} \tilde{U}_{j'}^\theta - \frac{\dot{a}(x_0, \theta)}{a(x_0, \theta)} (k+1). \end{aligned}$$

Now set

$$r_2 = \frac{\dot{a}(x_0, \theta)}{a^3(x_0, \theta)} \sum_{0 \leq j, j' \leq k} \tilde{U}_j^\theta \tilde{K}_{j, j'}^{-1} \tilde{U}_{j'}^\theta - \frac{\dot{a}(x_0, \theta)}{a^3(x_0, \theta)} \sum_{0 \leq j, j' \leq k} U_j^{n, \theta} \tilde{K}_{j, j'}^{-1} U_{j'}^{n, \theta},$$

take the conditional expectation in the equality $\delta(L^\theta) = \delta(\tilde{L}^\theta) + \delta(r_1)$: by Theorem 4, we get (27) with $r_{x_0}(u_0, \dots, u_k, \theta) = E\left(\delta(r_1) \mid (U_j^{n, \theta})_j = (u_j)_j\right) + E\left(r_2 \mid (U_j^{n, \theta})_j = (u_j)_j\right)$.

The final step in the proof is to show that r_{x_0} satisfies conditions (28)–(29). For the first condition, since the Skorohod integral is zero mean, we have $E[r_{x_0}(U_0^{n, \theta}, \dots, U_k^{n, \theta}, \theta)] = E(r_2)$ and we conclude using (23).

We now prove (29). The conditional expectation being a contraction in \mathbf{L}^p it is sufficient to prove

$$E(|\delta(r_1)|^p)^{\frac{1}{p}} \leq c(p, k)n^{-1/2}, \quad E(|r_2|^p)^{\frac{1}{p}} \leq c(p, k)n^{-1/2}.$$

The first estimate follows from (30) and the continuity of the operator δ from $\mathbb{D}^{1,p}(H)$ to \mathbf{L}^p . The second one is an immediate consequence of Lemma 1. \square

Remark 3. *Let us note that the constants $c(k), c(k, p)$ in Theorem 5 should increase as the block length k goes to infinity since the Gaussian approximation ceases to be valid in that case. However in the sequel we shall not need a precise evaluation of this dependence on k since we will have the possibility to conveniently choose the growth rate of $k = k_n$.*

In the following sections we will need this Corollary of Theorem 5.

Corollary 2. *We have for all $p > 1$,*

$$E \left[\left| \frac{\dot{p}_{x_0}}{p_{x_0}}(U_0^{n, \theta}, \dots, U_k^{n, \theta}, \theta) \right|^p \right] \leq c(k, p).$$

Proof. By Theorem 5 it is sufficient to show that $E \left[\left| \mathcal{L}_{x_0}(U_0^{n, \theta}, \dots, U_k^{n, \theta}, \theta) \right|^p \right] \leq c(k, p)$. But from the expression of \mathcal{L}_{x_0} , this estimate is clear. \square

3 Asymptotic study for the augmented model

In this paragraph we establish Theorem 2. Let us recall some notations: we now deal with the diffusion given by (1)–(2); k_n is some integer in $\{1, \dots, n\}$, $L_n = \lfloor n/k_n \rfloor$ and our observation consists of the $L_n + 1$ blocks $\mathbb{B}_0, \dots, \mathbb{B}_{L_n}$ described in Section 1. The length of the block \mathbb{B}_l is $k_{n,l} + 1$, where $k_{n,l} = k_n$ if $l \leq L_n - 1$ and $k_{n,L_n} = n - L_n k_n$. For sake of simplicity in the sequel we sometimes omit the dependence of the size of the block upon n and l , and

let $k_{n,l} = k$ with a slight abuse of notation in particular for the last block of data.

To be able to use the results of the Section 2, we introduce on each block the random variables corresponding to the definitions (10–13) for the first block. Hence for $l \in \{0, \dots, L_n\}$, we define the $k_{n,l}$ following variables:

$$\begin{aligned} U_{0,l} &= n^{\frac{1}{2}}(\bar{X}_{kl} - X_{\frac{kl}{n}}), \\ U_{1,l} &= n^{\frac{1}{2}}(\bar{X}_{kl+1} - \bar{X}_{kl}), \\ &\vdots \\ U_{k-1,l} &= n^{\frac{1}{2}}(\bar{X}_{kl+k-1} - \bar{X}_{kl+k-2}), \\ U_{k,l} &= n^{\frac{1}{2}}(X_{\frac{k(l+1)}{n}} - \bar{X}_{kl+k-1}). \end{aligned}$$

Clearly the observation of the $(U_{j,l})$ for $l \in \{0, \dots, L_n\}, j \in \{0, \dots, k_{n,l}\}$ is equivalent to the observation of the $L_n + 1$ blocks. Using the Markov property for the process X it appears that the law of the vector $(U_{j,l})_{j=0, \dots, k_{n,l}}$ conditionally to all the variables $U_{j,l'}$ with $l' < l, j \in \{0, \dots, k_{n,l'}\}$ is the same as conditionally to $X_{kl/n}$ only; moreover this law - conditionally to $X_{kl/n} = x_0$ - coincides with that of the vector $(U_0^{n,\theta}, \dots, U_k^{n,\theta})$ studied in Section 2. Thus it admits the density $p_{X_{\frac{kl}{n}}}(u_0, \dots, u_k, \theta)$ studied in Sections 2.2–2.3. Hence the log-likelihood of the augmented model admits the additive structure:

$$\begin{aligned} \ln(Z_{\theta_0, \theta_0 + u_n h}^{n, \text{aug}}(\mathcal{O}^{n, \text{aug}})) &= \sum_{l=0}^{L_n} \ln \frac{p_{X_{\frac{kl}{n}}}(U_{0,l}, \dots, U_{k,l}, \theta_0 + u_n h)}{p_{X_{\frac{kl}{n}}}(U_{0,l}, \dots, U_{k,l}, \theta_0)} \\ &= \sum_{l=0}^{L_n} \int_{\theta_0}^{\theta_0 + u_n h} \frac{\dot{p}_{X_{\frac{kl}{n}}}(U_{0,l}, \dots, U_{k,l}, s)}{p_{X_{\frac{kl}{n}}}(U_{0,l}, \dots, U_{k,l}, s)} ds. \end{aligned}$$

Owing to Theorem 5, we deduce the decomposition

$$\begin{aligned} \ln(Z_{\theta_0, \theta_0 + u_n h}^{n, \text{aug}}(\mathcal{O}^{n, \text{aug}})) &= \sum_{l=0}^{L_n} \int_{\theta_0}^{\theta_0 + u_n h} \mathcal{L}_{X_{\frac{kl}{n}}}(U_{0,l}, \dots, U_{k,l}, s) ds \\ &\quad + \sum_{l=0}^{L_n} \int_{\theta_0}^{\theta_0 + u_n h} r_{X_{\frac{kl}{n}}}(U_{0,l}, \dots, U_{k,l}, s) ds. \end{aligned}$$

In the above decomposition, we will show in Sections 3.1–3.2 that the explicit term involving \mathcal{L}_{x_0} governs the asymptotic behavior of the log-likelihood ratio; the other term does not contribute in the limit.

3.1 Proof of Theorem 2: the explicit term

Let us introduce a slight modification of \mathcal{L}_{x_0} , which has the advantage of being a smoother function w.r.t. θ :

$$\xi_{l,n}(\theta) = \frac{\dot{a}}{a}(x_0, \theta_0) \left\{ a(x_0, \theta)^{-2} \sum_{0 \leq j, j' \leq k} U_{j,l} \tilde{K}_{j,j'}^{-1} U_{j',l} - (k+1) \right\}, \quad (31)$$

and we set $N_n^{\text{aug}} = u_n \sum_{l=0}^{L_n} \xi_{l,n}(\theta_0)$ and $I_n^{\text{aug}} = u_n^2 \sum_{l=0}^{L_n} \frac{\partial \xi_{l,n}}{\partial \theta}(\theta_0)$.

Proposition 2. *If $k_n \rightarrow \infty$ slowly enough,*

$$\sum_{l=0}^{L_n} \int_{\theta_0}^{\theta_0 + u_n h} \mathcal{L}_{X_{\frac{kl}{n}}}(U_{0,l}, \dots, U_{k,l}, s) ds = h N_n^{\text{aug}} + \frac{h^2}{2} I_n^{\text{aug}} + R_n, \quad (32)$$

where $I_n^{\text{aug}} \xrightarrow[n \rightarrow \infty]{\mathbb{P}^{\theta_0}} -\mathcal{I}_{\theta_0}$, $R_n \xrightarrow[n \rightarrow \infty]{\mathbb{P}^{\theta_0}} 0$ and there exists an extra random variable $N \sim \mathcal{N}(0, 1)$ independent of \mathcal{I}_{θ_0} such that, N_n^{aug} converges stably in law under \mathbb{P}^{θ_0} to $N\sqrt{\mathcal{I}_{\theta_0}}$.

Proof. Comparing (26) with the definition of $\xi_{l,n}(\theta)$ above and using a Taylor expansion for $\xi_{l,n}(\theta)$ around θ_0 , we get the equation (32) with a remainder term $R_n = R_n^{(1)} + R_n^{(2)}$ satisfying:

$$R_n^{(1)} = \sum_{l=0}^{L_n} \int_{\theta_0}^{\theta_0 + u_n h} \left[\frac{\dot{a}}{a}(X_{\frac{kl}{n}}, s) - \frac{\dot{a}}{a}(X_{\frac{kl}{n}}, \theta_0) \right] \left\{ \sum_{0 \leq j, j' \leq k} \frac{U_{j,l} \tilde{K}_{j,j'}^{-1} U_{j',l}}{a(X_{\frac{kl}{n}}, s)^2} - (k+1) \right\} ds, \quad (33)$$

$$\left| R_n^{(2)} \right| \leq c \sum_{l=0}^{L_n} u_n^{2+\gamma} \left\{ \sum_{0 \leq j, j' \leq k} \left| U_{j,l} \tilde{K}_{j,j'}^{-1} U_{j',l} \right| \right\} \quad (34)$$

(for $R_n^{(2)}$ we have used that $\theta \mapsto \dot{a}(x, \theta)$ is γ -Hölder continuous). To complete the proof, we repeatedly use the following classical convergence result about triangular arrays of random variables.

Lemma 2 (Genon-Catalot and Jacod [5], Lemma 9). *Let $(\chi_l^n)_{0 \leq l \leq L_n}$, U be random variables, with χ_l^n being \mathcal{F}_{l+1}^n -measurable. The two following conditions imply $\sum_{l=0}^{L_n} \chi_l^n \xrightarrow{\mathbb{P}} U$:*

$$\sum_{l=0}^{L_n} E[\chi_l^n | \mathcal{F}_l^n] \xrightarrow{\mathbb{P}} U \quad \text{and} \quad \sum_{l=0}^{L_n} E[(\chi_l^n)^2 | \mathcal{F}_l^n] \xrightarrow{\mathbb{P}} 0.$$

• We first focus on N_n^{aug} . Let us introduce the sigma field $\mathcal{F}_l^n = \sigma(X_0; B_s, s \leq \frac{kl}{n})$ for $l = 0, \dots, L_n$ and $\mathcal{F}_{L_n+1}^n = \sigma(X_0; B_s, s \leq 1)$. Then the variable $\xi_{l,n}(\theta_0)$ is \mathcal{F}_{l+1}^n -measurable and the asymptotic behavior of N_n^{aug} will follow from Lemma 2. To make clearer this point we introduce the following approximation based on conditionally Gaussian variables:

$$\tilde{\xi}_{l,n}(\theta) = \frac{\dot{a}}{a}(X_{\frac{kl}{n}}, \theta) \left\{ a(X_{\frac{kl}{n}}, \theta)^{-2} \sum_{0 \leq j, j' \leq k} \tilde{U}_{j,l} \tilde{K}_{j,j'}^{-1} \tilde{U}_{j',l} - (k+1) \right\}. \quad (35)$$

Here, $\tilde{U}_{j,l}$ is the Gaussian approximation under \mathbb{P}^θ of $U_{j,l}$ corresponding on the block \mathbb{B}_l to the variables (18)–(20) on the block \mathbb{B}_0 :

$$\begin{aligned} \tilde{U}_{0,l} &:= a(X_{\frac{kl}{n}}, \theta) n^{\frac{1}{2}} \int_0^1 (B_{\frac{kl+s}{n}} - B_{\frac{kl}{n}}) d\mu(s), \\ \tilde{U}_{j,l} &:= a(X_{\frac{kl}{n}}, \theta) n^{\frac{1}{2}} \int_0^1 (B_{\frac{kl+j+s}{n}} - B_{\frac{kl+j-1+s}{n}}) d\mu(s) \quad \text{for } j = 1, \dots, k-1, \\ \tilde{U}_{k,l} &:= a(X_{\frac{kl}{n}}, \theta) n^{\frac{1}{2}} \int_0^1 (B_{\frac{k(l+1)}{n}} - B_{\frac{kl+k-1+s}{n}}) d\mu(s). \end{aligned}$$

Observe that this vector $(\tilde{U}_{j,l})_{j=0, \dots, k}$ has, under \mathbb{P}^θ and conditionally to $X_{\frac{kl}{n}} = x_0$, the same law as the vector $(\tilde{U}_j^\theta)_{j=0, \dots, k}$ defined in Section 2.3. Thus its conditional law is Gaussian with covariance matrix $a(X_{\frac{kl}{n}}, \theta)^2 \tilde{K}$ (by Lemma 10). Hence, the variable $\tilde{\xi}_{l,n}(\theta_0)$ is \mathcal{F}_{l+1}^n -measurable and under \mathbb{P}^{θ_0} , it is conditionally (to $X_{\frac{kl}{n}}$) distributed as a recentered $\chi^2(k+1)$ variable. Thus we deduce the four following properties:

- 1) $u_n \sum_{l=0}^{L_n} E_{\theta_0} [\tilde{\xi}_{l,n}(\theta_0) \mid \mathcal{F}_l^n] = 0$;
- 2) Using $u_n^2 = 1/n$, $L_n \sim n/k_n \rightarrow \infty$ and $k_n \rightarrow \infty$, one has

$$\begin{aligned} u_n^2 \sum_{l=0}^{L_n} E_{\theta_0} [(\tilde{\xi}_{l,n}(\theta_0))^2 \mid \mathcal{F}_l^n] &= u_n^2 \sum_{l=0}^{L_n} 2(k_n+1) \left(\frac{\dot{a}}{a}\right)^2 (X_{\frac{kl}{n}}, \theta_0) \\ &= \frac{2(k_n+1)}{k_n} \int_0^1 \left(\frac{\dot{a}}{a}\right)^2 (X_s, \theta_0) ds + o_{\mathbb{P}^{\theta_0}}(1) \quad (36) \\ &\xrightarrow{\mathbb{P}^{\theta_0}} 2 \int_0^1 \left(\frac{\dot{a}}{a}\right)^2 (X_s, \theta_0) ds = \mathcal{I}_{\theta_0}; \end{aligned}$$

- 3) $u_n^4 \sum_{l=0}^{L_n} E_{\theta_0} [|\tilde{\xi}_{l,n}(\theta_0)|^4 \mid \mathcal{F}_l^n] \leq cn^{-2} L_n k_n^4 \leq cn^{-1} k_n^3 \rightarrow 0$, if k_n goes to ∞ slowly enough;
- 4) $u_n \sum_{l=0}^{L_n} E_{\theta_0} [\tilde{\xi}_{l,n}(\theta_0) [B_{\frac{(k+1)l}{n}} - B_{\frac{kl}{n}}] \mid \mathcal{F}_l^n] = 0$.

From these four properties, it follows (see Jacod [14]) that $u_n \sum_{l=0}^{L_n} \tilde{\xi}_{l,n}(\theta_0)$ converges stably under \mathbb{P}^{θ_0} to a mixed Gaussian variable as in the statement of the proposition. To obtain the limit for N_n^{aug} , it is sufficient to prove that

$$N_n^{\text{aug}} - u_n \sum_{l=0}^{L_n} \tilde{\xi}_{l,n}(\theta_0) \xrightarrow{\mathbb{P}^{\theta_0}} 0. \quad (37)$$

Due to Lemma 2 a sufficient condition consists in the two following points:

- 1) $u_n \sum_{l=0}^{L_n} E_{\theta_0} \left[\tilde{\xi}_{l,n}(\theta_0) - \xi_{l,n}(\theta_0) \mid \mathcal{F}_l^n \right] \rightarrow 0$ in probability;
- 2) $u_n^2 \sum_{l=0}^{L_n} E_{\theta_0} \left[(\tilde{\xi}_{l,n}(\theta_0) - \xi_{l,n}(\theta_0))^2 \mid \mathcal{F}_l^n \right] \rightarrow 0$ in probability.

But these two points can be shown using (21) and (23) of Lemma 1 (for k_n slowly increasing).

• We now study I_n^{aug} . A direct differentiation of $\xi_{l,n}(\theta)$ (recall (31)) gives

$$\dot{\xi}_{l,n}(\theta) = \frac{\dot{a}}{a}(X_{\frac{kl}{n}}, \theta) \frac{-2\dot{a}}{a^3}(X_{\frac{kl}{n}}, \theta) \sum_{0 \leq j, j' \leq k} U_{l,j} \tilde{K}_{j,j'}^{-1} U_{l,j'}.$$

Then, with a few computations similar to the study of N_n^{aug} , we obtain (for appropriate k_n):

$$\begin{aligned} 1) \quad u_n^2 \sum_{l=0}^{L_n} E_{\theta_0} \left[\dot{\xi}_{l,n}(\theta_0) \mid \mathcal{F}_l^n \right] &= u_n^2 \sum_{l=0}^{L_n} -2(k_{n,l}+1) \frac{(\dot{a})^2}{a^2}(X_{\frac{kl}{n}}, \theta_0) + O_{\mathbb{P}^{\theta_0}}\left(\frac{c(k_n)}{\sqrt{n}}\right) \\ &= -2 \frac{(k_n+1)}{k_n} \int_0^1 \left(\frac{\dot{a}}{a}(X_s, \theta_0) \right)^2 ds + o_{\mathbb{P}^{\theta_0}}(1) \\ &\xrightarrow{\mathbb{P}^{\theta_0}} -\mathcal{I}_{\theta_0}; \end{aligned} \quad (38)$$

$$2) \quad u_n^4 \sum_{l=0}^{L_n} E_{\theta_0} \left[[\dot{\xi}_{l,n}(\theta_0)]^2 \mid \mathcal{F}_l^n \right] \leq cn^{-1}k_n^4 \rightarrow 0.$$

Combined with Lemma 2, these two convergences imply that of I_n^{aug} to $-\mathcal{I}_{\theta_0}$ under \mathbb{P}^{θ_0} .

• The remainder term R_n . Firstly concerning $R_n^{(2)}$, a direct use of (21) gives $E(|R_n|) \leq c(k_n)n^{-\gamma/2} \rightarrow 0$ if k_n slowly goes to ∞ . Secondly the convergence to zero of $R_n^{(1)} = \sum_{l=0}^{L_n} R_{n,l}^{(1)}$ is more delicate and Lemma 2 is helpful for this. To this end we evaluate the conditional expectation of $R_{n,l}^{(1)}$ using (23) and the fact the $(\tilde{U}_{j,l})_j$ have the conditional covariance matrix $a(X_{\frac{kl}{n}}, \theta)^2 \tilde{K}$:

$$\begin{aligned} E_{\theta_0}[R_{n,l}^{(1)} \mid \mathcal{F}_l^n] &= \int_{\theta_0}^{\theta_0+u_n h} \left[\frac{\dot{a}}{a}(X_{\frac{kl}{n}}, s) - \frac{\dot{a}}{a}(X_{\frac{kl}{n}}, \theta_0) \right] \left\{ \frac{a(X_{\frac{kl}{n}}, \theta_0)^2}{a(X_{\frac{kl}{n}}, s)^2} - 1 \right\} (k_n+1) ds \\ &\quad + O(n^{-1}u_n c(k_n)). \end{aligned}$$

The function a being $\mathcal{C}^{1+\gamma}$ in θ , one gets: $\sum_{l=0}^{L_n} |E_{\theta_0}[R_{n,l}^{(1)} | \mathcal{F}_l^n]| \leq cn^{-\gamma/2} + \frac{c(k_n)}{k_n} n^{-1/2} \rightarrow 0$ for appropriate k_n . With similar considerations we evaluate the second conditional moment and obtain $u_n^2 \sum_{l=0}^{L_n} E_{\theta_0}[(R_n^{(1)})^2 | \mathcal{F}_l^n] \leq c(k_n)L_n u_n^{2+2\gamma} \xrightarrow{n \rightarrow \infty} 0$. \square

3.2 Proof of Theorem 2: the negligible terms

It remains to prove that, as announced, there is convergence to zero of $\sum_{l=0}^{L_n} \eta_l$ with $\eta_l = \int_{\theta_0}^{\theta_0+u_n h} r_{X_{\frac{kl}{n}}}(U_{0,l}, \dots, U_{k,l}, s) ds$. We aim at applying Lemma 2 by computing the first two conditional moments of η_l under \mathbb{P}^{θ_0} . The main difficulty here comes from the fact that we do not have an explicit expression for $r_{x_0}((u_j)_j, \theta)$. Indeed by Theorem 5 we know bounds for the moments $E_{\theta, x_0}^n (|r_{x_0}((U_j)_j, \theta)|^p)$ where by $E_{x_0, \theta}^n$ we denote the expectation with respect to the law of $\mathcal{X}^{n, \theta}$ solution of (6). This is a priori insufficient to compute the conditional moments of η_l under \mathbb{P}^{θ_0} which involves quantities such as $E_{\theta_0, x_0}^n (|r_{x_0}((U_j)_j, s)|^p)$ for $s \neq \theta_0$. Thus in Lemmas 7-8 in the Appendix we study the transformation of such moments under change of measure.

Firstly, we evaluate the conditional expectation of η_l ,

$$E_{\theta_0}[\eta_l | \mathcal{F}_l^n] = \int_{\theta_0}^{\theta_0+u_n h} E_{\theta_0, x}^n[r_x((U_j)_j, s)]|_{x=X_{\frac{kl}{n}}} ds.$$

But $|E_{\theta_0, x}^n[r_x((U_j)_j, s)]| \leq |E_{s, x}^n[r_x((U_j)_j, s)]| + |E_{\theta_0, x}^n[r_x((U_j)_j, s)] - E_{s, x}^n[r_x((U_j)_j, s)]|$ can be bounded using (28) and Lemma 8 in the Appendix by $c(k)n^{-1} + |s - \theta_0| E_{s, x}^n[|r_x((U_j)_j, s)|^\alpha]^{\frac{1}{\alpha}}$ for some constant $\alpha \geq 1$. Then by (29) we deduce $|E[\eta_l | \mathcal{F}_l^n]| \leq c(k_n)[u_n n^{-1} + u_n^2 n^{-1/2}]$. Finally a block length k_n slowly increasing guarantees $\sum_{l=0}^{L_n} E_{\theta_0}[\eta_l | \mathcal{F}_l^n] \xrightarrow{\mathbb{P}^{\theta_0}} 0$.

Secondly and similarly, owing to Theorem 5 and Lemma 7 in the Appendix, we get $E[\eta_l^2 | \mathcal{F}_l^n] \leq c(k_n)u_n^2 n^{-1} \rightarrow 0$. Therefore, by Lemma 2, we have proved $\sum_{l=0}^{L_n} \eta_l \xrightarrow{\mathbb{P}^{\theta_0}} 0$. This ends the proof of Theorem 2.

3.3 What happens if k_n remains fixed?

If we assume now that $k_n = k \in \mathbb{N}$ remains fixed as $n \rightarrow \infty$, the number of data $(X_{\frac{kl}{n}})_{l=0, \dots, L_n}$ added to the model in order to force an additive structure to the log-likelihood is not negligible compared to the number of initial data. Hence the statistical properties of the augmented model shall depend on k and thus differ from the statistical properties of the initial model given in Theorem 1. Actually we have the following LAMN property for the augmented model in that case.

Theorem 6. *If a sequence $k_n = k$ is fixed, then the augmented model satisfies the LAMN property with a conditional variance equal to:*

$$\mathcal{I}_{k,\theta_0} = 2 \left(\frac{k+1}{k} \right) \int_0^1 \left(\frac{\dot{a}}{a} \right)^2 (X_s, \theta_0) ds.$$

Proof. The proof is essentially the same as that of Theorem 2, the difference in the asymptotic information comes from the difference in the limit of the quantities (36)–(38) when k is fixed. \square

As expected, the conditional information is greater by the a factor $(k+1)/k$, due to the non negligibility of the added observation. Actually this factor should be read as $1 + \frac{1}{k}$, meaning that an addition of $\frac{1}{k}\%$ of datas increases the information in the same way. Local means and values at discrete points are not redundant (as expected from the multiplicative Brownian case, see Example 1) and moreover, they bring an equal information. Considering $k = 1$ is interesting, since we observe then on each block $[i/n, (i+1)/n]$ both the exact value $X_{\frac{i}{n}}$ and a mean \bar{X}_i . It appears that the asymptotic information is then twice the information given by the observation of only the exact values (or only the means).

4 LAMN property for the initial model

In this Section we are back to the model where the observation is only $\mathcal{O}^n = (\bar{X}_j)_{j=0,\dots,n-1}$ and we will prove Theorem 1 by relying on the LAMN property for the augmented model.

A first intermediate result is that on can approximate the log-likelihood of the augmented model by a function of the observation \mathcal{O}^n .

Proposition 3. *There exist Γ_n random variables measurable with respect to \mathcal{O}^n such that:*

$$\ln(Z_{\theta_0, \theta_0 + u_n h}^{n, \text{aug}}(\mathcal{O}^{n, \text{aug}})) - \Gamma_n \xrightarrow[\mathbb{P}^{\theta_0}]{n \rightarrow \infty} 0.$$

Proof. We have seen in Section 3 that $\ln(Z_{\theta_0, \theta_0 + u_n h}^{n, \text{aug}}(\mathcal{O}^{n, \text{aug}})) = hN_n^{\text{aug}} + 1/2h^2 I_n^{\text{aug}} + o_{\mathbb{P}^{\theta_0}}(1)$ where the quantities N_n^{aug} and I_n^{aug} were defined in Section 3.1.

Thus the proof of the proposition consists in introducing a proper modification of these quantities which only depends on the observations. We let for $l = 0, \dots, k_n$

$$\xi_{l,n}^{\text{obs}}(\theta) = \frac{\dot{a}}{a}(\bar{X}_{kl-1}, \theta_0) \left\{ a^{-2}(\bar{X}_{kl-1}, \theta) \sum_{1 \leq j, j' \leq k-1} U_{l,j} \widehat{K}_{j,j'}^{-1} U_{l,j'} - (k-1) \right\},$$

with the convention $\bar{X}_{-1} = \xi_0$ is the known initial value of the diffusion and the matrix $a^2(x_0, \theta) \widehat{K}$ is the covariance matrix of the conditionally Gaussian vector $(\tilde{U}_1^\theta, \dots, \tilde{U}_{k-1}^\theta)$:

$$\widehat{K} = \begin{bmatrix} v_1 + v_2 & c & 0 & 0 \\ c & \ddots & \ddots & 0 \\ 0 & \ddots & \ddots & c \\ 0 & 0 & c & v_1 + v_2 \end{bmatrix}.$$

Clearly, $\xi_{l,n}^{\text{obs}}(\theta)$ only depends on the observation \mathcal{O}^n since we have suppressed all occurrences of the variables $U_{0,l}$ and $U_{k,l}$ and replaced $X_{\frac{kl}{n}}$ by \bar{X}_{kl-1} from the expression of $\xi_{l,n}(\theta)$ (compare with (31)). Then we let $N_n^{\text{obs}} = u_n \sum_{l=0}^{L_n} \xi_{l,n}^{\text{obs}}(\theta_0)$ and $I_n^{\text{obs}} = u_n^2 \sum_{l=0}^{L_n} \frac{\partial \xi_{l,n}^{\text{obs}}}{\partial \theta}(\theta_0)$.

• Study of $N_n^{\text{aug}} - N_n^{\text{obs}}$. The first step is to consider the conditionally re-centered chi square approximation of $\xi_{l,n}^{\text{obs}}(\theta)$ that we define as:

$$\tilde{\xi}_{l,n}^{\text{obs}}(\theta) = \frac{\dot{a}}{a} \left(X_{\frac{kl}{n}}, \theta_0 \right) \left\{ a^{-2} \left(X_{\frac{kl}{n}}, \theta \right) \sum_{1 \leq j, j' \leq k-1} \tilde{U}_{l,j} \widehat{K}_{j,j'}^{-1} \tilde{U}_{l,j'} - (k-1) \right\}. \quad (39)$$

The first step is to prove the validity of the approximation:

$$u_n \sum_{l=0}^{L_n} \{ \xi_{l,n}^{\text{obs}}(\theta_0) - \tilde{\xi}_{l,n}^{\text{obs}}(\theta_0) \} \xrightarrow{\mathbb{P}^{\theta_0}} 0. \quad (40)$$

This is done similarly to the proof of $N_n^{\text{aug}} - u_n \sum_{l=0}^{L_n} \tilde{\xi}_{l,n}(\theta_0) \rightarrow 0$ in proposition 2, by considering the first two conditional moments, but here the first moment is more delicate to handle: the conditional moment $E_{\theta_0}[\xi_{l,n}^{\text{obs}}(\theta_0) - \tilde{\xi}_{l,n}^{\text{obs}}(\theta_0) \mid \mathcal{F}_l^n]$ is of the form $(k-1)\{g(X_{\frac{kl}{n}}) - g(\bar{X}_{kl-1})\}h(\bar{X}_{kl-1}) + O(c(k_n)/n)$ for g and h two \mathcal{C}^2 functions. If we abruptly use the relation $\left\| X_{\frac{kl}{n}} - \bar{X}_{kl-1} \right\|_{\mathbf{L}^p} \leq c(p)n^{-1/2}$ then we only deduce that $u_n \sum_{l=0}^{L_n} E_{\theta_0}[\xi_{l,n}^{\text{obs}}(\theta_0) - \tilde{\xi}_{l,n}^{\text{obs}}(\theta_0) \mid \mathcal{F}_l^n]$ remains bounded in probability. To show that it actually converges to zero, we have to apply again Lemma 2 to the new triangular array of variables, $u_n \sum_{l=0}^{L_n} (k-1)\{g(X_{\frac{kl}{n}}) - g(\bar{X}_{kl-1})\}h(\bar{X}_{kl-1})$. Then by rather long computations, using that $\left\| \bar{X}_{kl-1} - \bar{X}_{kl-2} \right\|_{\mathbf{L}^p} \leq c(p)(k/n)^{1/2}$ and $\left| E_{\theta_0}[X_{\frac{kl}{n}} - \bar{X}_{kl-1} \mid \mathcal{F}_{l-1}^n] \right| \leq cn^{-1}$ by the centering property we can prove,

$$u_n \sum_{l=0}^{L_n} (k-1) \left| E_{\theta_0}[\{g(X_{\frac{kl}{n}}) - g(\bar{X}_{kl-1})\}h(\bar{X}_{kl-1}) \mid \mathcal{F}_{l-1}^n] \right| \leq c(k)n^{-1/2} \xrightarrow{\mathbb{P}^{\theta_0}} 0,$$

$$u_n^2 \sum_{l=0}^{L_n} (k-1)^2 E_{\theta_0}[\{g(X_{\frac{kl}{n}}) - g(\bar{X}_{kl-1})\}^2 h(\bar{X}_{kl-1})^2 \mid \mathcal{F}_{l-1}^n] \leq c(k)n^{-1} \xrightarrow{\mathbb{P}^{\theta_0}} 0.$$

Thus we deduce $u_n \sum_{l=0}^{L_n} E_{\theta_0} [\xi_{l,n}^{\text{obs}}(\theta_0) - \tilde{\xi}_{l,n}^{\text{obs}}(\theta_0) \mid \mathcal{F}_l^n] \rightarrow 0$. The second condition $u_n^2 \sum_{l=0}^{L_n} E_{\theta_0} [(\xi_{l,n}^{\text{obs}}(\theta_0) - \tilde{\xi}_{l,n}^{\text{obs}}(\theta_0))^2 \mid \mathcal{F}_l^n] \leq c(k)n^{-1} \rightarrow 0$ is easily obtained and we deduce (40).

Thus, in view of the equation (37), it remains to prove that $u_n \sum_{l=0}^{L_n} \{\tilde{\xi}_{l,n}^{\text{obs}}(\theta_0) - \tilde{\xi}_{l,n}(\theta_0)\}$ is negligible. But by Lemma 11 in the Appendix, comparing expressions (35) and (39), it appears that conditionally to \mathcal{F}_l^n the random variable $\tilde{\xi}_{l,n}^{\text{obs}}(\theta_0) - \tilde{\xi}_{l,n}(\theta_0)$ is a recentered $\chi^2(2)$ variable and hence the following properties hold:

$$u_n \sum_{l=0}^{L_n} E_{\theta_0} \left(\tilde{\xi}_{l,n}^{\text{obs}}(\theta_0) - \tilde{\xi}_{l,n}(\theta_0) \mid \mathcal{F}_l^n \right) = 0,$$

$$u_n^2 \sum_{l=0}^{L_n} E_{\theta_0} \left(\left\{ \tilde{\xi}_{l,n}^{\text{obs}}(\theta_0) - \tilde{\xi}_{l,n}(\theta_0) \right\}^2 \mid \mathcal{F}_l^n \right) = \sum_{l=0}^{L_n} u_n^2 4 \frac{\dot{a}^2}{a^2} (X_{\frac{kl}{n}}, \theta_0) \leq \frac{c}{k_n} \rightarrow 0.$$

These two properties imply by Lemma 2 the convergence to 0 under \mathbb{P}^{θ_0} of $u_n \sum_{l=0}^{L_n} \left\{ \tilde{\xi}_{l,n}^{\text{obs}}(\theta_0) - \tilde{\xi}_{l,n}(\theta_0) \right\}$, and thus $N_n^{\text{aug}} - N_n^{\text{obs}} \xrightarrow{\mathbb{P}^{\theta_0}} 0$.

• Study of $I_n^{\text{aug}} - I_n^{\text{obs}}$. Exactly as we proved that I_n^{aug} tends to $-\mathcal{I}_{\theta_0}$ we can show that $I_n^{\text{obs}} \rightarrow -\mathcal{I}_{\theta_0}$. Thus the difference is negligible.

Finally the proposition is obtained by setting $\Gamma_n = hN_n^{\text{obs}} + h^2/2I_n^{\text{obs}}$. \square

Then the Theorem 1 is a consequence of the following proposition combined with Proposition 3 and Theorem 2.

Proposition 4. *We have the convergence,*

$$Z_{\theta_0, \theta_0 + u_n h}^n - e^{\Gamma_n} \xrightarrow[n \rightarrow \infty]{\mathbb{P}^{\theta_0}} 0.$$

Proof. The starting point is the relation between the likelihood of the initial and of the augmented model: $Z_{\theta_0, \theta_0 + u_n h}^n = E_{\theta_0} \left[Z_{\theta_0, \theta_0 + u_n h}^{n, \text{aug}} \mid \mathcal{O}^n \right]$. By Proposition 3 we can write $Z_{\theta_0, \theta_0 + u_n h}^{n, \text{aug}} = e^{\Gamma_n} e^{\varepsilon_n}$ where ε_n tends to zero in \mathbb{P}_{θ_0} probability. Using that Γ_n is \mathcal{O}^n measurable we deduce,

$$Z_{\theta_0, \theta_0 + u_n h}^n - e^{\Gamma_n} = E_{\theta_0} \left[e^{\Gamma_n} (e^{\varepsilon_n} - 1) \mid \mathcal{O}^n \right].$$

We now use the inequality $|e^u - 1| \leq (|u| \wedge 1)(e^u + 1)$ to obtain that $\left| Z_{\theta_0, \theta_0 + u_n h}^n - e^{\Gamma_n} \right| \leq \alpha_n + \beta_n$ with:

$$\alpha_n = E_{\theta_0} \left[(|\varepsilon_n| \wedge 1) e^{\Gamma_n} \mid \mathcal{O}^n \right] = E_{\theta_0} \left[(|\varepsilon_n| \wedge 1) \mid \mathcal{O}^n \right] e^{\Gamma_n},$$

$$\beta_n = E_{\theta_0} \left[(|\varepsilon_n| \wedge 1) e^{\Gamma_n} e^{\varepsilon_n} \mid \mathcal{O}^n \right] = E_{\theta_0} \left[(|\varepsilon_n| \wedge 1) Z_{\theta_0, \theta_0 + u_n h}^{n, \text{aug}} \mid \mathcal{O}^n \right].$$

It now remains to show the convergence to zero of α_n and β_n .

For α_n , let us notice that $(e^{\Gamma_n})_n$ is a tight sequence and that $E_{\theta_0} [|\varepsilon_n| \wedge 1 \mid \mathcal{O}^n]$ converges in $\mathbf{L}^1(\mathbb{P}_{\theta_0})$ norm to zero since,

$$E_{\theta_0} [E_{\theta_0} [|\varepsilon_n| \wedge 1 \mid \mathcal{O}^n]] = E_{\theta_0} [|\varepsilon_n| \wedge 1] \xrightarrow{n \rightarrow \infty} 0.$$

For β_n , we have $E_{\theta_0}[\beta_n] = E_{\theta_0+u_n h}[|\varepsilon_n| \wedge 1]$. But the sequence of probabilities \mathbb{P}_{θ_0} and $\mathbb{P}_{\theta_0+u_n h}$ restricted to the sigma fields $\mathcal{O}^{n, \text{aug}}$ are contiguous (this is a consequence of the LAMN property for the augmented model, see for e.g. Proposition 1 in Jeganathan [15]); hence the sequence $(\varepsilon_n)_n$ which is measurable with respect to $\mathcal{O}^{n, \text{aug}}$ and converges to zero in \mathbb{P}_{θ_0} -probability converges also in $\mathbb{P}_{\theta_0+u_n h}$ -probability. This implies $E_{\theta_0}[\beta_n] = E_{\theta_0+u_n h}[|\varepsilon_n| \wedge 1] \rightarrow 0$. \square

5 Appendix

5.1 Proof of results of Section 2.1

Since the results of Section 2.1 concern only the study of a density for fixed values of θ , we omit the dependence upon θ in our notations. We will prove the results in the following order. First in section 5.1.1, we show that the law of the Wiener functional $(U^n, V^n) = \left(\int_0^1 \mathcal{X}_s^n d\mu(s), \mathcal{X}_1^n \right)$ admits a density. Then we prove the lower and upper bounds given in Theorem 3 (section 5.1.2) and eventually we deduce the Proposition 1 (section 5.1.3).

5.1.1 Existence of the density $\mathbf{p}_{x_0}^n$

We know [21] that under (\mathbf{R}) the random variable \mathcal{X}_t^n is an element of $\mathbb{D}^{3, \infty}$ and its first derivative is equal to

$$D_t \mathcal{X}_s^n = \mathbf{1}_{\{t \leq s\}} \mathcal{Y}_s^n (\mathcal{Y}_t^n)^{-1} a_n(\mathcal{X}_t^n), \quad (41)$$

where \mathcal{Y}^n is the solution of

$$d\mathcal{Y}_t^n = a'_n(\mathcal{X}_t^n) \mathcal{Y}_t^n dW_t + b'_n(\mathcal{X}_t^n) \mathcal{Y}_t^n dt, \quad \mathcal{Y}_0^n = 1. \quad (42)$$

In the sequel we will repeatedly use the positivity of \mathcal{Y}^n and the control

$$E\left(\sup_{t \in [0,1]} (\mathcal{Y}_t^n)^p \right) + E\left(\sup_{t \in [0,1]} (\mathcal{Y}_t^n)^{-p} \right) \leq c(p). \quad (43)$$

From this we can see that the random variables U^n and V^n are elements of $\mathbb{D}^{3, \infty}$ and using (41) with the linearity of the operator D , we have

$$\begin{aligned} D_t U^n &= \int_0^1 \mathbf{1}_{\{t \leq s\}} \mathcal{Y}_s^n (\mathcal{Y}_t^n)^{-1} a_n(\mathcal{X}_t^n) d\mu(s) = a_n(\mathcal{X}_t^n) (\mathcal{Y}_t^n)^{-1} \mathbf{1}_{\{t \leq 1\}} \int_{[t,1]} \mathcal{Y}_s^n d\mu(s), \\ D_t V^n &= a_n(\mathcal{X}_t^n) \mathcal{Y}_1^n (\mathcal{Y}_t^n)^{-1} \mathbf{1}_{\{t \leq 1\}}. \end{aligned}$$

Using Theorem 2.1.2 p. 86 in [21], a sufficient condition for the existence of a density for (U^n, V^n) is that its Malliavin covariance matrix γ_{U^n, V^n} satisfy a degeneracy condition given, for instance, by the following lemma.

Lemma 3. γ_{U^n, V^n} is a a.s invertible matrix and for all $p \geq 1$, we have

$$E(|\det(\gamma_{U^n, V^n})|^{-p}) \leq c(p).$$

Proof. To have shorter notations, during the proof we will denote by c_* any generic positive random variable which satisfies $E(c_*^{-p}) \leq c(p)$ with $c(p)$ as in the statement of the lemma. By direct computations we have,

$$\langle U^n, U^n \rangle_H = \int_0^1 a_n^2(\mathcal{X}_t^n)(\mathcal{Y}_t^n)^{-2} \left(\int_{[t,1]} \mathcal{Y}_s^n d\mu(s) \right)^2 dt, \quad (44)$$

$$\langle U^n, V^n \rangle_H = \int_0^1 a_n^2(\mathcal{X}_t^n)(\mathcal{Y}_t^n)^{-2} \left(\int_{[t,1]} \mathcal{Y}_s^n d\mu(s) \right) dt \mathcal{Y}_1^n, \quad (45)$$

$$\langle V^n, V^n \rangle_H = \int_0^1 a_n^2(\mathcal{X}_t^n)(\mathcal{Y}_t^n)^{-2} dt (\mathcal{Y}_1^n)^2. \quad (46)$$

Now, define the probability density on $[0, 1]$

$$m_t^n = a_n^2(\mathcal{X}_t^n)(\mathcal{Y}_t^n)^{-2} \left(\int_0^1 a_n^2(\mathcal{X}_s^n)(\mathcal{Y}_s^n)^{-2} ds \right)^{-1}, \quad (47)$$

and put $f^n(t) := \int_{[t,1]} \mathcal{Y}_s^n d\mu(s)$. Thus we can write:

$$\det(\gamma_{U^n, V^n}) = \langle V^n, V^n \rangle_H^2 (\mathcal{Y}_1^n)^{-2} \left[\int_0^1 m_t^n f^n(t)^2 dt - \left(\int_0^1 m_s^n f^n(s) ds \right)^2 \right].$$

Hence the above bracket can be interpreted as the variance of the function $f^n(t)$ under the probability measure $m_t^n dt$ and hence:

$$\det(\gamma_{U^n, V^n}) = \langle V^n, V^n \rangle_H^2 (\mathcal{Y}_1^n)^{-2} \int_0^1 m_t^n \left[f^n(t) - \left(\int_0^1 m_r^n f^n(r) dr \right) \right]^2 dt.$$

But clearly under Assumption **(R)**, $\langle V^n, V^n \rangle_H^2 \geq \underline{a}^2 \inf_{t \in [0,1]} (\mathcal{Y}_t^n)^{-2} \inf_{t \in [0,1]} (\mathcal{Y}_t^n)^2$ and hence by (43) this yields, $\langle V^n, V^n \rangle_H^2 \geq c_*$, using our convention about generic positive random variables c_* . Similarly, by (47), we have $m_t^n \geq c_*$ and thus,

$$\det(\gamma_{U^n, V^n}) \geq c_* \int_0^1 \left[f^n(t) - \left(\int_0^1 m_r^n f^n(r) dr \right) \right]^2 dt.$$

Then, writing the integral above as

$$\int_0^{1/2} \left[f^n(t) - \left(\int_0^1 m_r^n f^n(r) dr \right) \right]^2 + \left[f^n(t+1/2) - \left(\int_0^1 m_r^n f^n(r) dr \right) \right]^2 dt,$$

and using the simple inequality $x^2 + y^2 \geq (x - y)^2/2$, we get: $\det(\gamma_{U^n, V^n}) \geq c_* \int_0^{1/2} \left(\int_{[t, t+1/2)} \mathcal{Y}_s^n d\mu(s) \right)^2 dt$. Using again $\inf_{s \in [0, 1]} \mathcal{Y}_s^n \geq c_*$, we obtain: $\det(\gamma_{U^n, V^n}) \geq c_* \int_0^{1/2} \mu([t, t + 1/2))^2 dt$. But this integral is positive as soon as $\mu((0, 1)) > 0$ which is the case by assumption (3). Thus the lemma is proved. \square

5.1.2 Bounds for the density

For the proof of (9), we make a crucial use of the fact that the diffusion process \mathcal{X}^n is one dimensional by introducing the classical transformation:

$$s_n(x) := \int_0^x a_n^{-1}(y) dy, \quad \mathcal{W}_t^n := s_n(\mathcal{X}_t^n).$$

By the assumptions on a , the function s_n is one to one on \mathbb{R} and the derivatives of s_n and s_n^{-1} are bounded independently of n . By Itô's formula, \mathcal{W}^n solves the equation $d\mathcal{W}_t^n = dW_t + \tilde{b}_n(\mathcal{W}_t^n) dt$ where $\tilde{b}_n(w) := \frac{b_n}{a_n} \circ s_n^{-1}(w) - \frac{1}{2} a_n' \circ s_n^{-1}(w)$ and the initial value is $\mathcal{W}_0^n = s_n(\mathcal{X}_0^n) = 0$. We let \tilde{P} be the probability defined on (Ω, \mathcal{A}) by

$$\frac{d\tilde{P}}{dP} = \exp \left(- \int_0^1 \tilde{b}_n(\mathcal{W}_u^n) dW_u - \frac{1}{2} \int_0^1 \tilde{b}_n^2(\mathcal{W}_u^n) du \right).$$

The Girsanov theorem implies that the process \mathcal{W}^n is under \tilde{P} a standard Brownian motion. Note that the random variables (U^n, V^n) have the following expressions with respect to this \tilde{P} -Brownian motion:

$$U^n = \int_0^1 s_n^{-1}(\mathcal{W}_r^n) d\mu(r), \quad (48)$$

$$V^n = s_n^{-1}(\mathcal{W}_1^n). \quad (49)$$

Now let h_0 and h_1 be non negative real functions, then:

$$E_P [h_0(U^n) h_1(V^n)] = E_{\tilde{P}} [h_0(U^n) h_1(V^n) L^n], \quad (50)$$

where $L^n = \exp \left(\int_0^1 \tilde{b}_n(\mathcal{W}_r^n) d\mathcal{W}_r^n - \frac{1}{2} \int_0^1 \tilde{b}_n^2(\mathcal{W}_r^n) dr \right)$. But using Itô's formula, $L^n = \exp \left(\tilde{B}_n(\mathcal{W}_1^n) - \frac{1}{2} \int_0^1 (\tilde{b}_n^2 + \tilde{b}_n')(\mathcal{W}_r^n) dr \right)$ where \tilde{B}_n is the primitive function of \tilde{b}_n vanishing at zero. Since \tilde{b}_n and \tilde{b}_n' are clearly bounded by $cn^{-1/2}$ for some constant c only depending on a and b and $|\tilde{B}_n(x)| \leq cn^{-1/2} |x|$ we have: $c^{-1} \exp(-cn^{-1/2} |\mathcal{W}_1^n|) \leq L^n \leq c \exp(cn^{-1/2} |\mathcal{W}_1^n|)$. By (49) and the boundedness of s_n' we deduce $c^{-1} \exp(-cn^{-1/2} |V^n|) \leq L^n \leq$

$c \exp(cn^{-1/2}|V^n|)$. From this and (50), we obtain:

$$\begin{aligned} c^{-1} E_{\tilde{P}} \left[h_0(U^n) h_1(V^n) e^{-cn^{-\frac{1}{2}}|V^n|} \right] \\ \leq E_P [h_0(U^n) h_1(V^n)] \leq c E_{\tilde{P}} \left[h_0(U^n) h_1(V^n) e^{cn^{-\frac{1}{2}}|V^n|} \right]. \end{aligned}$$

Hence we have transformed the problem of finding bounds for the density of the law of (U^n, V^n) under P into an analogous problem under \tilde{P} . Consequently the bounds for $\mathfrak{p}_{x_0}^n$ stated in (9) will follow from the next lemma.

Lemma 4. *Let h_0, h_1 be two non negative functions. There exist some constants $c_1 > c_2 > 0$, depending only on the coefficients a and b such that:*

$$\begin{aligned} c_1^{-1} \int \int h_0(u) h_1(v) e^{-c_1(u^2+v^2)} du dv \leq \\ E_{\tilde{P}} [h_0(U^n) h_1(V^n)] \leq c_2^{-1} \int \int h_0(u) h_1(v) e^{-c_2(u^2+v^2)} du dv. \end{aligned}$$

Proof. We first show the lower bound. Using that the random variable V^n is measurable with respect to \mathcal{W}_1^n (by (49)), we can write:

$$\begin{aligned} E_{\tilde{P}} [h_0(U^n) h_1(V^n)] &= E_{\tilde{P}} [h_1(V^n) E_{\tilde{P}} [h_0(U^n) | \mathcal{W}_1^n]] \\ &= \int \mathfrak{g}(w) h_1(s_n^{-1}(w)) E_{\tilde{P}} [h_0(U^n) | \mathcal{W}_1^n = w] dw, \end{aligned} \quad (51)$$

where \mathfrak{g} is the density of the standard Gaussian law. Now let us admit temporarily the following relation on the conditional law of U^n :

$$E_{\tilde{P}} [h_0(U^n) | \mathcal{W}_1^n] \geq c^{-1} e^{-c(\mathcal{W}_1^n)^2} \int h_0(u) e^{-cu^2} du. \quad (52)$$

Then $E_{\tilde{P}} [h_0(U^n) h_1(V^n)]$ is greater than:

$$c^{-1} \int h_0(u) e^{-cu^2} du \times \int \mathfrak{g}(w) h_1(s_n^{-1}(w)) e^{-cw^2} dw.$$

The change of variable $v = s_n^{-1}(w)$ in the second integral above, the inequalities $|w| \leq c|v|$ and $s'_n(v) \geq c$ give the new lower bound

$$c^{-1} \int h_0(u) e^{-cu^2} du \times \int \mathfrak{g}(s_n(v)) h_1(v) e^{-cv^2} dv,$$

with a new constant c . Since \mathfrak{g} is the Gaussian kernel and thanks to the inequality $|s_n(v)| \leq c|v|$, we deduce the required lower bound for $E_{\tilde{P}} [h_0(U^n) h_1(V^n)]$.

We obtain the upper bound quite similarly. Let us temporarily admit that for all ε small enough there exists $c(\varepsilon)$ such that:

$$E_{\tilde{P}} [h_0(U^n) | \mathcal{W}_1^n] \leq c(\varepsilon)^{-1} e^{\varepsilon(\mathcal{W}_1^n)^2} \int h_0(u) e^{-c(\varepsilon)u^2} du. \quad (53)$$

Plugging this in equation (51), we deduce that $E_{\bar{P}}[h_0(U^n) | \mathcal{W}_1^n]$ is smaller than

$$c(\varepsilon)^{-1} \int h_0(u) e^{-c(\varepsilon)u^2} du \times \int \mathbf{g}(w) e^{\varepsilon w^2} h_1(s_n^{-1}(w)) dw.$$

Since $\mathbf{g}(w) = \exp(-w^2/2)/\sqrt{2\pi}$, any choice of ε smaller than $1/4$ implies that the second integral in the equation above is bounded by $c \int e^{-\frac{1}{4}w^2} h_1(s_n^{-1}(w)) dw$. As for the lower bound, we conclude by the change of variable $v = s_n^{-1}(w)$.

The final step is to show that (52)–(53) really holds. This is done in the following lemma. \square

Lemma 5. *For some constant $c > 0$ and $\bar{\varepsilon} > 0$, we have*

$$E_{\bar{P}}[h_0(U^n) | \mathcal{W}_1^n] \geq c^{-1} e^{-c(\mathcal{W}_1^n)^2} \int h_0(u) e^{-cu^2} du. \quad (54)$$

For all $\varepsilon \in]0, \bar{\varepsilon}[$, there exists $c(\varepsilon) > 0$ such that,

$$E_{\bar{P}}[h_0(U^n) | \mathcal{W}_1^n] \leq c(\varepsilon)^{-1} e^{\varepsilon(\mathcal{W}_1^n)^2} \int h_0(u) e^{-c(\varepsilon)u^2} du. \quad (55)$$

Proof. Let us recall that the process $\mathcal{W}_t^* := \mathcal{W}_t^n - t\mathcal{W}_1^n$ is a Brownian bridge on $[0, 1]$, independent of the variable \mathcal{W}_1^n . Thus, we can evaluate the conditional expectation $E_{\bar{P}}[h_0(U^n) | \mathcal{W}_1^n = w]$ as the expectation (recall (48)),

$$E \left[h_0 \left(\int_0^1 s_n^{-1}(\mathcal{W}_t^* + tw) d\mu(t) \right) \right], \quad (56)$$

for \mathcal{W}^* some Brownian bridge. This Brownian bridge itself admits a decomposition

$$\mathcal{W}_t^* = \xi \eta_t + \mathcal{W}_t^{**}, \quad (57)$$

where ξ is a $\mathcal{N}(0, 1)$ variable and η is the deterministic triangle shaped function:

$$\eta_t = \begin{cases} t & \text{if } t \in [0, 1/2] \\ (1-t) & \text{if } t \in [1/2, 1] \end{cases},$$

and \mathcal{W}^{**} is the process on $[0, 1]$ constructed as the concatenation of two independent Brownian bridges, one on $[0, 1/2]$ and another on $[1/2, 1]$. Furthermore in this decomposition the r.v. η and the process \mathcal{W}^{**} are independent.

For any realization of \mathcal{W}^{**} we can introduce the real function,

$$x \mapsto g_{\mathcal{W}^{**}}(x) = \int_0^1 \{s_n^{-1}(x\eta_t + \mathcal{W}_t^{**} + tw)\} d\mu(t).$$

Using (57) and the independence of ξ and \mathcal{W}^{**} , the quantity (56) now writes,

$$E_{(\mathcal{W}^{**})} E_{(\xi)} [h_0(g_{\mathcal{W}^{**}}(\xi))] \quad (58)$$

where the inner expectation denotes the expectation with respect to the random variable ξ and the outer one with respect to the process \mathcal{W}^{**} .

First we evaluate the inner expectation. Using that ξ is a standard Gaussian variable we have

$$E_{(\xi)} [h_0(g_{\mathcal{W}^{**}}(\xi))] = (2\pi)^{-1/2} \int h_0(g_{\mathcal{W}^{**}}(x)) e^{-\frac{x^2}{2}} dx. \quad (59)$$

Note now that for any realization of \mathcal{W}^{**} , the function $x \mapsto g_{\mathcal{W}^{**}}(x)$ is differentiable and using that $\frac{1}{c} \leq (s_n^{-1})' \leq c$ we get

$$\frac{1}{c} \int_0^1 \eta_t d\mu(t) \leq g'_{\mathcal{W}^{**}}(x) \leq c \int_0^1 \eta_t d\mu(t).$$

By assumption (3) on the measure μ the integral $\int_0^1 \eta_t d\mu(t)$ is positive. Thus the function $x \mapsto g_{\mathcal{W}^{**}}(x)$ is invertible on \mathbb{R} , with a derivative bounded from above and from below by some constant independent of \mathcal{W}^{**} and n . This allows us to make a change of variable in (59) to obtain the bounds

$$c^{-1} \int h_0(u) e^{-\frac{(g_{\mathcal{W}^{**}}^{-1}(u))^2}{2}} du \leq E_{(\xi)} [h_0(g_{\mathcal{W}^{**}}(\xi))] \leq c \int h_0(u) e^{-\frac{(g_{\mathcal{W}^{**}}^{-1}(u))^2}{2}} du. \quad (60)$$

Now the proofs of (54) and (55) are treated separately.

- For the lower bound, we have seen that $g_{\mathcal{W}^{**}}^{-1}$ is globally Lipschitz with a constant independent of \mathcal{W}^{**} and thus $|g_{\mathcal{W}^{**}}^{-1}(u)| \leq c|u| + |g_{\mathcal{W}^{**}}^{-1}(0)| \leq c|u| + c|g_{\mathcal{W}^{**}}(0)|$. In addition, a simple computation from the definition of $g_{\mathcal{W}^{**}}$ and the boundedness of $(s_n^{-1})'$ shows that

$$|g_{\mathcal{W}^{**}}(0)| \leq c[|w| + \sup_{t \in [0,1]} |\mathcal{W}_t^{**}|]. \quad (61)$$

Using this in (60) we find a new lower bound for the inner expectation:

$$E_{(\xi)} [h_0(g_{\mathcal{W}^{**}}(\xi))] \geq c^{-1} e^{-cw^2} e^{-c \sup_{t \in [0,1]} (\mathcal{W}_t^{**})^2} \int h_0(u) e^{-cu^2} du.$$

Taking the expectation with respect to \mathcal{W}^{**} proves that (58) is greater $c^{-1} E_{(\mathcal{W}^{**})} \left(e^{-c \sup_{t \in [0,1]} (\mathcal{W}_t^{**})^2} \right) \times e^{-cw^2} \int h_0(u) e^{-cu^2} du$. This gives (54).

- For the upper bound, we write using that $g_{\mathcal{W}^{**}}$ is Lipschitz $\|u\| - |g_{\mathcal{W}^{**}}(0)| \leq |g_{\mathcal{W}^{**}}(g_{\mathcal{W}^{**}}^{-1}(u)) - g_{\mathcal{W}^{**}}(0)| \leq c |g_{\mathcal{W}^{**}}^{-1}(u)|$. Together with the relation $(x - y)^2 \geq x^2 \frac{\varepsilon}{1+\varepsilon} - \varepsilon y^2$ (for $x, y \in \mathbb{R}, \varepsilon \in (0, 1)$), we deduce that $\exp(-\frac{1}{2}(g_{\mathcal{W}^{**}}^{-1}(u))^2)$ is upper bounded by

$$\exp\left(-\frac{\varepsilon u^2}{2c^2(1+\varepsilon)} + \frac{\varepsilon(g_{\mathcal{W}^{**}}(0))^2}{2c^2}\right) \leq \exp\left(-\frac{\varepsilon u^2}{2c^2(1+\varepsilon)} + \varepsilon w^2 + \varepsilon \sup_{t \in [0,1]} (\mathcal{W}_t^{**})^2\right)$$

where we have used (61). Combining this with (60) and taking the expectation with respect to \mathcal{W}^{**} , we get that the quantity (58) is smaller than: $c \int h_0(u) e^{-\frac{\varepsilon u^2}{2c(1+\varepsilon)}} e^{\varepsilon w^2} E_{\mathcal{W}^{**}} \left(e^{\varepsilon \sup_{t \in [0,1]} (\mathcal{W}_t^{**})^2} \right)$. The last expectation is finite as soon as ε is small enough, and thus (55) holds. \square

5.1.3 Proof of Proposition 1

To have shorter notations we set $\bar{\mathcal{X}}_j^n = \int_0^1 \mathcal{X}_{j+t}^n d\mu(t)$, for $j \geq 0$. First we prove:

Lemma 6. *Let us define Γ^k the Malliavin covariance matrix of the vector $\left(\bar{\mathcal{X}}_j^n, \mathcal{X}_{j+1}^n\right)_{j=0, \dots, k-1}$ of size $2k$.*

Then this matrix is a.s invertible and $E(\det(\Gamma^k)^{-p}) \leq c(p, k)$.

Proof. In the case $k = 1$, the lemma reduces to Lemma 3. For $k \geq 2$, we can proceed by induction on k using, indeed, that the Markov property of the process \mathcal{X}^n induces actually in specific relations between the columns of Γ^k .

To see this, notice that firstly by (41) if $t < k - 1$ and $s > k - 1$, we have $D_t \mathcal{X}_s^n = D_t \mathcal{X}_{k-1}^n \mathcal{Y}_s^n (\mathcal{Y}_{k-1}^n)^{-1}$; and secondly if $t > k - 1$ and $s < k - 1$, $D_t \mathcal{X}_s^n = 0$. Using these two properties, a calculation shows that if $(C_j)_{j=1, \dots, 2k}$ denote the columns of Γ^k , we have the relation,

$$[C_{2k-1}; C_{2k}] = \left[\left(\int_0^1 \mathcal{Y}_{k-1+s}^n (\mathcal{Y}_{k-1}^n)^{-1} d\mu(s) \right) C_{2k-2}; \mathcal{Y}_k^n (\mathcal{Y}_{k-1}^n)^{-1} C_{2k-2} \right] + \begin{pmatrix} 0 & 0 \\ \vdots & \vdots \\ 0 & 0 \\ [\gamma_k] \end{pmatrix}$$

where γ_k is the matrix of size 2×2 given by

$$\begin{pmatrix} \int_{k-1}^k (D_t \bar{\mathcal{X}}_{k-1}^n)^2 dt & \int_{k-1}^k (D_t \mathcal{X}_k^n) (D_t \bar{\mathcal{X}}_{k-1}^n) dt \\ \int_{k-1}^k (D_t \mathcal{X}_k^n) (D_t \bar{\mathcal{X}}_{k-1}^n) dt & \int_{k-1}^k (D_t \mathcal{X}_k^n)^2 dt \end{pmatrix}.$$

This proves that $\det \Gamma^k = \det \Gamma^{k-1} \det \gamma_k$. By the induction relation, it remains to prove $E((\det \gamma_k)^{-p}) \leq c(p)$. But direct computation using (41) shows that the component of γ_k have the following expression:

$$\begin{aligned} (\gamma_k)_{11} &= \int_{k-1}^k a_n(\mathcal{X}_t^n)^2 (\mathcal{Y}_t^n)^{-2} \left(\int_{[t,1]} \mathcal{Y}_{k-1+s}^n d\mu(s) \right)^2 dt, \\ (\gamma_k)_{12} &= (\gamma_k)_{21} = \int_{k-1}^k a_n(\mathcal{X}_t^n)^2 (\mathcal{Y}_t^n)^{-2} \left(\int_{[t,1]} \mathcal{Y}_{k-1+s}^n d\mu(s) \right) dt \mathcal{Y}_k^n, \\ (\gamma_k)_{22} &= \int_{k-1}^k a_n(\mathcal{X}_t^n)^2 (\mathcal{Y}_t^n)^{-2} dt (\mathcal{Y}_k^n)^2. \end{aligned}$$

These expressions are similar to the ones obtained for γ_{U^n, V^n} in lemma 3 (see (44)–(46)) and similarly, one gets $E((\det \gamma_k)^{-p}) \leq c(p)$. The lemma is proved. \square

Now we can deduce the Proposition 1. Recalling (10–13) we can find a invertible matrix M of size $2k \times 2k$ that maps $\left((\bar{\mathcal{X}}_j^n, \mathcal{X}_{j+1}^n) \right)_{j=0, \dots, k-1}$ into a vector whose $k+1$ first components are exactly (U_0^n, \dots, U_k^n) . Denoting $\hat{\Gamma}^k$ the Malliavin covariance matrix of the image by M of $\left((\bar{\mathcal{X}}_j^n, \mathcal{X}_{j+1}^n) \right)_{j=0, \dots, k-1}$, we have $\hat{\Gamma}^k = M\Gamma^k M^*$. Thus, Lemma 6 yields $E((\det(\hat{\Gamma}^k))^{-p}) \leq c(p, k)$ since M is invertible.

Observe that the covariance matrix K is the matrix extracted from the $k+1$ first rows and columns of $\hat{\Gamma}^k$. Hence, one has

$$[\det(K)]^{\frac{1}{k+1}} \geq \lambda_{\min}(K) \geq \lambda_{\min}(\hat{\Gamma}^k) \geq \frac{\det(\hat{\Gamma}^k)}{[\text{Tr}(\hat{\Gamma}^k)]^{2k-1}}$$

(where $\lambda_{\min}(A)$ is the smallest eigenvalue of the symmetric positive matrix A). This implies Proposition 1 since $E([\text{Tr}(\hat{\Gamma}^k)]^p) \leq c(p, k)$ for all $p, k \geq 1$.

5.2 Some estimates on the change of measures

For this section we denote by \mathcal{X} the canonical process on $\mathcal{C}([0, \infty))$ and we consider the random variable on this space defined by $H = f(U_0, \dots, U_k)$, where (U_0, \dots, U_k) is given by (10)–(13) with the canonical process \mathcal{X} in place of $\mathcal{X}^{\theta, n}$; we denote by E_{θ, x_0}^n the expectation with respect to the measure induced on the canonical space by the law of $\mathcal{X}^{\theta, n}$ solution of (6).

Lemma 7. *There exist $r \geq 1$ and a constant $c(k) \geq 0$, such that $\forall H = f(U_0, \dots, U_k) \geq 0$, $\forall \theta, \theta' \in \Theta$, $\forall x_0 \in \mathbb{R}$, we have*

$$E_{\theta', x_0}^n[H] \leq c(k) E_{\theta, x_0}^n[H^r]^{\frac{1}{r}}.$$

Proof. Recalling the notation of Section 2.1.1 we denote $\mathbf{p}_{x_0}^n(u, v, \theta)$ the density of the vector (8) and for $j = 0, \dots, k-1$ we let

$$Z_{j, \theta, \theta'} = \frac{\mathbf{p}_{\mathcal{X}_j}^n \left(\int_0^1 (\mathcal{X}_{j+s} - \mathcal{X}_j) d\mu(s), (\mathcal{X}_{j+1} - \mathcal{X}_j), \theta' \right)}{\mathbf{p}_{\mathcal{X}_j}^n \left(\int_0^1 (\mathcal{X}_{j+s} - \mathcal{X}_j) d\mu(s), (\mathcal{X}_{j+1} - \mathcal{X}_j), \theta \right)}. \quad (62)$$

Then using the Markov property of the process \mathcal{X} under the laws P_θ^n and $P_{\theta'}^n$, we have

$$E_{\theta', x_0}^n[H] = E_{\theta, x_0}^n \left[H \prod_{j=0}^{k-1} Z_{j, \theta, \theta'} \right] \leq E_{\theta, x_0}^n[H^r]^{\frac{1}{r}} E_{\theta, x_0}^n \left[\prod_{j=0}^{k-1} (Z_{j, \theta, \theta'})^{r'} \right]^{\frac{1}{r'}},$$

where r and r' are conjugate exponents. But we know by Theorem 3 that there exist two constants $0 < c_2 \leq c_1$ (uniform en θ, x_0, n) such that

$$c_1^{-1} e^{-c_1(u^2+v^2)} \leq \mathbf{p}_{x_0}^n(u, v, \theta) \leq c_2^{-1} e^{-c_2(u^2+v^2)}.$$

Then one can bound the conditional expectation $E_{\theta, x_0}^n [(Z_{k-1, \theta, \theta'})^{r'} \mid \mathcal{X}_s, s \leq (k-1)]$ by

$$\frac{c_1^{(r'-1)}}{c_2^{r'}} \int_{\mathbb{R}^2} e^{(u^2+v^2)(-r'c_2+(r'-1)c_1)} du dv.$$

But if r is chosen large enough such that r' is sufficiently close to 1 the latter integral converges and is equal to some constant κ . Proceeding by induction we get:

$$E_{\theta, x_0}^n \left[\prod_{j=0}^{k-1} (Z_{j, \theta, \theta'})^{r'} \right]^{\frac{1}{r'}} \leq E_{\theta, x_0}^n \left[\prod_{j=0}^{k-2} (Z_{j, \theta, \theta'})^{r'} \right]^{\frac{1}{r'}} \kappa^{\frac{1}{r'}} \leq \dots \leq \kappa^{k/r'}$$

which gives the result. \square

Lemma 8. *There exist $c(k) \geq 0$ and $\alpha \geq 1$ such that $\forall H = f(U_0, \dots, U_k)$ (with $E_{\theta, x_0}^n |H|^\alpha < +\infty$), $\forall \theta, \theta' \in \Theta$, $\forall x_0 \in \mathbb{R}$, we have*

$$|E_{\theta', x_0}^n [H] - E_{\theta, x_0}^n [H]| \leq c(k) |\theta - \theta'| [E_{\theta, x_0}^n |H|^\alpha]^{\frac{1}{\alpha}}. \quad (63)$$

Proof. Using the notations of Lemma 7, we write

$$\begin{aligned} E_{\theta', x_0}^n [H] - E_{\theta, x_0}^n [H] &= E_{\theta, x_0}^n \left[\left(\prod_{j=0}^{k-1} Z_{j, \theta, \theta'} - 1 \right) H \right] \\ &= \sum_{i=0}^{k-1} E_{\theta, x_0}^n \left[(Z_{i, \theta, \theta'} - 1) \prod_{j=i+1}^{k-1} Z_{j, \theta, \theta'} H \right]. \end{aligned}$$

Thus for conjugate exponents α and β , the left hand side of (63) is bounded by

$$\begin{aligned} &\sum_{i=0}^{k-1} [E_{\theta, x_0}^n |H|^\alpha]^{\frac{1}{\alpha}} E_{\theta, x_0}^n \left[|Z_{i, \theta, \theta'} - 1|^\beta \prod_{j=i+1}^{k-1} (Z_{j, \theta, \theta'})^\beta \right]^{\frac{1}{\beta}} \\ &= \sum_{i=0}^{k-1} [E_{\theta, x_0}^n |H|^\alpha]^{\frac{1}{\alpha}} E_{\theta, x_0}^n \left[|Z_{i, \theta, \theta'} - 1|^\beta E_{\theta, x_0}^n \left[\prod_{j=i+1}^{k-1} (Z_{j, \theta, \theta'})^\beta \mid \mathcal{X}_s, s \leq i+1 \right] \right]^{\frac{1}{\beta}} \end{aligned}$$

Using the Markov property of \mathcal{X} it can be shown exactly as in Lemma 7 that the conditional expectation in the equation above is finite, as soon as β is small enough and bounded by κ^{k-i-1} . Thus by Lemma 9 below, we deduce

$$|E_{\theta', x_0}^n [H] - E_{\theta, x_0}^n [H]| \leq c(\beta) |\theta - \theta'| [E_{\theta, x_0}^n |H|^\alpha]^{\frac{1}{\alpha}} \sum_{i=0}^{k-1} \kappa^{\frac{k-i-1}{\beta}}. \quad \square$$

Lemma 9. *There exists $\bar{\beta} > 1$ such that for all $1 < \beta \leq \bar{\beta}$ we have:*

$$E_{\theta, x_0}^n \left[|Z_{i, \theta, \theta'} - 1|^{\beta} \right]^{\frac{1}{\beta}} \leq c(\beta) |\theta - \theta'|.$$

Proof. Using expression of $Z_{i, \theta, \theta'}$, equation (8) with the Markov property it suffices to bound the quantity

$$E_{x_0, \theta}^n \left[\left| \frac{\mathbf{p}_{x_0}^n(U^n, V^n, \theta') - \mathbf{p}_{x_0}^n(U^n, V^n, \theta)}{\mathbf{p}_{x_0}^n(U^n, V^n, \theta)} \right|^{\beta} \right]^{\frac{1}{\beta}}. \quad (64)$$

By Theorem 4 with $k = 1$ the function $\theta \rightarrow \mathbf{p}_{x_0}(U^n, V^n, \theta)$ is absolutely continuous thus we can write the quantity above as: $E_{x_0, \theta}^n \left[\left| \frac{\int_{\theta}^{\theta'} \dot{\mathbf{p}}_{x_0}^n(U^n, V^n, s) ds}{\mathbf{p}_{x_0}^n(U^n, V^n, \theta)} \right|^{\beta} \right]^{\frac{1}{\beta}}$.

Using first the Minkowski inequality, a change of measure and then the Hölder inequality one find the following bounds for this quantity:

$$\begin{aligned} & \int_{\theta}^{\theta'} E_{x_0, s}^n \left[\left| \frac{\dot{\mathbf{p}}_{x_0}^n(U^n, V^n, s)}{\mathbf{p}_{x_0}^n(U^n, V^n, \theta)} \right|^{\beta} \frac{\mathbf{p}_{x_0}^n(U^n, V^n, \theta)}{\mathbf{p}_{x_0}^n(U^n, V^n, s)} \right]^{\frac{1}{\beta}} ds \\ & \leq \int_{\theta}^{\theta'} E_{x_0, s} \left[\left| \frac{\dot{\mathbf{p}}_{x_0}^n(U^n, V^n, s)}{\mathbf{p}_{x_0}^n(U^n, V^n, s)} \right|^{\beta \alpha'} \right]^{\frac{1}{\beta \alpha'}} E_{x_0, s} \left[\left| \frac{\mathbf{p}_{x_0}^n(U^n, V^n, s)}{\mathbf{p}_{x_0}^n(U^n, V^n, \theta)} \right|^{(\beta-1)\beta'} \right]^{\frac{1}{\beta \beta'}} ds \end{aligned}$$

for two conjugate exponents α' and β' . But the first expectation in the right hand side above is bounded by Corollary 2 (with $k = 1$) for all choice of α' , β and the second expectation can be bounded if $(\beta - 1)\beta'$ is close enough to zero by using (9) as in the proof of Lemma 7. This gives that (64) is smaller than $c|\theta - \theta'|$. \square

5.3 Technical lemmas

Lemma 10. *The covariance matrix of $(\tilde{U}_0^\theta, \dots, \tilde{U}_k^\theta)$ is $a^2(x_0, \theta)\tilde{K}$. This matrix is invertible.*

Proof. The first statement is easily checked from the expressions (18)–(20). Then we see that $(\lambda_0, \dots, \lambda_k)$ is element of the null space of $\tilde{K}(x_0, \theta)$ if the variance of $\sum_i \lambda_i \tilde{U}_i^\theta$ is null which gives, for all $j = 0, \dots, k - 1$:

$$\int_0^1 [\lambda_j \mu([s, 1]) + \lambda_{j+1} \mu([0, s])]^2 ds = 0.$$

But from the assumption (3) on μ , the condition, $\alpha \mu([0, s]) + \beta \mu([s, 1]) = 0$ ds-a.e on $[0, 1]$, implies that $\alpha = \beta = 0$ and hence we deduce $\lambda_j = 0, \forall j$. \square

Lemma 11. *Let (G_0, \dots, G_k) be a centered Gaussian vector with invertible covariance matrix C_{k+1} and let us denote by C_{k-1} the covariance matrix of (G_1, \dots, G_{k-1}) . Then,*

$$\sum_{0 \leq j, j' \leq k} G_j [C_{k+1}]_{j, j'}^{-1} G_{j'} - \sum_{1 \leq j, j' \leq k-1} G_j [C_{k-1}]_{j, j'}^{-1} G_{j'}, \quad (65)$$

is a $\chi^2(2)$ random variable.

Proof. Write the Gram-Schmidt orthonormalization procedure for the \mathbf{L}^2 vectors G_1, \dots, G_k, G_0 as:

$$\begin{bmatrix} \mathcal{H}_0 \\ \vdots \\ \mathcal{H}_k \end{bmatrix} = P_k \begin{bmatrix} G_0 \\ \vdots \\ G_k \end{bmatrix},$$

where the variables $\mathcal{H}_0, \dots, \mathcal{H}_k$ are i.i.d. with standard Gaussian law and P_k is some triangular matrix. Then a few linear algebra shows that (65) is equal to $\sum_{j=0}^k \mathcal{H}_j^2 - \sum_{j=1}^{k-1} \mathcal{H}_j^2 = \mathcal{H}_0^2 + \mathcal{H}_k^2$ and thus is chi-square distributed. \square

References

- [1] Barndorff-Nielsen O.E. and Shepard N. (2001), Non-Gaussian Ornstein–Uhlenbeck-based models and some of their uses in financial economics (with discussion) *Journal of the Royal Statistical Society, Series B* **63**, p.167–241.
- [2] Ditlevsen P. D., Ditlevsen S. and Andersen K. K. (2002), The fast climate fluctuations during the stadial and interstadial climate states *Ann. Glaciology* **35**, p.457–462.
- [3] Ditlevsen S. and Sørensen M. (2004) Inference for observations of integrated diffusion processes *Scandinavian Journal of Statistics* **31(3)**, p.417–429.
- [4] Dohnal G. (1987) On estimating the diffusion coefficient *J. Appl. Probab.* **24(1)**, p.105–114.
- [5] Genon–Catalot V. and Jacod J. (1993) On the estimation of the diffusion coefficient for multi–dimensional processes. *Ann. Inst. Henri Poincaré* **29**, p.119–151.
- [6] Gloter A. (2000) Discrete sampling of an integrated diffusion process and parameter estimation of the diffusion coefficient. *ESAIM: P&S*, **4**, p.205

- [7] Gloter A. (2001) Parameter estimation for a discrete sampling of an integrated Ornstein–Uhlenbeck process. *Statistics*, **23**, p.225–243.
- [8] Gloter A. and Jacod J. (2001) Diffusions with measurement errors. I. Local asymptotic normality. *ESAIM: P&S*, **5**, p.225–242.
- [9] Gobet E. (2001), Local asymptotic mixed normality property for elliptic diffusion: a Malliavin calculus approach. *Bernoulli* **7**, no. 6, p.899–912.
- [10] Gobet E. (2002), LAN property for ergodic diffusions with discrete observations. *Ann. Inst. H. Poincaré Probab. Statist.* **38**, no. 5, p.711–737.
- [11] Hirsch F. and Song S. (1997), Criteria of positivity for the density of a Wiener functional. *Bull. Sci. Math.*, **121**, p.261–273.
- [12] Hirsch F. and Song S. (1998), Properties of the set of positivity for the density of a regular Wiener functional. *Bull. Sci. Math.*, **122**, p.1–15.
- [13] Ibragimov I. A. And Has’minskii R. Z. (1981), *Statistical estimation. Asymptotic theory*. Translated from the Russian by Samuel Kotz. Applications of Mathematics, 16. Springer-Verlag, New York-Berlin.
- [14] Jacod J. (1996), On continuous conditional Gaussian martingales and stable convergence in law. *Séminaire de probabilité XXXI, Lecture notes in Mathematics* **1655**, p.232–246.
- [15] Jeganathan P. (1982), On the asymptotic theory of estimation when the limit of the log-likelihood ratios is mixed normal. *Sankhyā Ser. A* **44**, no. 2, p.173–212.
- [16] Jeganathan, P. (1983), Some asymptotic properties of risk functions when the limit of the experiment is mixed normal. *Sankhyā Ser. A* **45**, no. 1, p.66–87.
- [17] Kohatsu–Higa A. (2003), Lower bounds for densities of uniformly elliptic random variables on Wiener space. *Probab. Theory Rel. Fields*, **126**, p.421–457.
- [18] Krée P. and Soize C. (1986), *Mathematics of random phenomena. Random vibrations of mechanical structures.*, Mathematics and its Applications 32. D. Reidel Publishing Co., Reidel, Dordrecht.
- [19] Kunita H. (1984), Stochastic differentiable equations and stochastic flows of diffeomorphisms (Ecole d’Eté de Probabilités de Saint-Flour, 1982), 144–305, *Lecture Notes in Math. 1097*, Springer Verlag.
- [20] Le Cam L. and Lo Yang G. (2000), *Asymptotics in statistics*, Second edition, Springer, New York.

- [21] Nualart D. (1995), *The Malliavin Calculus and Related Topics*. Probability and its application, Springer–Verlag, New-York.
- [22] Nualart D. (1998), Analysis on Wiener space and anticipating stochastic calculus. *Lectures on probability theory and statistics (Saint-Flour, 1995)*, 123–227, *Lecture Notes in Math.*, 1690, Springer, Berlin.
- [23] Prakasa Rao B. L. S. (1999), *Statistical inference for diffusion type processes*, Kendall’s Library of Statistics 8, Edward Arnold, London.