

ECOLE POLYTECHNIQUE

CENTRE DE MATHÉMATIQUES APPLIQUÉES

UMR CNRS 7641

91128 PALAISEAU CEDEX (FRANCE). Tél: 01 69 33 41 50. Fax: 01 69 33 30 11

<http://www.cmap.polytechnique.fr/>

**Uniqueness of the Cheeger set of
a convex body**

V. Caselles, A. Chambolle, M. Novaga

R.I. 621

August 2007

Uniqueness of the Cheeger set of a convex body

V. Caselles^{*}, A. Chambolle[†], M. Novaga[‡]

Abstract

We prove that if $C \subset \mathbb{R}^N$ is of class C^2 and uniformly convex, then the Cheeger set of C is unique. The Cheeger set of C is the set which minimizes, inside C , the ratio perimeter over volume.

1 Introduction

Given an nonempty open bounded subset Ω of \mathbb{R}^N , we call Cheeger constant of Ω the quantity

$$h_\Omega = \min_{K \subseteq \Omega} \frac{P(K)}{|K|}. \quad (1)$$

Here $|K|$ denotes the N -dimensional volume of K and $P(K)$ denotes the perimeter of K . The minimum in (1) is taken over all nonempty sets of finite perimeter contained in Ω . A Cheeger set of Ω is any set $G \subseteq \Omega$ which minimizes (1). If Ω minimizes (1), we say that it is Cheeger in itself. We observe that the minimum in (1) is attained at a subset G of Ω such that ∂G intersects $\partial\Omega$: otherwise we would diminish the quotient $P(G)/|G|$ by dilating G .

For any set of finite perimeter K in \mathbb{R}^N , let us denote

$$\lambda_K := \frac{P(K)}{|K|}.$$

Notice that for any Cheeger set G of Ω , $\lambda_G = h_\Omega$. Observe also that G is a Cheeger set of Ω if and only if G minimizes

$$\min_{K \subseteq \Omega} P(K) - \lambda_G |K|. \quad (2)$$

We say that a set $\Omega \subset \mathbb{R}^N$ is calibrable if Ω minimizes the problem

$$\min_{K \subseteq \Omega} P(K) - \lambda_\Omega |K|. \quad (3)$$

In particular, if G is a Cheeger set of Ω , then G is calibrable. Thus, Ω is a Cheeger set of itself if and only if it is calibrable.

^{*}Departament de Tecnologia, Universitat Pompeu-Fabra, Barcelona, Spain,
e-mail: vicent.caselles@tecn.upf.es

[†]CMAP, Ecole Polytechnique, CNRS, 91128 Palaiseau, France,
e-mail: antonin.chambolle@polytechnique.fr

[‡]Dipartimento di Matematica, Università di Pisa, Largo B. Pontecorvo 5, 56127 Pisa, Italy,
e-mail: novaga@dm.unipi.it

Finding the Cheeger sets of a given Ω is a difficult task. This task is simplified if Ω is a convex set and $N = 2$. In that case, the Cheeger set in Ω is unique and is identified with the set $\Omega^R \oplus B(0, R)$ where $\Omega^R := \{x \in \Omega : \text{dist}(x, \partial\Omega) > R\}$ is such that $|\Omega^R| = \pi R^2$ and $A \oplus B := \{a + b : a \in A, b \in B\}$, $A, B \subset \mathbb{R}^2$ [2, 19]. We see in particular that it is convex. Moreover, a convex set $\Omega \subseteq \mathbb{R}^2$ is Cheeger in itself if and only if $\max_{x \in \partial\Omega} \kappa_\Omega(x) \leq \lambda_\Omega$ where $\kappa_\Omega(x)$ denotes the curvature of $\partial\Omega$ at the point x . This has been proved in [14, 9, 19, 2, 20], though it was stated in terms of calibrability in [9, 2]. The proof in [14] had also a complement result: if Ω is Cheeger in itself then Ω is strictly calibrable, that is, for any set $K \subset \Omega$, $K \neq \Omega$, then

$$0 = P(\Omega) - \lambda_\Omega |\Omega| < P(K) - \lambda_\Omega |K|,$$

and this implies that the capillary problem in absence of gravity (with vertical contact angle at the boundary)

$$\begin{aligned} -\operatorname{div} \left(\frac{Du}{\sqrt{1 + |Du|^2}} \right) &= \lambda_\Omega \quad \text{in } \Omega \\ -\frac{Du}{\sqrt{1 + |Du|^2}} \cdot \nu^\Omega &= 1 \quad \text{in } \partial\Omega \end{aligned} \tag{4}$$

has a solution. Indeed, both problems are equivalent [14, 18].

Our purpose in this paper is to extend the above result to \mathbb{R}^N , that is, to prove the uniqueness and convexity of the Cheeger set contained in a convex set $\Omega \subset \mathbb{R}^N$. We have to assume, in addition, that Ω is uniformly convex and of class C^2 . This regularity assumption is probably too strong, and its removal is the subject of current research [1]. The characterization of a convex set $\Omega \subset \mathbb{R}^N$ of class $C^{1,1}$ which is Cheeger in itself (also called calibrable) in terms of the mean curvature of its boundary was proved in [3]. The precise result states that such a set Ω is Cheeger in itself if and only if $\kappa_\Omega(x) \leq \lambda_\Omega$ for any $x \in \partial\Omega$, where $\kappa_\Omega(x)$ denotes the sum of the principal curvatures of the boundary of Ω , i.e. $(N - 1)$ times the mean curvature of $\partial\Omega$ at x . Moreover, in [3], the authors also proved that for any convex set $\Omega \subset \mathbb{R}^N$ there exists a maximal Cheeger set contained in Ω which is convex. These results were extended to convex sets Ω satisfying a regularity condition and anisotropic norms in \mathbb{R}^N (including the crystalline case) in [12].

In particular, we obtain that $\Omega \subset \mathbb{R}^N$ is the unique Cheeger set of itself, whenever Ω is a C^2 , uniformly convex calibrable set. We point out that, by Theorems 1.1 and 4.2 in [14], this uniqueness result is equivalent to the existence of a solution $u \in W_{\text{loc}}^{1,\infty}(\Omega)$ of the capillary problem (4).

Let us explain the plan of the paper. In Section 2 we collect some definitions and recall some results about the mean curvature operator in (4) and the subdifferential of the total variation. In Section 3 we state and prove the uniqueness result.

2 Preliminaries

2.1 BV functions

Let Ω be an open subset of \mathbb{R}^N . A function $u \in L^1(\Omega)$ whose gradient Du in the sense of distributions is a (vector valued) Radon measure with finite total variation in Ω is called a function of bounded variation. The class of such functions will be denoted by $BV(\Omega)$. The total variation of Du on Ω turns out to be

$$\sup \left\{ \int_{\Omega} u \operatorname{div} z \, dx : z \in C_0^\infty(\Omega; \mathbb{R}^N), \|z\|_{L^\infty(\Omega)} := \operatorname{ess\,sup}_{x \in \Omega} |z(x)| \leq 1 \right\}, \quad (5)$$

(where for a vector $v = (v_1, \dots, v_N) \in \mathbb{R}^N$ we set $|v|^2 := \sum_{i=1}^N v_i^2$) and will be denoted by $|Du|(\Omega)$ or by $\int_{\Omega} |Du|$. The map $u \rightarrow |Du|(\Omega)$ is $L^1_{\text{loc}}(\Omega)$ -lower semicontinuous. $BV(\Omega)$ is a Banach space when endowed with the norm $\int_{\Omega} |u| \, dx + |Du|(\Omega)$. We recall that $BV(\mathbb{R}^N) \subseteq L^{N/(N-1)}(\mathbb{R}^N)$.

A measurable set $E \subseteq \mathbb{R}^N$ is said to be of finite perimeter in \mathbb{R}^N if (5) is finite when u is substituted with the characteristic function χ_E of E and $\Omega = \mathbb{R}^N$. The perimeter of E is defined as $P(E) := |D\chi_E|(\mathbb{R}^N)$. For a complete monograph on functions of bounded variation we refer to [5].

Finally, let us denote by \mathcal{H}^{N-1} the $(N-1)$ -dimensional Hausdorff measure. We recall that when E is a finite-perimeter set with regular boundary (for instance, Lipschitz), its perimeter $P(E)$ also coincides with the more standard definition $\mathcal{H}^{N-1}(\partial E)$.

2.2 A generalized Green's formula

Let Ω be an open subset of \mathbb{R}^N . Following [7], let

$$X_2(\Omega) := \{z \in L^\infty(\Omega; \mathbb{R}^N) : \operatorname{div} z \in L^2(\Omega)\}.$$

If $z \in X_2(\Omega)$ and $w \in L^2(\Omega) \cap BV(\Omega)$ we define the functional $(z \cdot Dw) : C_0^\infty(\Omega) \rightarrow \mathbb{R}$ by the formula

$$\langle (z \cdot Dw), \varphi \rangle := - \int_{\Omega} w \varphi \operatorname{div} z \, dx - \int_{\Omega} w z \cdot \nabla \varphi \, dx.$$

Then $(z \cdot Dw)$ is a Radon measure in Ω ,

$$\int_{\Omega} (z \cdot Dw) = \int_{\Omega} z \cdot \nabla w \, dx \quad \forall w \in L^2(\Omega) \cap W^{1,1}(\Omega).$$

Recall that the outer unit normal to a point $x \in \partial\Omega$ is denoted by $\nu^\Omega(x)$. We recall the following result proved in [7].

Theorem 1. *Let $\Omega \subset \mathbb{R}^N$ be a bounded open set with Lipschitz boundary. Let $z \in X_2(\Omega)$. Then there exists a function $[z \cdot \nu^\Omega] \in L^\infty(\partial\Omega)$ satisfying $\|[z \cdot \nu^\Omega]\|_{L^\infty(\partial\Omega)} \leq \|z\|_{L^\infty(\Omega; \mathbb{R}^N)}$, and such that for any $u \in BV(\Omega) \cap L^2(\Omega)$ we have*

$$\int_{\Omega} u \operatorname{div} z \, dx + \int_{\Omega} (z \cdot Dw) = \int_{\partial\Omega} [z \cdot \nu^\Omega] u \, d\mathcal{H}^{N-1}.$$

Moreover, if $\varphi \in C^1(\overline{\Omega})$ then $[(\varphi z) \cdot \nu^\Omega] = \varphi [z \cdot \nu^\Omega]$.

This result is complemented with the following result proved by Anzellotti in [8].

Theorem 2. *Let $\Omega \subset \mathbb{R}^N$ be a bounded open set with a boundary of class C^1 . Let $z \in C(\bar{\Omega}; \mathbb{R}^N)$ with $\operatorname{div} z \in L^2(\Omega)$. Then*

$$[z \cdot \nu^\Omega](x) = z(x) \cdot \nu^\Omega(x) \quad \mathcal{H}^{N-1} \text{ a.e. on } \partial\Omega.$$

2.3 Some auxiliary results

Let Ω be an open bounded subset of \mathbb{R}^N with Lipschitz boundary, and let $\varphi \in L^1(\Omega)$. For all $\epsilon > 0$, we let $\Psi_\varphi^\epsilon : L^2(\Omega) \rightarrow (-\infty, +\infty]$ be the functional defined by

$$\Psi_\varphi^\epsilon(u) := \begin{cases} \int_\Omega \sqrt{\epsilon^2 + |Du|^2} + \int_{\partial\Omega} |u - \varphi| & \text{if } u \in L^2(\Omega) \cap BV(\Omega) \\ +\infty & \text{if } u \in L^2(\Omega) \setminus BV(\Omega). \end{cases} \quad (6)$$

As it is proved in [15], if $f \in W^{1,\infty}(\Omega)$, then the minimum $u \in BV(\Omega)$ of the functional

$$\Psi_\varphi^\epsilon(u) + \int_\Omega |u(x) - f(x)|^2 dx \quad (7)$$

belongs to $u \in C^{2+\alpha}(\Omega)$, for every $\alpha < 1$. The minimum u of (7) is a solution of

$$\begin{cases} u - \frac{1}{\lambda} \operatorname{div} \frac{Du}{\sqrt{\epsilon^2 + |Du|^2}} = f(x) & \text{in } \Omega \\ u = \varphi & \text{on } \partial\Omega \end{cases} \quad (8)$$

where the boundary condition is taken in a generalized sense [21], i.e.,

$$\left[\frac{Du}{\sqrt{\epsilon^2 + |Du|^2}} \cdot \nu^\Omega \right] \in \operatorname{sign}(\varphi - u) \quad \mathcal{H}^{N-1} \text{ a.e. on } \partial\Omega.$$

Observe that (8) can be written as

$$u + \frac{1}{\lambda} \partial \Psi_\varphi^\epsilon(u) \ni f. \quad (9)$$

We are particularly interested in the case where $\varphi = 0$. As we shall show below (see also [3]) in the case of interest to us we have $u > 0$ on $\partial\Omega$ and, thus, $\left[\frac{Du}{\sqrt{\epsilon^2 + |Du|^2}} \cdot \nu^\Omega \right] = -1$ \mathcal{H}^{N-1} a.e. on $\partial\Omega$. It follows that u is a solution of the first equation in (8) with vertical contact angle at the boundary.

As $\epsilon \rightarrow 0^+$, the solution of (8) converges to the solution of

$$\begin{cases} u + \frac{1}{\lambda} \partial \Psi_\varphi(u) = f(x) & \text{in } \Omega \\ u = \varphi & \text{on } \partial\Omega. \end{cases} \quad (10)$$

where $\Psi : L^2(\Omega) \rightarrow (-\infty, +\infty]$ is given by

$$\Psi_\varphi(u) := \begin{cases} \int_{\mathbb{R}^N} |Du| + \int_{\partial\Omega} |u - \varphi| & \text{if } u \in L^2(\Omega) \cap BV(\Omega) \\ +\infty & \text{if } u \in L^2(\Omega) \setminus BV(\Omega). \end{cases} \quad (11)$$

In this case $\partial\Psi_\varphi$ represents the operator $-\operatorname{div}(Du/|Du|)$ with the boundary condition $u = \varphi$ in $\partial\Omega$, and this connection is precisely given by the following Lemma (see [6]).

Lemma 2.1. *The following assertions are equivalent:*

- (a) $v \in \partial\Psi_\varphi(u)$;
- (b) $u \in L^2(\Omega) \cap BV(\Omega)$, $v \in L^2(\Omega)$, and there exists $z \in X_2(\Omega)$ with $\|z\|_\infty \leq 1$, such that

$$v = -\operatorname{div} z \quad \text{in } \mathcal{D}'(\Omega),$$

$$(z \cdot Du) = |Du|,$$

and

$$[z \cdot \nu^\Omega] \in \operatorname{sign}(\varphi - u) \quad \mathcal{H}^{N-1} \text{ a.e. on } \partial\Omega.$$

Notice that the solution $u \in L^2(\Omega)$ of (10) minimizes the problem

$$\min_{u \in BV(\Omega)} \int_{\Omega} |Du| + \int_{\partial\Omega} |u(x) - \varphi(x)| d\mathcal{H}^{N-1}(x) + \frac{\lambda}{2} \int_{\Omega} |u(x) - f(x)|^2 dx, \quad (12)$$

and the two problems are equivalent.

3 The uniqueness theorem

We now state our main result.

Theorem 3. *Let C be a convex body in \mathbb{R}^N . Assume that C is uniformly convex, with boundary of class C^2 . Then the Cheeger set of C is convex and unique.*

We do not believe that the regularity and the uniform convexity of C is essential for this result (see [1]).

Let us recall the following result proved in [3] (Theorems 6 and 8 and Proposition 4):

Theorem 4. *Let C be a convex body in \mathbb{R}^N with boundary of class $C^{1,1}$. For any $\lambda, \varepsilon > 0$, there is a unique solution u_ε of the equation:*

$$\begin{cases} u_\varepsilon - \frac{1}{\lambda} \operatorname{div} \frac{Du_\varepsilon}{\sqrt{\varepsilon^2 + |Du_\varepsilon|^2}} = 1 & \text{in } C \\ u_\varepsilon = 0 & \text{on } \partial C, \end{cases} \quad (13)$$

such that $0 \leq u_\varepsilon \leq 1$. Moreover, there exist λ_0 and ε_0 , depending only on ∂C , such that if $\lambda \geq \lambda_0$ and $\varepsilon \leq \varepsilon_0$, then u_ε is a concave function such that $u_\varepsilon \geq \alpha > 0$ on ∂C for some $\alpha > 0$. Hence, u_ε satisfies

$$\left[\frac{Du^\varepsilon}{\sqrt{\varepsilon^2 + |Du^\varepsilon|^2}} \cdot \nu^C \right] = \text{sign}(0 - u^\varepsilon) = -1 \quad \text{on } \partial C. \quad (14)$$

As $\varepsilon \rightarrow 0$, the functions u_ε converge to the concave function u which minimizes the problem

$$\min_{u \in BV(C)} \int_C |Du| + \int_{\partial C} |u(x)| d\mathcal{H}^{N-1}(x) + \frac{\lambda}{2} \int_C |u(x) - 1|^2 dx \quad (15)$$

or, equivalently, if u is extended with zero out of C , u minimizes

$$\int_{\mathbb{R}^N} |Du| + \frac{\lambda}{2} \int_{\mathbb{R}^N} |u - \chi_C|^2 dx.$$

The function u satisfies $0 \leq u < 1$. Moreover, the superlevel set $\{u \geq t\}$, $t \in (0, 1]$, is contained in C and minimizes the problem

$$\min_{F \subset C} P(F) - \lambda(1-t)|F|. \quad (16)$$

It was proved in [3] (see also [12]) that the set $C^* = \{u = \max_C u\}$ is the maximal Cheeger set contained in C , that is, the maximal set that solves (1). Moreover, one has $u = 1 - h_C/\lambda > 0$ in C^* and $h_C = \lambda C^*$.

If we want to consider what happens inside C^* and, in particular, if there are other Cheeger sets, we have to analyze the level sets of u_ε before passing to the limit as $\varepsilon \rightarrow 0^+$. In order to do this, let us introduce the following rescaling of u_ε :

$$v_\varepsilon = \frac{u_\varepsilon - m_\varepsilon}{\varepsilon} \leq 0,$$

where $m_\varepsilon = \max_C u_\varepsilon \rightarrow 1 - h_C/\lambda$ as $\varepsilon \rightarrow 0$. The function v_ε is a generalized solution of the equation:

$$\begin{cases} \varepsilon v_\varepsilon - \frac{1}{\lambda} \text{div} \frac{Dv_\varepsilon}{\sqrt{1 + |Dv_\varepsilon|^2}} = 1 - m_\varepsilon & \text{in } C \\ v_\varepsilon = -m_\varepsilon/\varepsilon & \text{on } \partial C. \end{cases} \quad (17)$$

We let $z_\varepsilon = Du_\varepsilon/\sqrt{\varepsilon^2 + |Du_\varepsilon|^2} = Dv_\varepsilon/\sqrt{1 + |Dv_\varepsilon|^2}$. Notice that z_ε is a vector field in $L^\infty(C)$, with uniformly bounded divergence, such that $|z_\varepsilon| \leq 1$ a.e. in C and, by (14), $[z_\varepsilon \cdot \nu_C] = -1$ on ∂C .

Let us study the limit of v_ε and z_ε as $\varepsilon \rightarrow 0$. Let us observe that, by concavity of v_ε , for each $\varepsilon > 0$ small enough and each $s \in (0, |C|)$, there exists a (convex) superlevel set C_s^ε of v_ε such that $|C_s^\varepsilon| = s$. We also observe that $\{v_\varepsilon = 0\}$ is a null set. Otherwise, since v_ε is concave, it would be a convex set of positive measure, hence with nonempty interior. We would then have that $v_\varepsilon = \text{div} z_\varepsilon = 0$, hence $1 - m_\varepsilon = 0$ in the interior of $\{v_\varepsilon = 0\}$. This is a contradiction with Theorem 4 for $\varepsilon > 0$ small enough. Hence we may take $C_0^\varepsilon := \{v_\varepsilon = 0\}$

and $C_{|C|}^\varepsilon := C$. The boundaries $\partial C_s^\varepsilon \cap C$ define in C a foliation, in the sense that for all $x \in C$, there exists a unique value of $s \in [0, |C|]$ such that $x \in \partial C_s^\varepsilon$.

We observe that a sequence of uniformly bounded convex sets is compact both for the L^1 and Hausdorff topologies. Hence, up to a subsequence, we may assume that C_s^ε converge to convex sets C_s , each of volume s , first for any $s \in \mathbb{Q} \cap (0, |C|)$ and then by continuity for any s . Possibly extracting a further subsequence, we may assume that there exists $s_* \in [0, |C|]$ such that v_ε goes to a concave function v in C_s for any $s < s_*$, and to $-\infty$ outside $C_* := C_{s_*}$. We may also assume that $z_\varepsilon \rightharpoonup z$ weakly* in $L^\infty(C)$, for some vector field z , satisfying $|z| \leq 1$ a.e. in C . From (13) we have in the limit

$$-\operatorname{div} z = \lambda(1 - u) \quad \text{in } \mathcal{D}'(C). \quad (18)$$

Moreover, by the results recalled in Section 2, it holds $-\operatorname{div} z \in \partial \Psi_0(u)$. We see from (18) that

$$-\operatorname{div} z = h_C \quad \text{in } C^*, \quad (19)$$

while $-\operatorname{div} z > h_C$ a.e. on $C \setminus C^*$. We let $s^* := |C^*|$, so that $C^* = C_{s^*}$. By Theorem 4, for $s \geq s^*$, the set C_s is a minimizer of the variational problem

$$\min_{E \subseteq C} P(E) - \mu_s |E|, \quad (20)$$

for some $\mu_s \geq h_C$ (μ_s is equal to the constant value of $-\operatorname{div} z = \lambda(1 - u)$ on $\partial C_s \cap C$, see eq. (16)). Notice that μ_s is bounded from above by $P(C)/(|C| - s)$: indeed, for $\varepsilon > 0$, one has

$$-\int_{C \setminus C_s^\varepsilon} \operatorname{div} z_\varepsilon(x) dx = \mathcal{H}^{N-1}(\partial C \setminus \partial C_s^\varepsilon) - \int_{\partial C_s^\varepsilon \cap C} \frac{|Du_\varepsilon|}{\sqrt{1 + |Du_\varepsilon|^2}} \leq P(C)$$

(since the inner normal to C_s^ε at $x \in \partial C_s^\varepsilon \cap C$ is $Du_\varepsilon(x)/|Du_\varepsilon(x)|$). On the other hand,

$$-\int_{C \setminus C_s^\varepsilon} \operatorname{div} z_\varepsilon(x) dx = \int_{C \setminus C_s^\varepsilon} \lambda(1 - u_\varepsilon(x)) dx \geq \mu_s^\varepsilon (|C| - s),$$

where μ_s^ε is the constant value of $\lambda(1 - u_\varepsilon)$ on the level set $\partial C_s^\varepsilon \cap C$, and goes to μ_s as $\varepsilon \rightarrow 0$. A more careful analysis would show, in fact, that $\mu_s \leq (P(C) - P(C_s))/(|C| - s)$.

For $s > s^*$, we have $\mu_s > h_C$ and the set C_s is the unique minimizer of the variational problem (20). As a consequence (see [3, 12]) for any $s > s^*$ the set C_s is also the unique minimizer of $P(E)$ among all $E \subseteq C$ of volume s .

Lemma 3.1. *We have $s_* > 0$ and the sets C_s are Cheeger sets in C for any $s \in [s_*, s^*]$.*

Proof. Let $s_* < s \leq |C|$. If $x \in \partial C_s^\varepsilon \setminus \partial C$, then

$$0 - v_\varepsilon(x) \leq Dv_\varepsilon(x) \cdot (\bar{x}_\varepsilon - x)$$

where $v_\varepsilon(\bar{x}_\varepsilon) = \max_C v_\varepsilon$. Hence, $\lim_{\varepsilon \rightarrow 0} \inf_{\partial C_s^\varepsilon \setminus \partial C} |Dv_\varepsilon| = +\infty$. Since $[z_\varepsilon \cdot \nu^C] = -1$ on ∂C and $P(C_s^\varepsilon) \rightarrow P(C_s)$, we deduce

$$\begin{aligned} & - \int_{\partial C_s^\varepsilon} [z_\varepsilon(x) \cdot \nu^{C_s^\varepsilon}(x)] d\mathcal{H}^{N-1}(x) \\ &= \int_{\partial C_s^\varepsilon \setminus \partial C} \frac{|Dv_\varepsilon(x)|}{\sqrt{1 + |Dv_\varepsilon(x)|^2}} d\mathcal{H}^{N-1}(x) + \mathcal{H}^{N-1}(\partial C_s^\varepsilon \cap \partial C) \rightarrow P(C_s) \end{aligned}$$

as $\varepsilon \rightarrow 0^+$. Hence,

$$\begin{aligned} \int_{\partial C_s} [z \cdot \nu^{C_s}] d\mathcal{H}^{N-1} &= \int_{C_s} \operatorname{div} z = \lim_{\varepsilon \rightarrow 0} \int_{C_s^\varepsilon} \operatorname{div} z_\varepsilon \\ &= \lim_{\varepsilon \rightarrow 0} \int_{\partial C_s^\varepsilon} [z_\varepsilon \cdot \nu_{C_s^\varepsilon}] d\mathcal{H}^{N-1} = -P(C_s). \end{aligned}$$

Since $|z| \leq 1$ a.e. in C , we deduce that $[z \cdot \nu^{C_s}] = -1$ on ∂C_s for any $s > s_*$ (in particular, we have $|z| = 1$ a.e. in $C \setminus C_*$). Using this and (19), for all $s_* < s \leq s^*$ we have

$$\frac{P(C_s)}{|C_s|} = h_C. \quad (21)$$

This has two consequences. First, from the isoperimetric inequality, we obtain

$$h_C = \frac{P(C_s)}{|C_s|} \geq \frac{P(B_1)}{|B_1|^{\frac{N-1}{N}} s^{\frac{1}{N}}},$$

if $s \in (s_*, s^*]$, so that $s_* > 0$. Moreover, C_s is a Cheeger set for any $s \in (s_*, s^*]$, and by continuity C_* is also a Cheeger set. \square

We point out that, since the sets C_s are convex minimizers of $P(E) - \mu_s|E|$ among all $E \subseteq C$, for $s \geq s_*$, their boundary is of class $C^{1,1}$ [10, 22], with curvature less than or equal to μ_s , and equal to μ_s in the interior of C (note that $\mu_s = h_C$ for $s \in [s_*, s^*]$).

Remark 3.2. Observe that we have either $s_* = s^*$ and therefore $C_* = C^*$, or $s_* < s^*$, and we have $C^* = \bigcup_{s \in (s_*, s^*)} C_s$. In the latter case, the supremum of κ_{C^*} on ∂C^* is equal to h_C . Indeed, if it were not the case, by considering $C' \subset \operatorname{int}(C^*)$, with curvature strictly below h_C , and the smallest set C_s , with $s > s_*$, which contains C' , we would have $\kappa_{C'}(x) \geq \kappa_{C_s}(x) = h_C$ at all $x \in \partial C' \cap \partial C_s$, a contradiction. In particular, if the supremum of κ_C on ∂C is strictly less than $P(C)/|C|$ (which implies $C = C^*$ by [3]) then $C = C_*$.

From the strong convergence of Dv_ε to Dv (in $L^2(C_s)$ for any $s < s_*$), we deduce that $z = \frac{Dv}{\sqrt{1+|Dv|^2}}$ in C_* . It follows that v satisfies the equation

$$-\operatorname{div} \frac{Dv}{\sqrt{1+|Dv|^2}} = h_C \quad \text{in } C_*. \quad (22)$$

Integrating both terms of (22) in C_* , we deduce that

$$\left[\frac{Dv}{\sqrt{1+|Dv|^2}} \cdot \nu^{C_*} \right] = -1 \quad \text{on } \partial C_*.$$

Lemma 3.3. *The set C_* is the minimal Cheeger set of C , i.e., any other Cheeger set of C must contain C_* .*

Proof. Let $K \subseteq C^*$ be a Cheeger set in C . We have

$$h_C|K| = - \int_K \operatorname{div} z = - \int_{\partial K} [z \cdot \nu^K] d\mathcal{H}^{N-1} = P(K)$$

so that $[z \cdot \nu^K] = -1$ a.e. on ∂K . Let ν^ϵ and ν be the vector fields of unit normals to the sets C_s^ϵ and C_s , $s \in [0, |C|]$, respectively. Observe that, by the Hausdorff convergence of C_s^ϵ to C_s as $\epsilon \rightarrow 0^+$ for any $s \in [0, |C|]$, we have that $\nu^\epsilon \rightarrow \nu$ a.e. in C . On the other hand, $|z_\epsilon + \nu^\epsilon| \rightarrow 0$ locally uniformly in $C \setminus \overline{C_*}$: indeed, we have in C

$$|z_\epsilon + \nu^\epsilon| = \left| \frac{Dv_\epsilon}{\sqrt{1 + |Dv_\epsilon|^2}} - \frac{Dv_\epsilon}{|Dv_\epsilon|} \right| = \left| \frac{|Dv_\epsilon|}{\sqrt{1 + |Dv_\epsilon|^2}} - 1 \right|.$$

Since (see the first lines of the proof of Lemma 3.1) $|Dv_\epsilon| \rightarrow \infty$ uniformly in any subset of C at positive distance from C_* , it shows the uniform convergence of $|z_\epsilon + \nu^\epsilon|$ to 0 in such subsets.

These two facts imply that $z = -\nu$ a.e. on $C \setminus C_*$. By modifying z in a set of null measure, we may assume that $z = -\nu$ on $C \setminus C_*$. We recall that the sets C_s , $s \geq s_*$ are minimizers of variational problems of the form $\min_{K \subseteq C} P(K) - \mu|K|$, for some values of μ (with $\mu = h_C$ as long as $s \leq s^*$ and $\mu = \mu_s > h_C$ continuously increasing with $s \geq s^*$). Since these sets are convex, with boundary (locally) uniformly of class $C^{1,1}$, and the map $s \rightarrow C_s$ is continuous in the Hausdorff topology, we obtain that the normal $\nu(x)$ is a continuous function in $C \setminus \text{int}(C_*)$.

Since $|z| < 1$ inside C_* and $[z \cdot \nu^K] = -1$ a.e. on ∂K , by [7, Theorem 1]) we have that the boundary of K must be outside the interior of C_* , hence either $K \supseteq C_*$ or $K \cap C_* = \emptyset$ (modulo a null set). Let us prove that the last situation is impossible. Indeed, assume that $K \cap C_* = \emptyset$ (modulo a null set). Since ∂K is of class C^1 out of a closed set of zero \mathcal{H}^{N-1} -measure (see [16]) and z is continuous in $C \setminus \text{int}(C_*)$, by Theorem 2 we have

$$z(x) \cdot \nu^K(x) = -1 \quad \mathcal{H}^{N-1}\text{-a.e. on } \partial K. \quad (23)$$

Now, since $K \cap C_* = \emptyset$ (modulo a null set), then there is some $s \geq s_*$ and some $x \in \partial C_s \cap \partial K$ such that $\nu^K(x) + \nu(x) = 0$. Fix $0 < \epsilon < 2$. By a slight perturbation, if necessary, we may assume that $x \in \partial C_s \cap \partial K$ with $s > s_*$, (23) holds at x and

$$|\nu^K(x) + \nu(x)| < \epsilon. \quad (24)$$

Since by (23) we have $\nu(x) = -z(x) = \nu^K(x)$ we obtain a contradiction with (24). We deduce that $K \supseteq C_*$. \square

Therefore, in order to prove uniqueness of the Cheeger sets of C , it is enough to show that

$$C_* = C^*. \quad (25)$$

Recall that the boundary of both C_* and C^* is of class $C^{1,1}$, and the sum of its principal curvatures is less than or equal h_C , and constantly equal to h_C in the interior of C . We now show that if $C_* \neq C^*$ and under additional assumptions, the sum of the principal curvatures of the boundary of C^* (or of any C_s for $s \in (s_*, s^*]$) must be h_C out of C_* .

Lemma 3.4. *Assume that C has C^2 boundary. Let $s \in (s_*, s^*]$ and $x \in \partial C_s \setminus \partial C_*$. If the sum of the principal curvatures of ∂C_s at x is strictly below h_C , then the Gaussian curvature of ∂C at x is 0.*

Proof. Let $x \in \partial C_s \setminus \partial C_*$ and assume the sum of the principal curvatures of ∂C_s at x is strictly below h_C (assuming x is a Lebesgue point for the curvature on ∂C_s). Necessarily, this implies that $x \in \partial C$. Assume then that the Gauss curvature of ∂C at x is positive: by continuity, in a neighborhood of x , C is uniformly convex and the sum of the principal curvatures is less than h_C . We may assume that near x , ∂C is the graph of a non-negative, C^2 and convex function $f : B \rightarrow \mathbb{R}$ where B is an $(N-1)$ -dimensional ball centered at x . We may as well assume that ∂C_s is the graph of $f_s : B \rightarrow \mathbb{R}$, which is $C^{1,1}$ [10, 22], and also nonnegative and convex. In B , we have $f_s \geq f \geq 0$, and

$$D^2 f \geq \alpha I \quad \text{and} \quad \operatorname{div} \frac{Df}{\sqrt{1+|Df|^2}} = h$$

with $h \in C^0(\overline{B})$, $h < h_C$, $\alpha > 0$, while

$$\operatorname{div} \frac{Df_s}{\sqrt{1+|Df_s|^2}} = h\chi_{\{f=f_s\}} + h_C\chi_{\{f_s>f\}}$$

(where $\chi_{\{f=f_s\}}$ has positive density at x).

We let $g = f_s - f \geq 0$. Introducing now the Lagrangian $\Psi : \mathbb{R}^{N-1} \rightarrow [0, +\infty)$ given by $\Psi(p) = \sqrt{1+|p|^2}$, we have that for a.e. $y \in B$

$$\begin{aligned} (h_C - h(y))\chi_{\{g>0\}}(y) &= \operatorname{div} (D\Psi(Df_s(y)) - D\Psi(Df(y))) \\ &= \operatorname{div} \left(\left(\int_0^1 D^2\Psi(Df(y) + t(Df_s(y) - Df(y))) dt \right) Dg(y) \right) \end{aligned}$$

so that, letting $A(y) := \int_0^1 D^2\Psi(Df(y) + tDg(y)) dt$ (which is a positive definite matrix and Lipschitz continuous inside B), we see that g is the minimizer of the functional

$$w \mapsto \int_B \left(A(y)Dw(y) \cdot Dw(y) + (h_C - h(y))w(y) \right) dy$$

under the constraint $w \geq 0$ and with boundary condition $w = f_s - f$ on ∂B . Adapting the results in [11] we get that $\{f = f_s\} = \{g = 0\}$ is the closure of a nonempty open set with boundary of zero \mathcal{H}^{N-1} -measure.

We therefore have found an open subset $D \subset \partial C \cap \partial C_s$, disjoint from ∂C_* , on which C is uniformly convex, with curvature less than h_C . Let φ be a smooth, nonnegative function with compact support in D . One easily shows that if $\varepsilon > 0$ is small enough, $\partial C_s - \varepsilon\varphi\nu^{C_s}$ is the boundary of a set C'_ε which is still convex, with $P(C'_\varepsilon)/|C'_\varepsilon| > P(C_s)/|C_s| = h_C$ (just differentiate the map $\varepsilon \rightarrow P(C'_\varepsilon)/|C'_\varepsilon|$), and the sum of its principal curvatures is less than h_C . This implies that for $\varepsilon > 0$ small enough, the set $C' := C'_\varepsilon$ is calibrable [3], which in turn implies that $\min_{K \subset C'} P(K)/|K| = P(C')/|C'|$. But this contradicts $C_* \subset C'$, which is true for ε small enough. \square

Proof of Theorem 3. Assume that C is C^2 and uniformly convex. Let us prove that its Cheeger set is unique. Assume by contradiction that $C^* \neq C_*$. From Lemma 3.4 we have that the sum of the principal curvatures of ∂C^* is h_C outside of C_* .

Let now $\bar{x} \in \partial C^* \cap \partial C_*$ be such that $\partial C^* \cap B_\rho(\bar{x}) \neq \partial C_* \cap B_\rho(\bar{x})$ for all $\rho > 0$ ($\partial C^* \cap \partial C_* \neq \emptyset$ since otherwise both C^* and C_* would be balls, which is impossible). Letting T be the tangent hyperplane to ∂C^* at \bar{x} , we can write ∂C^* and ∂C_* as the graph of two positive convex functions v^* and v_* , respectively, over $T \cap B_\rho(\bar{x})$ for $\rho > 0$ small enough. Identifying $T \cap B_\rho(\bar{x})$ with $B_\rho \subset \mathbb{R}^{N-1}$, we have that $v_*, v^* : B_\rho \rightarrow \mathbb{R}$ both solve the equation

$$-\operatorname{div} \frac{Dv}{\sqrt{1 + |Dv|^2}} = f, \quad (26)$$

for some function $f \in L^\infty(B_\rho)$. Moreover, it holds $v_* \geq v^*$, $v_*(0) = v^*(0)$ and $v_*(y) > v^*(y)$ for some $y \in B_\rho$. Notice that $f = \lambda_C$ in the (open) set where $v_* > v^*$, in particular both functions are smooth in this set. Let D be an open ball such that $\bar{D} \subset B_\rho$, $v_* > v^*$ on D and $v_*(y) = v^*(y)$ for some $y \in \partial D$. Notice that, since both v^* and v_* belong to $C^\infty(D) \cap C^1(\bar{D})$, the fact that $v_*(y) = v^*(y)$ also implies that $Dv_*(y) = Dv^*(y)$. In D , both functions solve (26) with $f = \lambda_C$. Letting now $w = v_* - v^*$, we have that $w(y) = 0$ and $Dw(y) = 0$, while $w > 0$ inside D . Recalling the function $\Psi(p) = \sqrt{1 + |p|^2}$, we have that for any $x \in D$

$$\begin{aligned} 0 &= \operatorname{div} (D\Psi(Dv_*(x)) - D\Psi(Dv^*(x))) \\ &= \operatorname{div} \left(\left(\int_0^1 D^2\Psi(Dv^*(x) + t(Dv_*(x) - Dv^*(x))) dt \right) Dw(x) \right) \end{aligned}$$

so that w solves a linear, uniformly elliptic equation with smooth coefficients. Then Hopf's lemma [13] implies that $Dw(y) \cdot \nu_D(y) < 0$, a contradiction. Hence $C_* = C^*$. \square

Remark 3.5. Notice that, as a consequence of Theorem 3 and the results of Giusti [14], we get that if C is of class C^2 and uniformly convex, equation (22) has a solution on the whole of C , if and only if C is a Cheeger set of itself, i.e. if and only if the the sum of the principal curvatures of ∂C is less than or equal to $P(C)/|C|$.

Remark 3.6. The results of this paper can be easily extended to the anisotropic setting (see [12]) provided the anisotropy is smooth and uniformly elliptic.

Acknowledgement. The first author acknowledges partial support by the Departament d'Universitats, Recerca i Societat de la Informació de la Generalitat de Catalunya and by PNP GC project, reference BFM2003-02125.

References

- [1] F. Alter. Uniqueness of the Cheeger set of a convex body. *In preparation*.
- [2] F. Alter, V. Caselles, A. Chambolle. Evolution of Convex Sets in the Plane by the Minimizing Total Variation Flow. *Interfaces and Free Boundaries* **7**, 29-53 (2005).
- [3] F. Alter, V. Caselles, A. Chambolle. A characterization of convex calibrable sets in \mathbb{R}^N . *Math. Ann.* **332**, 329-366 (2005).

- [4] L. Ambrosio. Corso introduttivo alla teoria geometrica della misura ed alle superfici minime. Scuola Normale Superiore, Pisa, 1997.
- [5] L. Ambrosio, N. Fusco, D. Pallara. Functions of Bounded Variation and Free Discontinuity Problems. Oxford Mathematical Monographs, 2000.
- [6] F. Andreu, C. Ballester, V. Caselles, J.M. Mazón. The Dirichlet Problem for the Total Variation Flow. *J. Funct. Anal.* **180** (2001), 347-403.
- [7] G. Anzellotti. Pairings between measures and bounded functions and compensated compactness. *Ann. Mat. Pura Appl.* **135** (1983), 293-318.
- [8] G. Anzellotti. Traces of bounded vector fields and the divergence theorem. Unpublished preprint (1983).
- [9] G. Bellettini, V. Caselles, M. Novaga. The Total Variation Flow in \mathbb{R}^N . *J. Differential Equations* **184**, 475-525 (2002).
- [10] H. Brézis and D. Kinderlehrer. The smoothness of solutions to nonlinear variational inequalities. *Indiana Univ. Math. J.* **23**, 831-844 (1973/74).
- [11] L.A. Caffarelli and N.M. Riviere. On the rectifiability of domains with finite perimeter. *Ann. Scuola Normale Superiore di Pisa* **3**, 177-186 (1976).
- [12] V. Caselles, A. Chambolle, S. Moll and M. Novaga. A characterization of convex calibrable sets in \mathbb{R}^N with respect to anisotropic norms. Preprint 2005.
- [13] D. Gilbarg and N.S. Trudinger. Elliptic partial Differential Equations of Second Order, Springer Verlag, 1998.
- [14] E. Giusti. On the equation of surfaces of prescribed mean curvature. Existence and uniqueness without boundary conditions. *Invent. Math.* **46**, 111-137 (1978).
- [15] E. Giusti. Boundary Value Problems for Non-Parametric Surfaces of Prescribed Mean Curvature. *Ann. Sc. Norm. Sup. Pisa (4)* **3**, 501-548 (1976).
- [16] E. Gonzalez, U. Massari, and I. Tamanini. On the regularity of sets minimizing perimeter with a volume constraint. *Indiana Univ. Math. Journal*, **32**, 25-37 (1983).
- [17] B. Kawohl, V. Fridman. Isoperimetric estimates for the first eigenvalue of the p -Laplace operator and the Cheeger constant. *Comment. Math. Univ. Carolinae* **44**, 659-667 (2003).
- [18] B. Kawohl, N. Kutev. Global behaviour of solutions to a parabolic mean curvature equation. *Differential and Integral Equations* **8**, 1923-1946 (1995).
- [19] B. Kawohl, T. Lachand-Robert. Characterization of Cheeger sets for convex subsets of the plane. *Pacific J. Math.*, to appear.

- [20] B. Kawohl, M. Novaga. The p -Laplace eigenvalue problem as $p \rightarrow 1$ and Cheeger sets in a Finsler metric. Preprint (2006).
- [21] A. Lichnerowski, R. Temam. Pseudosolutions of the Time Dependent Minimal Surface Problem. *J. Differential Equations* **30** (1978), 340-364.
- [22] E. Stredulinsky, W.P. Ziemer. Area Minimizing Sets Subject to a Volume Constraint in a Convex Set. *J. Geom. Anal.* **7**, 653-677 (1997).