

ECOLE POLYTECHNIQUE

CENTRE DE MATHÉMATIQUES APPLIQUÉES

UMR CNRS 7641

91128 PALAISEAU CEDEX (FRANCE). Tel: 01 69 33 46 00. Fax: 01 69 33 46 46

[http : //www.cmap.polytechnique.fr/](http://www.cmap.polytechnique.fr/)

**On the reconstruction of conductivity
of bordered two-dimensional surface in \mathbb{R}^3
from electrical current measurements on its boundary**

G.M. Henkin, R.G. Novikov

R.I. 678

Mars 2010

**On the reconstruction of conductivity
of bordered two-dimensional surface in \mathbb{R}^3
from electrical current measurements on its boundary**

G.M. Henkin¹ and R.G. Novikov²

¹Université Pierre et Marie Curie, case 247, 4 place Jussieu, 75252, Paris, France
e-mail: henkin@math.jussieu.fr

²CNRS (UMR 7641), Centre de Mathématiques Appliquées, Ecole Polytechnique,
91128, Palaiseau, France
e-mail: novikov@cmap.polytechnique.fr

Abstract.

An electrical potential U on a bordered real surface X in \mathbb{R}^3 with isotropic conductivity function $\sigma > 0$ satisfies equation $d(\sigma d^c U)|_X = 0$, where $d^c = i(\bar{\partial} - \partial)$, $d = \bar{\partial} + \partial$ are real operators associated with complex (conforme) structure on X induced by Eucliden metric of \mathbb{R}^3 . This paper gives exact reconstruction of conductivity function σ on X from Dirichlet-to-Neumann mapping $U|_{bX} \rightarrow \sigma d^c U|_{bX}$. This paper extends to the case of the Riemann surfaces the reconstruction schemes of R.Novikov [N2] and of A.Bukhgeim [B], given for the case $X \subset \mathbb{R}^2$. The paper extends and corrects the statements of [HM], where the inverse boundary value problem on the Riemann surfaces was firstly considered.

Keywords. Riemann surface. Electrical current. Inverse conductivity problem. $\bar{\partial}$ -method.

Mathematics Subject Classification (2000) 32S65, 32V20, 35R05, 35R30, 58J32, 81U40.

0. Introduction

0.1. Reduction of inverse boundary value problem on a surface in \mathbb{R}^3 to the corresponding problem on affine algebraic Riemann surface in \mathbb{C}^3 .

Let X be bordered oriented two-dimensional manifold in \mathbb{R}^3 . Manifold X is equipped by complex (conformal) structure induced by Euclidean metric of \mathbb{R}^3 . We say that X possesses an isotropic conductivity function $\sigma > 0$, if any electric potential u on bX generates electrical potential U on X , solving the Dirichlet problem:

$$U|_{bX} = u \quad \text{and} \quad d\sigma d^c U|_X = 0, \tag{0.1}$$

where $d^c = i(\bar{\partial} - \partial)$, $d = \bar{\partial} + \partial$ and the Cauchy-Riemann operator $\bar{\partial}$ corresponds to complex (conformal) structure on X . Inverse conductivity problem consists in the reconstruction of $\sigma|_X$ from the mapping potential $U|_{bX} \rightarrow$ current $j = \sigma d^c U|_{bX}$ for solutions of (0.1). This mapping is called Dirichlet-to-Neumann mapping.

This problem is the special case of the following more general inverse boundary value problem, going back to I.M.Gelfand [Ge] and A.Calderon [C]: to find potential (2-forme) q on X in the equation

$$dd^c \psi = q\psi \tag{0.2}$$

from knowledge of Dirichlet-to-Neumann mapping $\psi|_{bX} \rightarrow d^c\psi|_{bX}$ for solutions of (0.2). Equation (0.2) is called in some context by stationary Schrödinger equation, in other context by monochromatic acoustic equation etc. Equation (0.1) can be reduced to the equation (0.2) with

$q = \frac{dd^c\sqrt{\sigma}}{\sqrt{\sigma}}$ by the substitution $\psi = \sqrt{\sigma}U$.

Let restriction of Euclidean metric of \mathbb{R}^3 on X have (in local coordinates) the form

$$ds^2 = E dx^2 + 2F dx dy + G dy^2 = A dz^2 + 2B dz d\bar{z} + \bar{A} d\bar{z}^2,$$

where $z = x + iy$, $B = \frac{E+G}{4}$, $A = \frac{E-G-2iF}{4}$. Put $\mu = \frac{\bar{A}}{B + \sqrt{B^2 - |A|^2}}$. By classical results (going back to Gauss and Riemann) one can construct holomorphic embedding $\varphi : X \rightarrow \mathbb{C}^3$, using some solution of Beltrami equation: $\bar{\partial}\varphi = \mu\partial\varphi$ on X . Moreover, embedding φ can be chosen in such a way that $\varphi(X)$ belongs to smooth algebraic curve V in \mathbb{C}^3 . Using existence of embedding φ we can identify further X with $\varphi(X)$.

0.2. Reconstruction schemes for the case $X \subset \mathbb{R}^2 \simeq \mathbb{C}$.

For the case $X = \Omega \subset \mathbb{R}^2$ the exact reconstruction scheme for formulated inverse problems was given in [N2], [N3] under some restriction (smallness assumption) for σ or q (see Corollary 2 of [N2]). For the case of inverse conductivity problem, see (0.1), (0.2), when $q = \frac{dd^c\sqrt{\sigma}}{\sqrt{\sigma}}$, restriction on σ in this scheme was eliminated by A.Nachman [Na] by the reduction to the equivalent question for the first order system studied by R.Beals and R.Coifman [BC2]. Recently A.Bukhgeim [B] has found new original reconstruction scheme for inverse boundary value problem, see (0.2), without smallness assumption on q .

In a particular case, the scheme of [N2] for the inverse conductivity problem consists in the following. Let $\sigma(x) > 0$ for $x \in \bar{\Omega}$ and $\sigma \in C^{(2)}(\bar{\Omega})$. Put $\sigma(x) = 1$ for $x \in \mathbb{R}^2 \setminus \bar{\Omega}$.

Let $q = \frac{dd^c\sqrt{\sigma}}{\sqrt{\sigma}}$.

From L.Faddeev [F1] result it follows: \exists compact set $E \subset \mathbb{C}$ such that for each $\lambda \in \mathbb{C} \setminus E$ there exists a unique solution $\psi(z, \lambda)$ of the equation $dd^c\psi = q\psi = \frac{dd^c\sqrt{\sigma}}{\sqrt{\sigma}}\psi$, with asymptotics

$$\psi(z, \lambda)e^{-\lambda z} \stackrel{\text{def}}{=} \mu(z, \lambda) = 1 + o(1), \quad z \rightarrow \infty.$$

Such solution can be found from the integral equation

$$\mu(z, \lambda) = 1 + \frac{i}{2} \int_{\xi \in \Omega} g(z - \xi, \lambda) \frac{\mu(\xi, \lambda) dd^c\sqrt{\sigma}}{\sqrt{\sigma}}, \quad (0.3)$$

where the function

$$g(z, \lambda) = \frac{i}{(2\pi)^2} \int_{w \in \mathbb{C}} \frac{e^{\lambda w - \bar{\lambda} \bar{w}} dw \wedge d\bar{w}}{(w + z)\bar{w}} = \frac{i}{2(2\pi)^2} \int_{w \in \mathbb{C}} \frac{e^{i(w\bar{z} + \bar{w}z)} dw \wedge d\bar{w}}{w(\bar{w} - i\lambda)}$$

is called the Faddeev-Green function for the operator

$$\mu \mapsto \bar{\partial}(\partial + \lambda dz)\mu.$$

From [N2] it follows that $\forall \lambda \in \mathbb{C} \setminus E$ the function $\psi|_{b\Omega}$ can be found through Dirichlet-to-Neumann mapping by integral equation

$$\psi(z, \lambda)|_{b\Omega} = e^{\lambda z} + \int_{\xi \in b\Omega} e^{\lambda(z-\xi)} g(z-\xi, \lambda) (\hat{\Phi}\psi(\xi, \lambda) - \hat{\Phi}_0\psi(\xi, \lambda)), \quad (0.4)$$

where $\hat{\Phi}\psi = \bar{\partial}\psi|_{b\Omega}$, $\hat{\Phi}_0\psi = \bar{\partial}\psi_0|_{b\Omega}$, $\psi_0|_{b\Omega} = \psi|_{b\Omega}$ and $\partial\bar{\partial}\psi_0|_{\Omega} = 0$.

By results of [BC1], [GN] and [N2] it follows that $\psi(z, \lambda)$ satisfies $\bar{\partial}$ -equation of Bers-Vekua type with respect to $\lambda \in \mathbb{C} \setminus E$:

$$\frac{\partial\psi}{\partial\bar{\lambda}} = b(\lambda)\bar{\psi}, \quad \text{where} \quad (0.5)$$

$$\bar{\lambda}b(\lambda) = -\frac{1}{2\pi i} \int_{z \in b\Omega} e^{\lambda z - \bar{\lambda}\bar{z}} \bar{\partial}_z \mu(z, \lambda) = \frac{1}{4\pi} \int_{\Omega} e^{\lambda z - \bar{\lambda}\bar{z}} q\mu, \quad (0.6)$$

$$\psi(z, \lambda)e^{-\lambda z} = \mu(z, \lambda) \rightarrow 1, \quad \lambda \rightarrow \infty, \quad \forall z \in \mathbb{C}. \quad (0.7)$$

From [BC2] and [Na] it follows that for $q = \frac{dd^c\sqrt{\sigma}}{\sqrt{\sigma}}$, $\sigma > 0$, $\sigma \in C^{(2)}(\bar{\Omega})$ the exceptional set $E = \{\emptyset\}$ and function $\lambda \mapsto b(\lambda)$ belongs to $L^{2+\varepsilon}(\mathbb{C}) \cap L^{2-\varepsilon}(\mathbb{C})$ for some $\varepsilon > 0$. As a consequence function $\mu = e^{-\lambda z}\psi$ is a unique solution of the Fredholm integral equation

$$\mu(z, \lambda) + \frac{1}{2\pi} \int_{\lambda' \in \mathbb{C}} b(\lambda') e^{\bar{\lambda}'\bar{z} - \lambda'z} \bar{\mu}(z, \lambda') \frac{d\lambda' \wedge d\bar{\lambda}'}{\lambda' - \lambda} = 1. \quad (0.8)$$

Integral equations (0.4), (0.8) permit, starting from the Dirichlet-to-Neumann mapping, to find firstly the boundary values $\psi|_{b\Omega}$, secondly " $\bar{\partial}$ -scattering data" $b(\lambda)$ and thirdly function $\psi|_{\Omega}$. From equality $dd^c\psi = \frac{dd^c\sqrt{\sigma}}{\sqrt{\sigma}}\psi$ on X we find finally $\frac{dd^c\sqrt{\sigma}}{\sqrt{\sigma}}$ on X .

The scheme of the Bukhgeim type [B] can be presented in the following way. Let $q = Qdd^c|z|^2$, where $Q \in C^{(1)}(\bar{\Omega})$, but potential Q is not necessary of the conductivity form $\frac{dd^c\sqrt{\sigma}}{\sqrt{\sigma}}$. By variation of Faddeev statement and proof we obtain that $\forall a \in \mathbb{C} \exists$ compact set $E \subset \mathbb{C}$ such that $\forall \lambda \in \mathbb{C} \setminus E$ there exists a unique solution $\psi_a(z, \lambda)$ of the equation $dd^c\psi = q\psi$ with asymptotics

$$\psi_a(z, \lambda)e^{-\lambda(z-a)^2} = \mu_a(z, \lambda) = 1 + o(1), \quad z \rightarrow \infty.$$

Such a solution can be found from integral equation (0.3), where kernel $g(z, \lambda)$ is replaced by kernel

$$g_a(z, \zeta, \lambda) = \frac{ie^{\lambda a^2 - \bar{\lambda}\bar{a}^2}}{2\pi^2} \int_{\mathbb{C}} \frac{e^{-\lambda(\zeta-\eta+a)^2 + \bar{\lambda}(\bar{\zeta}-\bar{\eta}+\bar{a})^2}}{(\eta-z)(\bar{\zeta}-\bar{\eta})} d\eta \wedge d\bar{\eta}.$$

Kernel $g_a(z, \zeta, \lambda)$ can be called the Faddeev type Green function for the operator

$\mu \rightarrow \bar{\partial}(\partial + \lambda d(z-a)^2)\mu$. Equation $\bar{\partial}(\partial + \lambda d(z-a)^2)\mu = \frac{i}{2}q\mu$ and Green formula implies

$$\int_{b\Omega} e^{\lambda(z-a)^2 - \bar{\lambda}(\bar{z}-\bar{a})^2} \bar{\partial}\mu = \int_{\Omega} e^{\lambda(z-a)^2 - \bar{\lambda}(\bar{z}-\bar{a})^2} \frac{q\mu}{2i}. \quad (0.9)$$

Stationary phase method, applied to the integral in the right-hand side of (0.9), gives for $\tau \rightarrow \infty$, $\tau \in \mathbb{R}$, equality

$$\lim_{\substack{\tau \rightarrow \infty \\ \tau \in \mathbb{R}}} \frac{4\tau}{\pi i} \int_{z \in b\Omega} e^{i\tau[(z-a)^2 + (\bar{z}-\bar{a})^2]} \bar{\partial}_z \mu_a(z, i\tau) = Q(a). \quad (0.10)$$

Formula (0.10) means that values of potential Q in the arbitrary point a of Ω can be reconstructed from Dirichlet-to-Neumann mapping $\mu_a|_{b\Omega} \mapsto \bar{\partial}_z \mu_a|_{b\Omega}$ for family of functions $\mu_a(z, \lambda)$ depending on parameter $\lambda = i\tau$, $\tau > \text{const}$, where we assume that $\mu_a|_{b\Omega}$ is found using an analog of (0.4) for $\psi_a|_{b\Omega}$.

Bukhgeim's scheme works well at least $\forall Q \in C^1(\bar{\Omega})$.

More constructive scheme of [N2] works quite well only in the absence of exceptional set E in the λ -plane for Faddeev type functions. In papers [BLMP], [Ts], [N3] it was constructed modified Faddeev-Green function permitting to solve inverse boundary problem (0.2), on the $\mathbb{R}^2 = \mathbb{C}$, at least, under some smallness assumptions on potential Q .

Let us note that the first uniqueness results in the two-dimensional inverse boundary value or scattering problems for (0.1) or (0.2) goes back to A.Calderon [C], V.Druskin [D], R.Kohn, M.Vogelius [KV], J.Sylvester, G.Uhlmann [SU] and R.Novikov [N1].

Note in this connection that the first seminal results on reconstruction of the two-dimensional Schrödinger operator H on the torus from the data "extracted" from the family of eigenfunctions (Bloch-Floquet) of single energy level $H\psi = E\psi$ were obtained in series of papers starting from B.Dubrovin, I.Krichever, S.P.Novikov [DKN], S.P.Novikov, A.Veselov [NV]. These results were obtained in connection with (2+1)- dimensional evolution equations.

This paper extends to the case of Riemann surfaces reconstruction procedures of [N2] and of [B]. The paper extends (and also corrects) the recent paper [HM2] where the inverse boundary value problem on Riemann surface was firstly considered. Earlier in [HM1] it was proved that if $X \subset \mathbb{R}^3$ possesses a constant conductivity then X with complex structure can be effectively reconstructed by at most three generic potential \rightarrow current measurements on bX .

Very recently, motivated by [B] and [HM1], [HM2], C.Guillarmou and L.Tzou [GT] have obtained general identifiability result (without reconstruction procedure): if for all solutions of equations $dd^c u + q_j u = 0$, $q_j \in C^{(2)}(X)$, $j = 1, 2$, Cauchy datas $u|_{bX}$, $d^c u|_{bX}$, coincide, then $q_1 = q_2$ on X .

1. Preliminaries and main results

Let $\mathbb{C}P^3$ be complex projective space with homogeneous coordinates

$w = (w_0 : w_1 : w_2 : w_3)$. Let $\mathbb{C}P_\infty^2 = \{w \in \mathbb{C}P^3 : w_0 = 0\}$. Then $\mathbb{C}P^3 \setminus \mathbb{C}P_\infty^2$ can be considered as the complex affine space with coordinates $z_k = w_k/w_0$, $k = 1, 2, 3$. By classical result of G. Halphen (see R.Hartshorne [H], ch.IV, § 6) any compact Riemann surface of genus g can be embedded in $\mathbb{C}P^3$ as projective algebraic curve \tilde{V} , which intersects $\mathbb{C}P_\infty^2$ transversally in $d > g$ points, where $d \geq 1$ if $g = 0$, $d \geq 3$ if $g = 1$ and $d \geq g + 3$ if $g \geq 2$. Without loss of generality one can suppose that

- i) $V = \tilde{V} \setminus \mathbb{C}P_\infty^2$ is connected affine algebraic curve in \mathbb{C}^3 defined by polynomial equations $V = \{z \in \mathbb{C}^3 : p_1(z) = p_2(z) = p_3(z) = 0\}$ such that the rang of the matrix $[\frac{\partial p_1}{\partial z}(z), \frac{\partial p_2}{\partial z}(z), \frac{\partial p_3}{\partial z}(z)] \equiv 2 \forall z \in V$.
- ii) $\tilde{V} \cap \mathbb{C}P_\infty^2 = \{\beta_1, \dots, \beta_d\}$, where

$$\beta_l = (0 : \beta_l^1 : \beta_l^2 : \beta_l^3), \left(\frac{\beta_l^2}{\beta_l^1}, \frac{\beta_l^3}{\beta_l^1} \right) \in \mathbb{C}^2, l = 1, 2, \dots, d.$$

- iii) For $r_0 > 0$ large enough

$$\det \begin{vmatrix} \frac{\partial p_\alpha}{\partial z_2} & \frac{\partial p_\alpha}{\partial z_3} \\ \frac{\partial p_\beta}{\partial z_2} & \frac{\partial p_\beta}{\partial z_3} \end{vmatrix} \neq 0 \text{ for } z \in V : |z_1| \geq r_0 \text{ and } \alpha \neq \beta.$$

- iv) For $|z|$ large enough:

$$\frac{dz_2}{dz_1}|_{V_l} = \gamma_l + \frac{\gamma_l^0}{z_1^2} + O\left(\frac{1}{z_1^3}\right), \quad \frac{dz_3}{dz_1}|_{V_l} = \tilde{\gamma}_l + \frac{\tilde{\gamma}_l^0}{z_1^2} + O\left(\frac{1}{z_1^3}\right),$$

where $\gamma_l, \tilde{\gamma}_l, \gamma_l^0, \tilde{\gamma}_l^0 \neq 0$, for $l = 1, \dots, d$, $d \geq 2$.

Let $V_0 = \{z \in V : |z_1| \leq r_0\}$ and $V \setminus V_0 = \cup_{l=1}^d V_l$, where $\{V_l\}$ are connected components of $V \setminus V_0$. Let us equip V by Euclidean volume form $dd^c|z|^2$. Let $\tilde{W}^{1, \tilde{p}}(V) = \{F \in L^\infty(V) : \bar{\partial}F \in L_{0,1}^{\tilde{p}}(V)\}$, $\tilde{W}_{1,0}^{1, \tilde{p}}(V) = \{f \in L_{1,0}^\infty(V) : \bar{\partial}f \in L_{1,1}^{\tilde{p}}(V)\}$, $\tilde{p} > 2$. Let $H_{0,1}(V)$ denotes the space of antiholomorphic $(0,1)$ -forms on V . Let $H_{0,1}^p(V) = H_{0,1}(V) \cap L_{0,1}^p(V)$, $1 < p < 2$.

Let $W^{1,p}(V) = \{F \in L^p(V) : \bar{\partial}F \in L_{0,1}^p(V)\}$.

From the Hodge-Riemann decomposition theorem (see [GH], [Ho]) $\forall \Phi_0 \in W_{0,1}^{1,p}(\tilde{V})$ we have

$\Phi_0 = \bar{\partial}(\bar{\partial}^*G\Phi_0) + \mathcal{H}\Phi_0$, where $\mathcal{H}\Phi_0 \in H_{0,1}(\tilde{V})$ and G is the Hodge-Green operator for the Laplacian $\bar{\partial}\bar{\partial}^* + \bar{\partial}^*\bar{\partial}$ on \tilde{V} with the properties: $G(H_{0,1}(\tilde{V})) = 0$, $\bar{\partial}G = G\bar{\partial}$, $\bar{\partial}^*G = G\bar{\partial}^*$.

Straight generalization of Proposition 1 from [He] gives explicit operators:

$R_1 : L_{0,1}^p(V) \rightarrow L^{\tilde{p}}(V)$, $R_0 : L_{0,1}^p(V) \rightarrow \tilde{W}_{1,0}^{1, \tilde{p}}(V)$ and $\mathcal{H} : L_{0,1}^p(V) \rightarrow H_{0,1}^p(V)$, $1 < p < 2$, $\frac{1}{\tilde{p}} = \frac{1}{p} - \frac{1}{2}$, such that $\forall \Phi \in L_{0,1}^p(V)$ we have decomposition of Hodge-Riemann type:

$$\Phi = \bar{\partial}R\Phi + \mathcal{H}\Phi, \text{ where } R = R_1 + R_0,$$

$$R_1\Phi(z) = \frac{1}{2\pi i} \int_{\xi \in V} \Phi(\xi) \wedge (dp_\alpha \wedge dp_\beta) \rfloor d\xi_1 \wedge d\xi_2 \wedge d\xi_3 \det \left[\frac{\partial p_\alpha(\xi)}{\partial \xi}, \frac{\partial p_\beta(\xi)}{\partial \xi}, \frac{\bar{\xi} - \bar{z}}{|\xi - z|^2} \right],$$

$$R_0\Phi(z) = (\bar{\partial}^*G(\bar{\partial}R_1\Phi - \Phi))(z) - (\bar{\partial}^*G(\bar{\partial}R_1\Phi - \Phi))(\beta_1),$$

$(\bar{\partial}R_1\Phi - \Phi) \in W_{0,1}^{1,p}(\tilde{V})$, G is the Hodge - Green operator for Laplacian $\bar{\partial}\bar{\partial}^*$

for $(0,1)$ - forms on \tilde{V} ,

(1,1)-form under sign of integral does not depend on the choice of indexes $\alpha, \beta = 1, 2, 3$, $\alpha \neq \beta$,

$$\mathcal{H}\Phi = \sum_{j=1}^g \left(\int_V \Phi \wedge \omega_j \right) \bar{\omega}_j,$$

$\{\omega_j\}$ is orthonormal basis of holomorphic (1,0)-forms on \tilde{V} , i.e.

$$\int_V \omega_j \wedge \bar{\omega}_k = \delta_{jk}, \quad j, k = 1, 2, \dots, g.$$

Note that as a corollary of construction of R we have that $\lim_{\substack{z \in V_1 \\ z \rightarrow \infty}} R\Phi(z) = R\Phi(\beta_1) = 0$.

Remark 1.1. If $V = \{z \in \mathbb{C}^2 : P(z) = 0\}$ be algebraic curve in \mathbb{C}^2 then formula for operator R_1 is reduced to the following:

$$R_1\Phi(z) = \frac{1}{2\pi i} \int_{\xi \in V} \Phi(\xi) \frac{d\xi_1}{\partial \xi_2} \det \left[\frac{\partial P}{\partial \xi}(\xi), \frac{\bar{\xi} - \bar{z}}{|\xi - z|^2} \right].$$

Remark 1.2. Based on [HP] one can construct an explicit formula not only for the main part R_1 of the R -operator, but for the whole operator $R = R_1 + R_0$.

Let $\varphi \in L_{1,1}^1(V) \cap L_{1,1}^\infty(V)$, $f \in \tilde{W}_{1,0}^{1,\tilde{p}}(V)$, $\lambda \in \mathbb{C}$, $\theta \in \mathbb{C}$.

Let

$$\begin{aligned} \hat{R}_\theta \varphi &= R((dz_1 + \theta dz_2) \rfloor \varphi)(dz_1 + \theta dz_2), \\ R_{\lambda,\theta} f &= e_{-\lambda,\theta} \overline{R(e_{\lambda,\theta} f)}, \quad \text{where } e_{\lambda,\theta}(z) = e^{\lambda(z_1 + \theta z_2) - \bar{\lambda}(\bar{z}_1 + \bar{\theta} \bar{z}_2)}. \end{aligned}$$

By straight generalization of Propositions 2, 3 from [He] the form $f = \hat{R}_\theta \varphi$ is a solution of $\bar{\partial} f = \varphi$ on V , function $u = R_{\lambda,\theta} f$ is a solution of

$$\begin{aligned} (\partial + \lambda(dz_1 + \theta dz_2))u &= f - \mathcal{H}_{\lambda,\theta} f, \quad \text{where} \\ \mathcal{H}_{\lambda,\theta} f &\stackrel{\text{def}}{=} e_{-\lambda,\theta} \overline{\mathcal{H}(e_{\lambda,\theta} f)}, \quad u \in W^{1,\tilde{p}}(V), \quad \tilde{p} > 2. \end{aligned}$$

In addition, by straight generalization of Proposition 4 from [He] we have that

$$\bar{\partial}(\partial + \lambda(dz_1 + \theta dz_2))u = \varphi + \bar{\lambda}(d\bar{z}_1 + \bar{\theta} d\bar{z}_2) \wedge \mathcal{H}_{\lambda,\theta}(\hat{R}_\theta \varphi) \quad \text{on } V.$$

Definition 1.1. The kernel $g_{\lambda,\theta}(z, \xi)$, $z, \xi \in V$, $\lambda \in \mathbb{C}$, of integral operator $R_{\lambda,\theta} \circ \hat{R}_\theta$ is called in [He] the Faddeev type Green function for operator $\bar{\partial}(\partial + \lambda(dz_1 + \theta dz_2))$.

Definition 1.2. Let $g = \text{genus } \tilde{V}$. Let $\{\omega_j\}$, $j = 1, \dots, g$, be orthonormal basis of holomorphic forms on \tilde{V} . Let $\{a_1, \dots, a_g\}$ be different points (or effective divisor) on $V \setminus V_0$. Let

$$\Delta_\theta(\lambda) = \det \left[\int_{\xi \in V} \hat{R}_\theta(\delta(\xi, a_j)) \wedge \bar{\omega}_k(\xi) e_{\lambda,\theta}(\xi), \quad j, k = 1, \dots, g, \right]$$

where $\delta(\xi, a_j)$ - Dirac (1,1)-form concentrated in $\{a_j\}$.

Let $E_\theta = \{\lambda \in \mathbb{C} : \Delta_\theta(\lambda) = 0\}$.

Definition 1.3. Parameter $\theta \in \mathbb{C}$ will be called generic if $\theta \notin \{\theta_1, \dots, \theta_d\}$, where $\theta_l = -1/\gamma_l$. Divisor $\{a_1, \dots, a_g\}$ on $V \setminus V_0$ will be called generic if

$$\det \left[\frac{\omega_j}{dz_1}(a_k) \right]_{j,k=1, \dots, g} \neq 0.$$

Proposition 1.1. Let parameter $\theta \in \mathbb{C}$ and divisor $\{a_1, \dots, a_g\}$ on $V \setminus V_0$ be generic, where $V_0 = \{z \in V : |z_1| \leq r_0\}$, $g \geq 1$. Then for r_0 large enough we have inequalities:

$$\overline{\lim}_{\lambda \rightarrow \infty} |\lambda^g \Delta_\theta(\lambda)| < \infty \quad \text{and}$$

$$\forall \varepsilon > 0 \quad \underline{\lim}_{\lambda \rightarrow \infty} |\lambda^g \Delta_\theta(\lambda)|_\varepsilon > 0, \quad \text{where} \quad |\lambda^g \Delta_\theta(\lambda)|_\varepsilon = \sup_{\{\lambda' : |\lambda' - \lambda| < \varepsilon\}} |(\lambda')^g \cdot \Delta_\theta(\lambda')|$$

Besides, the set E_θ is a closed nowhere dense subset of \mathbb{C} .

Let X be domain containing V_0 and relatively compact on V . Let $\sigma \in C^{(3)}(V)$, $\sigma > 0$, on V , $\sigma = 1$ on $V \setminus X$. Let Y be domain containing \bar{X} and relatively compact on V . Let divisor $\{a_1, \dots, a_g\}$ on $Y \setminus X$ and parameter $\theta \in \mathbb{C}$ be generic.

Definition 1.4. The functions $\psi_\theta(z, \lambda) = \sqrt{\sigma} F_\theta(z, \lambda) = \mu_\theta(z, \lambda) e^{\lambda(z_1 + \theta z_2)}$, $z \in V$, $\theta \in \mathbb{C} \setminus \{\theta_1, \dots, \theta_d\}$, $\lambda \in \mathbb{C} \setminus E_\theta$, will be called the Faddeev type functions, associated with σ , θ and $\{a_1, \dots, a_g\}$ if ψ_θ , F_θ , μ_θ satisfy correspondingly properties:

$$\begin{aligned} d\sigma d^c F_\theta &= 2\sqrt{\sigma} e^{\lambda(z_1 + \theta z_2)} \sum_{j=1}^g C_{j,\theta}(\lambda) \delta(z, a_j), \\ dd^c \psi_\theta &= q\psi_\theta + 2e^{\lambda(z_1 + \theta z_2)} \sum_{j=1}^g C_{j,\theta}(\lambda) \delta(z, a_j), \\ \bar{\partial}(\partial + \lambda(dz_1 + \theta dz_2))\mu_\theta &= \frac{i}{2} q\mu_\theta + i \sum_{j=1}^g C_{j,\theta}(\lambda) \delta(z, a_j), \end{aligned} \tag{1.1}$$

and the normalization condition

$$\lim_{\substack{z \in V_1 \\ z \rightarrow \infty}} \mu_\theta(z, \lambda) = 1, \tag{1.2}$$

where $\mu_\theta|_Y \in L^{\tilde{p}}(Y)$, $\mu_\theta|_{V \setminus Y} \in L^\infty(V \setminus Y)$, $\tilde{p} > 2$, $q = \frac{dd^c \sqrt{\sigma}}{\sqrt{\sigma}}$, $\{C_{j,\theta}\}$ are some functions of $\lambda \in \mathbb{C} \setminus E_\theta$.

Theorem 1.1. Under the aforementioned notations and conditions, \forall generic $\theta \in \mathbb{C}$, \forall generic divisor $\{a_1, \dots, a_g\} \subset V \setminus X$ and $\forall \lambda \in \mathbb{C} \setminus E_\theta : |\lambda| > \text{const}(V, \{a_j\}, \theta, \sigma)$ there exists unique Faddeev type function

$$\psi_\theta(z, \lambda) = \sqrt{\sigma} F_\theta(z, \lambda) = e^{\lambda(z_1 + \theta z_2)} \mu_\theta(z, \lambda),$$

associated with conductivity function σ and divisor $\{a_1, \dots, a_g\}$. Moreover:

A) function $z \rightarrow \psi_\theta(z, \lambda)$ and parameters $\{C_{j,\theta}(\lambda)\}$ can be found from the following equations, depending on parameters $\theta \in \mathbb{C}$, $\lambda \in \mathbb{C} \setminus E_\theta$,

$$\begin{aligned} \psi_\theta(z, \lambda) - \frac{i}{2} \int_{\xi \in X} e^{\lambda((z_1 - \xi_1) + \theta(z_2 - \xi_2))} g_{\lambda,\theta}(z, \xi) \frac{dd^c \sqrt{\sigma}}{\sqrt{\sigma}} \psi_\theta(z, \lambda) = \\ e^{\lambda(z_1 + \theta z_2)} + i \sum_{j=1}^g C_{j,\theta}(\lambda) g_{\lambda,\theta}(z, a_j) e^{\lambda(z_1 + \theta z_2)}, \end{aligned} \quad (1.3)$$

$$\begin{aligned} 2 \sum_{j=1}^g C_{j,\theta}(\lambda) e_{\lambda,\theta}(a_j) \frac{\bar{\omega}_k}{d\bar{z}_1 + \theta d\bar{z}_2}(a_j) = \\ - \int_{z \in V} e^{-\bar{\lambda}(\bar{z}_1 + \bar{\theta}\bar{z}_2)} \frac{dd^c \sqrt{\sigma}}{\sqrt{\sigma}} \psi_\theta(z, \lambda) \frac{\bar{\omega}_k}{d\bar{z}_1 + \theta d\bar{z}_2}(z), \end{aligned} \quad (1.4)$$

where $k = 1, 2, \dots, g$ and $\{\omega_j\}$ is orthonormal basis of holomorphic forms on \tilde{V} ;

B) functions $z \rightarrow \psi_\theta(z, \lambda)$ and parameters $\{C_{j,\theta}(\lambda)\}$ satisfy the following properties for $\lambda \in \mathbb{C} \setminus E_\theta$: $|\lambda| \geq \text{const}(V, \{a_j\}, \theta, \sigma)$

$$\exists \lim_{\substack{z \rightarrow \infty, \\ i=1,2,\dots,d}} \frac{\bar{z}_1 + \bar{\theta}\bar{z}_2}{\bar{\lambda}} e^{-\bar{\lambda}(\bar{z}_1 + \bar{\theta}\bar{z}_2)} \left(\frac{\partial \psi_\theta}{\partial \bar{z}_1} + \bar{\theta} \frac{\partial \psi_\theta}{\partial \bar{z}_2} \right) = \lim_{\substack{z \rightarrow \infty \\ z \in V_i}} \psi_\theta e^{-\lambda(z_1 + \theta z_2)} b_\theta(\lambda), \quad (1.5)$$

$$iC_{j,\theta}(\lambda) = (2\pi i) \text{Res}_{a_j} e^{-\lambda(z_1 + \theta z_2)} \partial \psi_\theta \stackrel{\text{def}}{=} 2\pi i \lim_{\varepsilon \rightarrow 0} \int_{|z - a_j| = \varepsilon} e^{-\lambda(z_1 + \theta z_2)} \partial \psi_\theta, \quad (1.6)$$

$$\frac{\partial \psi_\theta(z, \lambda)}{\partial \bar{\lambda}} = b_\theta(\lambda) \overline{\psi_\theta(z, \lambda)}, \quad (1.7)$$

$$\frac{\partial C_{j,\theta}(\lambda)}{\partial \bar{\lambda}} e^{\lambda(a_{j,1} + \theta a_{j,2})} = b_\theta(\lambda) \overline{C_{j,\theta}(\bar{\lambda})} e^{\bar{\lambda}(\bar{a}_{j,1} + \bar{\theta}\bar{a}_{j,2})}. \quad (1.8)$$

Besides,

$$\begin{aligned} \bar{\lambda} b_\theta(\lambda) d = -\frac{1}{2\pi i} \int_{z \in bX} e_{\lambda,\theta}(z) \bar{\partial} \mu(z) + i \sum_{j=1}^g C_{j,\theta} e_{\lambda,\theta}(a_j), \\ |\lambda| \cdot |b_\theta(\lambda)| \leq \text{const}(V, \{a_j\}, \sigma) \frac{1}{(|\lambda| + 1)^{1/3}} \frac{1}{|\Delta_\theta(\lambda)|(1 + |\lambda|)^g}, \\ |C_{j,\theta}(\lambda)| \leq \text{const}(V, \{a_j\}, \sigma) \frac{1}{(|\lambda| + 1)^{1/3}} \frac{1}{|\Delta_\theta(\lambda)|(1 + |\lambda|)^g}. \end{aligned} \quad (1.9)$$

Remark 1.3. If $\|\ln \sqrt{\sigma}\|_{C^{(2)}(X)} \leq \text{const}(V, \{a_j\}, \theta)$ then the condition $\lambda \in \mathbb{C} \setminus E_\theta$: $|\lambda| \geq \text{const}(V, \{a_j\}, \theta, \sigma)$ in Theorem 1.1 can be replaced by the condition $\lambda \in \mathbb{C} \setminus E_\theta$. Dependence of $\text{const}(V, \{a_j\}, \theta, \sigma)$ of σ means its dependence only of $\|\ln \sqrt{\sigma}\|_{C^{(2)}(X)}$.

Definition 1.5. The functions $b_\theta(\lambda)$ and $\{C_{j,\theta}\}$ will be called "scattering" data for potential q .

Let $\hat{\Phi}(\psi|_{bX}) = \bar{\partial}\psi|_{bX}$ for all sufficiently regular solutions ψ of (0.2) in \bar{X} , where $q = \frac{dd^c\sqrt{\sigma}}{\sqrt{\sigma}}$. The operator $\hat{\Phi}$ is equivalent to the Dirichlet-to-Neumann operator for (0.1). Let $\hat{\Phi}_0$ denote $\hat{\Phi}$ for $q \equiv 0$ on \bar{X} .

Theorem 1.2. Under the conditions of Proposition 1.1 and Theorem 1.1, the following statements are valid:

A) $\forall \lambda \in \mathbb{C} \setminus E_\theta : |\lambda| \geq \text{const}(V, \{a_j\}, \theta, \sigma)$ the restriction of $\psi_\theta(z, \lambda)$ on bX and data $\{C_{j,\theta}(\lambda)\}$ can be reconstructed from Dirichlet-to-Neumann data as unique solution of the Fredholm integral equation

$$\psi_\theta(z, \lambda)|_{bX} + \int_{\xi \in bX} e^{\lambda[(z_1 - \xi_1) + \theta(z_2 - \xi_2)]} g_{\lambda,\theta}(z, \xi) (\hat{\Phi} - \hat{\Phi}_0) \psi_\theta(\xi, \lambda) = \quad (1.10)$$

$$e^{\lambda(z_1 + \theta z_2)} + i \sum_{j=1}^g C_{j,\theta}(\lambda) g_{\lambda,\theta}(z, a_j) e^{\lambda(z_1 + \theta z_2)}, \quad \text{where}$$

$$\int_{z \in bX} (z_1 + \theta z_2)^{-k} (\partial + \lambda(dz_1 + \theta dz_2)) \mu_\theta(z, \lambda) = - \sum_{j=1}^g (a_{j,1} + \theta a_{j,2})^{-k} C_{j,\theta}(\lambda), \quad (1.11)$$

$k = 2, \dots, g+1$, where (without restriction of generality) we suppose that values $\{a_{j,1}\}$ of the first coordinates of points $\{a_j\}$ are mutually different;

B) Function $\sigma(w)$, $w \in X$, can be reconstructed from Dirichlet-to-Neumann data

$$\psi_\theta|_{bX} \stackrel{\text{def}}{=} \mu_\theta|_{bX} e^{\lambda(z_1 + \theta z_2)} \rightarrow \bar{\partial}\psi_\theta|_{bX}$$

by explicit formulas, where we assume that $\psi_\theta|_{bX}$ is found using (1.10), (1.11).

For the case $V = \{z \in \mathbb{C}^2 : P(z) = 0\}$, where P is a polynomial of degree N , this formula has the following form. Let $\{w_m\}$ be points of V , where $(dz_1 + \theta dz_2)|_V(w_m) = 0$, $m = 1, \dots, M$. Then for almost all θ values $\frac{dd^c\sqrt{\sigma}}{\sqrt{\sigma}dd^c|z|^2}|_V(w_m)$ can be found from the following linear system

$$\tau(1 + o(1)) \frac{d^k}{d\tau^k} \left(\int_{z \in bX} e_{i\tau,\theta}(z) \bar{\partial}\mu_\theta(z, i\tau) \right) =$$

$$\sum_{m=1}^M \frac{i\pi(1 + |\theta|^2)}{2} \frac{dd^c\sqrt{\sigma}}{\sqrt{\sigma}dd^c|z|^2} \Big|_V(w_m) \times \quad (1.12)$$

$$\frac{|\frac{\partial P}{\partial z_1}(w)|^3 \frac{d^k}{d\tau^k} \exp i\tau[(w_{m,1} + \theta w_{m,2}) + (\bar{w}_{m,1} + \bar{\theta} \bar{w}_{m,2})]}{|\frac{\partial^2 P}{\partial z_1^2} (\frac{\partial P}{\partial z_2})^2 - 2 \frac{\partial^2 P}{\partial z_1 \partial z_2} (\frac{\partial P}{\partial z_2}) (\frac{\partial P}{\partial z_1}) + \frac{\partial^2 P}{\partial z_2^2} (\frac{\partial P}{\partial z_1})^2| (w_m)},$$

where $m, k = 1, \dots, M$; $M = N(N-1)$, $\tau \in \mathbb{R}$, $\tau \rightarrow \infty$, $|\tau|^g |\Delta_\theta(i\tau)| \geq \varepsilon > 0$, ε -small enough. Determinant of system (1.12) is proportional to the determinant of Vandermonde.

C) If $g = 0$ and if $\theta = \theta(\lambda) = \lambda^{-2}$, then $\forall z \in X$ and $\forall \lambda \in \mathbb{C}$ function $\mu_\theta(z, \lambda) = \psi_\theta(z, \lambda)e^{-\lambda(z_1 + \theta z_2)}$ is unique solution of Fredholm integral equation

$$\mu_{\theta(\lambda)}(z, \lambda) + \frac{1}{2\pi i} \int_{\xi \in \mathbb{C}} b_{\theta(\xi)}(\xi) e^{\bar{\xi}(z_1 + \bar{\theta}(\xi)z_2) - \xi(z_1 + \theta(\xi)z_2)} \bar{\mu}_{\theta(\xi)}(z, \xi) \frac{d\xi \wedge d\bar{\xi}}{\xi - \lambda} = 1,$$

$$\text{where } |b_{\theta(\xi)}(\xi)| \leq \frac{\text{const}(V)}{(1 + |\xi|)^2},$$

and function $z \rightarrow \sigma(z)$, $z \in X$, can be found from equality

$$dd^c \psi_{\theta(\lambda)}(z, \lambda) = \frac{dd^c \sqrt{\sigma}}{\sqrt{\sigma}}(z) \psi_{\theta(\lambda)}(z, \lambda), \quad z \in X.$$

Remark 1.4. Using the Faddeev type Green function constructed in [He], in [HM2] were obtained natural analogues of the main steps of the reconstruction scheme of [N2] on the Riemann surface V . In particular, under a smallness assumption on $\partial \log \sqrt{\sigma}$ the existence (and uniqueness) of the solution $\mu(z, \lambda)$ of the Faddeev type integral equation

$$\mu_\theta(z, \lambda) = 1 + \frac{i}{2} \int_{\xi \in V} g_{\lambda, \theta}(z, \xi) \frac{\mu_\theta(\xi, \lambda) dd^c \sqrt{\sigma}}{\sqrt{\sigma}} + i \sum_{j=1}^g C_j g_{\lambda, \theta}(z, a_j), \quad z \in V, \quad \lambda \in \mathbb{C},$$

holds for any a priori fixed constants C_1, \dots, C_g . However (and this fact was overlooked in [HM]) for $\lambda \in \mathbb{C} \setminus E$ there exists unique choice of constants $C_j(\lambda, \sigma)$ for which the integral equation above is equivalent to the differential equation

$$\bar{\partial}(\partial + \lambda(dz_1 + \theta dz_2))\mu - \frac{i}{2} \left(\frac{dd^c \sqrt{\sigma}}{\sqrt{\sigma}} \mu \right) + i \sum_{j=1}^g C_j \delta(z, a_j),$$

where $\delta(z, a_j)$ are Dirac measures concentrated in the points a_j .

2. Faddeev type functions on Riemann surfaces. Uniqueness

Let projective algebraic curve \tilde{V} be embedded in $\mathbb{C}P^3$ and intersect $\mathbb{C}P_\infty^2 = \{w \in \mathbb{C}P^3 : w_0 = 0\}$ transversally in $d > g$ points. Let $V = \tilde{V} \setminus \mathbb{C}P_\infty^2$, $V_0 = \{z \in V : |z_1| \leq r_0\}$ and properties i)-iv) from § 1 be valid.

Proposition 2.1. *Let σ be positive function belonging to $C^{(2)}(V)$ such that $\sigma \equiv \text{const} = 1$ on $V \setminus X \subset V \setminus V_0 = \cup_{l=1}^d V_l$, where $\{V_l\}$ are connected components of $V \setminus \bar{V}_0$. Put $q = \frac{dd^c \sqrt{\sigma}}{\sqrt{\sigma}}$. Let $\{a_1, \dots, a_g\}$ be generic divisor with support in $Y \setminus \bar{X}$, $\bar{X} \subset Y \subset \bar{Y} \subset V$. Let for generic $\theta \in \mathbb{C}$ and $\lambda \in \mathbb{C} : |\lambda| \geq \text{const}(V, \{a_j\}, \theta, \sigma)$ function $z \mapsto \mu = \mu_\theta(z, \lambda)$ be such that:*

$$\begin{aligned} \mu|_Y &\in L^{\tilde{p}}(Y), \quad \mu|_{V \setminus Y} \in L^\infty(V \setminus \bar{Y}), \\ \bar{\partial} \mu|_Y &\in L^p(Y), \quad \bar{\partial} \mu|_{V \setminus \bar{Y}} \in L^{\tilde{p}}(V \setminus Y), \quad 1 \leq p < 2, \quad \tilde{p} > 2, \end{aligned} \tag{2.1}$$

$$\bar{\partial}(\partial + \lambda(dz_1 + \theta dz_2))\mu = \frac{i}{2}q\mu + i \sum_{j=1}^g C_j \delta(z, a_j) \quad \text{with some } C_j = C_{j,\theta}(\lambda) \quad \text{and} \quad (2.2)$$

$$\mu_\theta(z, \lambda) \rightarrow 0, \quad z \rightarrow \infty, \quad z \in V_1. \quad (2.3)$$

Then $\mu_\theta(z, \lambda) \equiv 0, z \in V$.

Remark 2.1. Proposition 2.1 is a corrected version of Proposition 2.1 of [HM2]. For the case $V = \mathbb{C}$ the equivalent result goes back to [BC2].

Lemma 2.1. Let $\psi = \sqrt{\sigma}F = e^{\lambda(z_1 + \theta z_2)}\mu$, where μ satisfies (2.1), (2.2) and

$$F_1 = \sqrt{\sigma}\partial F, \quad F_2 = \sqrt{\sigma}\bar{\partial}F. \quad (2.4)$$

Then forms F_1, F_2 satisfy the system of equations

$$\begin{aligned} \bar{\partial}F_1 + F_2 \wedge \partial \ln \sqrt{\sigma} &= ie^{\lambda(z_1 + \theta z_2)} \sum_{j=1}^g C_j \delta(z, a_j), \\ \partial F_2 + F_1 \wedge \bar{\partial} \ln \sqrt{\sigma} &= -ie^{\lambda(z_1 + \theta z_2)} \sum_{j=1}^g C_j \delta(z, a_j). \end{aligned} \quad (2.5)$$

Proof of Lemma 2.1. From definition of F_1 and F_2 it follows that

$$\begin{aligned} d\sigma d^c F &= i[2\sigma\partial\bar{\partial}F - \bar{\partial}\sigma \wedge \partial F + \partial\sigma \wedge \bar{\partial}F] = \\ &= 2i\sqrt{\sigma}(\partial F_2 + F_1 \wedge \bar{\partial} \ln \sqrt{\sigma}) = -2i\sqrt{\sigma}(\bar{\partial}F_1 + F_2 \wedge \partial \ln \sqrt{\sigma}). \end{aligned}$$

From (2.4) and (2.2) we deduce also that

$$d(\sigma d^c F) = \sqrt{\sigma}(dd^c\psi - \psi \frac{dd^c\sqrt{\sigma}}{\sqrt{\sigma}}) = 2\sqrt{\sigma}e^{\lambda(z_1 + \theta z_2)} \sum_{j=1}^g C_j \delta(z, a_j).$$

These equalities imply (2.5).

Lemma 2.1 is proved.

Lemma 2.2. Let $\{b_m\}$ be the points of X , where $(dz_1 + \theta dz_2)|_X(b_m) = 0$.

Let $B^0 = \cup_m \{b_m\}$ and $A^0 = \cup_j \{a_j\}$.

Let $u_\pm = m_1 \pm e_{-\lambda, \theta}(z)\bar{m}_2$, where $m_1 = e^{-\lambda(z_1 + \theta z_2)}f_1$, $m_2 = e^{-\lambda(z_1 + \theta z_2)}f_2$, $f_1 = \sqrt{\sigma}\frac{\partial F}{\partial z_1}$, $f_2 = \sqrt{\sigma}\frac{\partial F}{\partial \bar{z}_1}$. Let also $q_1 = \frac{\partial \ln \sqrt{\sigma}}{\partial z_1}$ and $\delta_0(z, a_j) = \frac{\delta(z, a_j)}{dz_1 \wedge d\bar{z}_1}$. Then in conditions of Lemma 2.1

$$\begin{aligned} \sup_{z \in X} |\bar{\partial}u_\pm|_X(z) \cdot \text{dist}^2(z, B^0) &= O\left(\sup_{z \in X} |u_\pm \text{dist}(z, B^0)|\right) < \infty; \\ u_\pm|_{V \setminus X} &\in L^1(V \setminus X) \cap O(V \setminus (X \cup A^0)) \end{aligned} \quad (2.6)$$

and system (2.5) is equivalent to the system

$$\begin{aligned} \frac{\partial u_{\pm}}{\partial \bar{z}_1} d\bar{z}_1 &= \mp(e_{-\lambda, \theta}(z)q_1 \bar{u}_{\pm})d\bar{z}_1 + \\ &i \sum_{j=1}^g (C_j \pm \bar{C}_j e_{-\lambda, \theta}(z)) \delta_0(z, a_j) d\bar{z}_1. \end{aligned} \quad (2.7)$$

Proof of Lemma 2.2. From (2.1) we deduce the property

$$\begin{aligned} u_{\pm}|_Y &\in L^p(Y), \quad 1 \leq p < 2, \\ u_{\pm}|_{V \setminus Y} &\in L^{\tilde{p}}(V \setminus Y) \oplus L^{\infty}(V \setminus Y), \quad \tilde{p} > 2. \end{aligned}$$

System (2.5) is equivalent to the system of equations

$$\begin{aligned} \frac{\partial f_1}{\partial \bar{z}_1} &= -f_2 q_1 + i e^{\lambda(z_1 + \theta z_2)} \sum_{j=1}^g C_j \delta_0(z, a_j), \\ \frac{\partial f_2}{\partial z_1} &= -f_1 \bar{q}_1 + i e^{\lambda(z_1 + \theta z_2)} \sum_{j=1}^g C_j \delta_0(z, a_j). \end{aligned}$$

This system and definition of m_1, m_2 imply

$$\begin{aligned} \frac{\partial m_1}{\partial \bar{z}_1} &= -q_1 m_2 + i \sum_{j=1}^g C_j \delta_0(z, a_j), \\ \frac{\partial m_2}{\partial z_1} + \lambda m_2 (1 + \theta \frac{\partial z_2}{\partial z_1}) &= -\bar{q}_1 m_1 + i \sum_{j=1}^g C_j \delta_0(z, a_j). \end{aligned}$$

From the last equalities and definition of u_{\pm} we deduce

$$\begin{aligned} \frac{\partial u_{\pm}}{\partial \bar{z}_1} &= \frac{\partial m_1}{\partial \bar{z}_1} \pm e_{-\lambda, \theta}(z) \left(\frac{\partial \bar{m}_2}{\partial \bar{z}_1} + \bar{\lambda} (1 + \bar{\theta} \frac{\partial \bar{z}_2}{\partial \bar{z}_1}) \bar{m}_2 \right) = -q_1 m_2 + i \sum_{j=1}^g C_j \delta_0(z, a_j) \pm \\ &e_{-\lambda, \theta}(z) \left(\bar{\lambda} (1 + \bar{\theta} \frac{\partial \bar{z}_2}{\partial \bar{z}_1}) \bar{m}_2 - \bar{\lambda} \bar{m}_2 (1 + \bar{\theta} \frac{\partial \bar{z}_2}{\partial \bar{z}_1}) - q_1 \bar{m}_1 + i \sum_{j=1}^g \bar{C}_j \delta_0(z, a_j) \right) = \\ &\mp (e_{-\lambda, \theta}(z) q_1 \bar{u}_{\pm}) + i \sum_{j=1}^g (C_j \pm \bar{C}_j e_{-\lambda, \theta}(z)) \delta_0(z, a_j). \end{aligned}$$

Property (2.7) is proved.

For proving (2.6) we will use construction coming back to Bers and Vekua (see [Ro], [V]). Let β_{\pm} be continuous on Y solutions of $\bar{\partial}$ - equations

$$\bar{\partial} \beta_{\pm} = \pm e_{-\lambda, \theta}(z) q_1 \frac{\bar{u}_{\pm}}{u_{\pm}} d\bar{z}_1,$$

where the right-hand side belongs to $L_{0,1}^\infty(Y)$.

Functions $v_\pm = u_\pm e^{-\beta_\pm}$ belongs to $\mathcal{O}(Y)$. Indeed, from (2.1), (2.2) it follows that $\mu \in W^{1,p}(Y) \cap W_{loc}^{1,\tilde{p}}(Y \setminus (A^0 \cup B^0))$. From this and from definition of v_\pm we deduce that $\bar{\partial}v_\pm = q_1 \bar{u}_\pm d\bar{z}_1 e^{-\beta_\pm} - q_1 u_\pm \frac{\bar{u}_\pm}{u_\pm} e^{-\beta_\pm} d\bar{z}_1 = 0$ on $Y \setminus (A^0 \cup B^0)$ and the following formula for u_\pm is valid

$$u_\pm(z) = v_\pm(z) e^{\beta_\pm(z)}. \quad (2.8)$$

From this and (2.7), (2.8) we obtain (2.6).

Lemma 2.2 is proved.

Lemma 2.3. *Let u_\pm be the functions from Lemma 2.2 and μ be the function from Lemma 2.1. Then*

$$u_\pm = \frac{\partial \mu}{\partial z_1} + \lambda(1 + \theta \frac{\partial z_2}{\partial z_1}) \mu - q_1 \mu \pm e_{-\lambda, \theta}(z) \left(\frac{\partial \bar{\mu}}{\partial z_1} - q_1 \bar{\mu} \right).$$

Proof of Lemma 2.3. We have

$$u_\pm = e^{-\lambda(z_1 + \theta z_2)} f_1 \pm e^{-\lambda(z_1 + \theta z_2)} \bar{f}_2 = e^{-\lambda(z_1 + \theta z_2)} (f_1 \pm \bar{f}_2),$$

where

$$\begin{aligned} f_1 &= \sqrt{\sigma} \frac{\partial F}{\partial z_1} = \sqrt{\sigma} \frac{\partial}{\partial z_1} \left(\frac{1}{\sqrt{\sigma}} e^{\lambda(z_1 + \theta z_2)} \mu \right) = \\ &e^{\lambda(z_1 + \theta z_2)} \left(\frac{\partial \mu}{\partial z_1} + \lambda(1 + \theta \frac{\partial z_2}{\partial z_1}) \mu - q_1 \mu \right), \\ \bar{f}_2 &= \sqrt{\sigma} \frac{\partial \bar{F}}{\partial z_1} = \sqrt{\sigma} \frac{\partial}{\partial z_1} \left(\frac{1}{\sqrt{\sigma}} e^{\bar{\lambda}(\bar{z}_1 + \bar{\theta} \bar{z}_2)} \bar{\mu} \right) = \\ &e^{\bar{\lambda}(\bar{z}_1 + \bar{\theta} \bar{z}_2)} \left(\frac{\partial \bar{\mu}}{\partial z_1} - q_1 \bar{\mu} \right). \end{aligned}$$

This imply Lemma 2.3.

Lemma 2.4. *Let $\omega_1, \dots, \omega_g$ be orthonormal basis of holomorphic 1-forms on \tilde{V} . Let $\{a_1, \dots, a_g\}$ be generic divisor on $Y \setminus \bar{X}$, where $V_0 \subset \bar{X} \subset Y \subset V$. Put $\omega_{j,k}^0 = \frac{\omega_k}{dz_1}(a_j)$. Let for some generic $\theta \in \mathbb{C}$ and $\lambda \in \mathbb{C}$ functions u_\pm from Lemmas 2.2-2.3 satisfy (2.6), (2.7) with some $C_j = C_{j,\theta}(\lambda)$. Then*

$$\sup_j |C_{j,\theta}(\lambda)| \leq \text{const}(V, \{a_j\}, \theta) \|\ln \sqrt{\sigma}\|_{W^{2,\infty}(X)}^2 (1 + |\lambda|)^{-1/3} \|u_\pm\|_{L^\infty(X, B^0)},$$

$$\text{where } \|u_\pm\|_{L^\infty(X, B^0)} \stackrel{\text{def}}{=} \sup_{z \in X} |u_\pm(z) \text{dist}(z, B^0)|.$$

Proof of Lemma 2.4. From condition iv) of section 1 we deduce $|\omega_{j,k}^0| < \infty$. From definition of generic divisor we obtain $\det[\omega_{j,k}^0] \neq 0$. From (2.7) and from definition of

Dirac measure $\forall k = 1, \dots, g$ we deduce

$$\begin{aligned}
& \overline{\lim}_{r \rightarrow \infty} \left(\int_{\{z \in V: |z_1|=r\}} u_{\pm} \wedge \omega_k \right) \pm \int_X e_{-\lambda, \theta}(z) \frac{\partial \ln \sqrt{\sigma}}{\partial z_1} \bar{u}_{\pm} d\bar{z}_1 \wedge \omega_k = \\
& i \int_Y \sum_{j=1}^g (C_j \pm \bar{C}_j e_{-\lambda, \theta}(z)) \delta_0(z, a_j) d\bar{z}_1 \wedge \omega_k = \\
& i \sum_{j=1}^g (C_j \pm \bar{C}_j e_{-\lambda, \theta}(a_j)) \omega_{j,k}^0, \quad j, k = 1, 2, \dots, g.
\end{aligned} \tag{2.9}$$

From estimates $\overline{\lim}_{r_n \rightarrow \infty} \sup_{\{z \in V: |z_1|=r_n\}} |u_{\pm}(z)| < \infty$, for some sequence $r_n \rightarrow \infty$, and $\frac{|\omega_k|}{dz_1} \leq O(|\frac{1}{z_1^2}|)$, $z \in V \setminus Y$, $k = 1, \dots, g$, we obtain

$$\overline{\lim}_{r \rightarrow \infty} \left| \int_{\{z \in V: |z_1|=r\}} u_{\pm} \wedge \omega_k \right| = 0. \tag{2.10}$$

From (2.9), (2.10) and Kramers's formula we obtain

$$\begin{aligned}
& i(C_j \pm \bar{C}_j e_{-\lambda, \theta}(a_j)) = \\
& \frac{\det[\omega_{1,k}^0; \dots; \omega_{j-1,k}^0; \int_X \pm e_{-\lambda, \theta}(z) \frac{\partial \ln \sqrt{\sigma}}{\partial z_1} \bar{u}_{\pm} d\bar{z}_1 \wedge \omega_k; \omega_{j+1,k}^0; \dots; \omega_{g,k}^0]}{\det[\omega_{j,k}^0]},
\end{aligned} \tag{2.11}$$

where $j, k = 1, \dots, g$.

Let us prove estimate

$$\begin{aligned}
& \left| \int_X e_{-\lambda, \theta}(z) \frac{\partial \ln \sqrt{\sigma}}{\partial z_1} \bar{u}_{\pm} d\bar{z}_1 \wedge \omega_k \right| \leq \\
& \text{const}(X, \theta) (1 + |\lambda|)^{-1/3} \|\ln \sqrt{\sigma}\|_{W^{2, \infty}(X)}^2 \cdot \|u_{\pm}\|_{L^{\infty}(X, B^0)}.
\end{aligned} \tag{2.12}$$

For $|\lambda| \leq 1$ estimate follows directly, using that $\ln \sqrt{\sigma} \in W^{1, \infty}(X)$.

Let $B^{\varepsilon} = \cup_{m=1}^M \{z \in X : |z - b_m| \leq \varepsilon\}$.

Let $\chi_{\varepsilon, \nu}$, $\nu = 1, 2$, be functions from $C^{(1)}(V)$ such that $\chi_{\varepsilon, 1} + \chi_{\varepsilon, 2} \equiv 1$ on V , $\text{supp } \chi_{\varepsilon, 1} \subset B^{2\varepsilon}$, $\text{supp } \chi_{\varepsilon, 2} \subset V \setminus B^{\varepsilon}$, $|d\chi_{\varepsilon, \nu}| = O(\frac{1}{\varepsilon})$, $\nu = 1, 2$.

Put $J_{\nu}^{\varepsilon} u_{\pm} = \int_X \chi_{\varepsilon, \nu}(z) e_{-\lambda, \theta}(z) \frac{\partial \ln \sqrt{\sigma}}{\partial z_1} \bar{u}_{\pm} d\bar{z}_1 \wedge \omega_k$, $\nu = 1, 2$. We have directly:

$$|J_1^{\varepsilon} u_{\pm}| \leq \text{const}(X) \varepsilon \|\ln \sqrt{\sigma}\|_{W^{1, 1}(X)} \cdot \|u_{\pm}\|_{L^{\infty}(X, B^0)}. \tag{2.13}$$

For $J_2^\varepsilon u_\pm$ we obtain by integration by parts:

$$\begin{aligned} J_2^\varepsilon u_\pm &= -\frac{1}{\lambda} \int_X \chi_{\varepsilon,2} \partial e_{-\lambda,\theta}(z) \frac{\partial \ln \sqrt{\sigma}}{\partial z_1} \bar{u}_\pm d\bar{z}_1 \wedge \frac{\omega_k}{dz_1 + \theta dz_2} = \\ &\frac{1}{\lambda} \int_X e_{-\lambda,\theta}(z) \partial \left(\chi_{\varepsilon,2} \frac{\partial \ln \sqrt{\sigma}}{\partial z_1} \bar{u}_\pm d\bar{z}_1 \wedge \frac{\omega_k}{dz_1 + \theta dz_2} \right). \end{aligned} \quad (2.14)$$

To estimate (2.14) we use (2.6) and the following properties: $|\partial \chi_{\varepsilon,2}| = O(\frac{1}{\varepsilon})$, $\text{supp}(\partial \chi_{\varepsilon,2}) \subset B^{2\varepsilon}$,

$$\begin{aligned} &\left\| \frac{\partial \ln \sqrt{\sigma}}{\partial z_1} d\bar{z}_1 \wedge \partial \chi_{\varepsilon,2} u_\pm \frac{\omega_k}{dz_1 + \theta dz_2} \right\|_{L^1_{0,1}(X)} \leq \\ &\frac{\text{const}(X, \theta)}{\varepsilon} \|\ln \sqrt{\sigma}\|_{W^{1,\infty}(X)} \|u_\pm\|_{L^\infty(X, B^0)} \\ &\left\| \frac{\partial^2 \ln \sqrt{\sigma}}{\partial z_1^2} dz_1 \wedge d\bar{z}_1 \chi_{\varepsilon,2} u_\pm \frac{\omega_k}{dz_1 + \theta dz_2} \right\|_{L^1_{1,1}(X)} \leq \\ &|\ln \varepsilon| \text{const}(X, \theta) \|\ln \sqrt{\sigma}\|_{W^{2,\infty}(X)} \|u_\pm\|_{L^\infty(X, B^0)} \\ &\left\| \frac{\partial \ln \sqrt{\sigma}}{\partial z_1} d\bar{z}_1 \chi_{\varepsilon,2} u_\pm \wedge \partial \left(\frac{\omega_k}{dz_1 + \theta dz_2} \right) \right\|_{L^1_{0,1}(X)} \leq \\ &\frac{\text{const}(X, \theta)}{\varepsilon} \|\ln \sqrt{\sigma}\|_{W^{1,\infty}(X)} \|u_\pm\|_{L^\infty(X, B^0)} \\ &\partial \bar{u}_\pm \Big|_X = \mp(e_{\lambda,\theta}(z) \bar{q}_1 \bar{u}_\pm) dz_1. \end{aligned}$$

From (2.14), (2.6) and these properties we obtain

$$\begin{aligned} |J_2^\varepsilon u_\pm| &\leq |\ln \varepsilon| \frac{\text{const}(X, \theta)}{|\lambda|} \|\ln \sqrt{\sigma}\|_{W^{2,\infty}(X)} \cdot \|u_\pm\|_{L^\infty(X, B^0)} + \\ &\frac{\text{const}(X, \theta)}{\varepsilon |\lambda|} \|\ln \sqrt{\sigma}\|_{W^{1,\infty}(X)} \cdot \|u_\pm\|_{L^\infty(X, B^0)} + \\ &\frac{\text{const}(X, \theta, \delta)}{\varepsilon^{1+\delta} |\lambda|} \|\ln \sqrt{\sigma}\|_{W^{1,\infty}(X)} \cdot \|u_\pm\|_{L^\infty(X, B^0)}. \end{aligned} \quad (2.15)$$

Putting in (2.13), (2.15) $\varepsilon = \frac{1}{\sqrt{\lambda}}$ and $\delta = 1/3$ we obtain (2.12) for $|\lambda| \geq 1$.

Inequalities (2.11), (2.12) imply estimate

$$|C_j \pm \bar{C}_j e_{-\lambda,\theta}(a_j)| \leq \text{const}(X, \{a_j\}, \theta) (1 + |\lambda|)^{-1/3} \|\ln \sqrt{\sigma}\|_{W^{2,\infty}(X)}^2 \cdot \|u_\pm\|_{L^\infty(X, B^0)}.$$

We obtained statement of Lemma 2.4.

Lemma 2.5. *Let functions u_\pm satisfy (2.6), (2.7) and R - operator from section 1. Then*

$$\|R[e_{-\lambda,\theta} q_1 \bar{u}_\pm d\bar{\xi}_1]\|_{L^\infty(X, B^0)} \leq \text{const}(X, \theta) (1 + |\lambda|)^{-1/5} \|\ln \sqrt{\sigma}\|_{W^{2,\infty}(X)} \cdot \|u_\pm\|_{L^\infty(X, B^0)}.$$

Proof of Lemma 2.5.

Let $\chi_{\varepsilon, \nu}$, $\nu = 1, 2$, be partition of unity from Lemma 2.4. Put $S_\nu^\varepsilon u_\pm = R[\chi_{\varepsilon, \nu} q_1 \bar{u}_\pm d\bar{\xi}_1]$, $\nu = 1, 2$. Using (2.6) and formula for operator R we deduce estimate

$$\|S_1^\varepsilon u_\pm\|_{L^\infty(X, B^0)} = O(\varepsilon) \|\ln \sqrt{\sigma}\|_{W^{1, \infty}(X)} \|u_\pm\|_{L^\infty(X, B^0)}. \quad (2.16)$$

Let $R_{1,0}(\xi, z)$ be kernel of operator R . It means, in particular, that $\bar{\partial}_\xi R_{1,0}(\xi, z) = -\delta(\xi, z)$, where $\delta(\xi, z)$ - Dirac (1,1)- measure, concentrated in the point $\xi = z$. We have

$$S_2^\varepsilon u_\pm = \int_X \chi_{\varepsilon, 2} e_{-\lambda, \theta} q_1 \bar{u}_\pm d\bar{\xi}_1 R_{1,0}(\xi, z). \quad (2.17)$$

Integration by parts in (2.17) gives the following

$$\begin{aligned} S_2^\varepsilon u_\pm &= \frac{1}{\lambda} \int_X \bar{\partial} e_{-\lambda, \theta}(\xi) \frac{d\bar{\xi}_1}{d\bar{\xi}_1 + \theta d\bar{\xi}_2} \chi_{\varepsilon, 2}(\xi) q_1(\xi) \bar{u}_\pm(\xi) R_{1,0}(\xi, z) = \\ &- \frac{1}{\lambda} \int_X e_{-\lambda, \theta}(\xi) \bar{\partial} \left(\frac{d\bar{\xi}_1}{d\bar{\xi}_1 + \theta d\bar{\xi}_2} \chi_{\varepsilon, 2}(\xi) q_1(\xi) \bar{u}_\pm(\xi) \right) R_{1,0}(\xi, z) + \\ &\frac{1}{\lambda} e_{-\lambda, \theta}(z) \frac{d\bar{\xi}_1}{d\bar{\xi}_1 + \theta d\bar{\xi}_2}(z) \chi_{\varepsilon, 2}(z) q_1(z) \bar{u}_\pm(z). \end{aligned} \quad (2.18)$$

To estimate (2.18) we use (2.6), properties of partition of unity $\{\chi_{\varepsilon, \nu}\}$ and inequalities

$$\begin{aligned} \left| \frac{d\bar{\xi}_1}{d\bar{\xi}_1 + \theta d\bar{\xi}_2}(\xi) \right| &= O\left(\frac{1}{\text{dist}(\xi, B^0)}\right), \quad \left| \bar{\partial} \frac{d\bar{\xi}_1}{d\bar{\xi}_1 + \theta d\bar{\xi}_2}(\xi) \right| = O\left(\frac{1}{(\text{dist}(\xi, B^0))^2}\right), \\ |q_1(\xi)| &= O\left(\frac{1}{\text{dist}(\xi, B^0)}\right), \quad |\bar{\partial} q_1(\xi)| = O\left(\frac{1}{(\text{dist}(\xi, B^0))^2}\right), \quad \xi \in X. \end{aligned} \quad (2.19)$$

From (2.19), (2.8) and from the formula for operator R we deduce estimate

$$\|S_2^\varepsilon u_\pm\|_{L^\infty(X)} = O\left(\frac{1}{\varepsilon^4 |\lambda|}\right) \|\ln \sqrt{\sigma}\|_{W^{2, \infty}(X)} \|u_\pm\|_{L^\infty(X, B^0)}. \quad (2.20)$$

Putting in (2.16), (2.20) $\varepsilon = \frac{1}{|\lambda|^{1/5}}$ we obtain statement of Lemma 2.5.

Proof of Proposition 2.1.

Let function μ satisfy conditions (2.1)-(2.3) and u_\pm be functions defined in Lemma 2.2. Then by Lemma 2.3 we have

$$\begin{aligned} \lim_{\substack{z \rightarrow \infty \\ z \in V_1}} u_\pm(z, \lambda) &= \lim_{\substack{z \rightarrow \infty \\ z \in V_1}} (m_1 \pm e_{-\lambda, \theta}(z) \bar{m}_2) = \\ \lim_{\substack{z \rightarrow \infty \\ z \in V_1}} \left[\lambda \left(1 + \theta \frac{dz_2}{dz_1}\right) \mu + \frac{\partial \mu}{\partial z_1} \pm e_{-\lambda, \theta}(z) \frac{\partial \bar{\mu}}{\partial z_1} \right] &\rightarrow 0. \end{aligned} \quad (2.21)$$

Let

$$h_{\pm} = u_{\pm} \pm R[(e_{-\lambda, \theta}(z)q_1 \bar{u}_{\pm})d\bar{z}_1 - i \sum_{j=1}^g (C_j \pm \bar{C}_j e_{-\lambda, \theta}(z))\delta_0(z, a_j)d\bar{z}_1], \quad (2.22)$$

where R is the operator from section 1.

By Lemmas 2.2-2.5 and properties of operator R we have $h_{\pm} \in \mathcal{O}(V) \cap L^{\infty}(V)$ and $h_{\pm}(z, \lambda) \rightarrow 0, z \rightarrow \infty, z \in V_1$. By Liouville theorem, $h_{\pm}(z, \lambda) \equiv 0$ on $V, \lambda \in \mathbb{C}$. Then from (2.22) with $h_{\pm}(z, \lambda) \equiv 0$ and Lemmas 2.4, 2.5 it follows that $u_{\pm}(z, \lambda) \equiv 0, z \in V$, if $\lambda \in \mathbb{C} \setminus E_{\theta} : |\lambda| \geq \text{const}(V, \{a_j\}, \theta) \|\ln \sqrt{\sigma}\|_{W^{2, \infty}(X)}^2$. Property $u_{\pm}(z, \lambda) \equiv 0, z \in V$, implies by Lemma 2.3 equality $\frac{\partial \mu}{\partial \bar{z}_1} - \bar{q}_1 \mu = 0, z \in V$, where $\mu(z) \rightarrow \infty$ if $z \in V_1, z \rightarrow \infty$. The Liouville type theorem for generalized holomorphic functions ([Ro], theorem 7.1) implies $\mu \equiv 0$. Proposition 2.1 is proved.

3. Faddeev type functions on Riemann surface. Existence.

Proof of Theorem 1.1A

Proposition 3.1. *Let conductivity σ and divisor $\{a_1, \dots, a_g\}$ satisfy conditions of Proposition 2.1. Then \forall generic $\theta \in \mathbb{C}$ and $\forall \lambda \in \mathbb{C} \setminus E_{\theta} : |\lambda| \geq \text{const}(V, \{a_j\}, \theta, \sigma)$ there exists unique Faddeev type function*

$$\begin{aligned} \psi &\stackrel{\text{def}}{=} \sqrt{\sigma} F \stackrel{\text{def}}{=} e^{\lambda(z_1 + \theta z_2)} \mu, \quad \text{where} \\ \psi &= \psi_{\theta}(z, \lambda), \quad F = F_{\theta}(z, \lambda), \quad \mu = \mu_{\theta}(z, \lambda), \end{aligned} \quad (3.1)$$

associated with σ and divisor $\{a_1, \dots, a_g\}$, i.e.

$$\begin{aligned} \bar{\partial}(\partial + \lambda(dz_1 + \theta dz_2))\mu &= \frac{i}{2}q\mu + \sum_{j=1}^g C_j \delta(z, a_j), \quad \text{for some } C_j = C_{j, \theta}(\lambda), \quad \text{where} \\ q &= \frac{dd^c \sqrt{\sigma}}{\sqrt{\sigma}}, \quad \mu|_Y \in L^{\tilde{p}}(Y), \quad \mu|_{V \setminus \bar{Y}} \in L^{\infty}(V \setminus \bar{Y}), \quad \lim_{\substack{z \rightarrow \infty \\ z \in V_1}} \mu_{\theta}(z, \lambda) = 1. \end{aligned} \quad (3.1a)$$

In addition,

$$\begin{aligned} \|\mu_{\theta}(z, \lambda) - \mu_{\theta}(\infty_l, \lambda)\|_{L^{\tilde{p}}(V)} &\leq \frac{\text{const}(V, \{a_j\}, \theta, \sigma, \tilde{p}, \varepsilon)}{|\Delta_{\theta}(\lambda)| \cdot (1 + |\lambda|)^{g+1-\varepsilon}}, \\ \text{where } \mu_{\theta}(\infty_l, \lambda) &\stackrel{\text{def}}{=} \lim_{\substack{z \rightarrow \infty \\ z \in V_l}} \mu_{\theta}(z, \lambda), \quad l = 1, \dots, d, \end{aligned} \quad (3.1b)$$

$$\|\partial \mu\|_{L^p_{1,0}(Y)} + \|\partial \mu\|_{L^{\tilde{p}}_{1,0}(V \setminus Y)} \leq \frac{\text{const}(V, \{a_j\}, \theta, \sigma, p, \tilde{p}, \varepsilon)}{|\Delta_{\theta}(\lambda)| \cdot (1 + |\lambda|)^{g-\varepsilon}}, \quad p < 2, \quad \tilde{p} > 2,$$

$$\begin{aligned} \forall \text{ generic } \theta \in \mathbb{C} \text{ and } \lambda \in \mathbb{C} \setminus E_{\theta} : |\lambda| \geq \text{const}(V, \{a_j\}, \theta, \sigma), \\ \frac{\partial \mu}{\partial \lambda}|_Y \in W^{1,p}(Y), \quad \frac{\partial \mu}{\partial \lambda}|_{V_l \setminus Y} \in L^{\infty}(V_l \setminus Y) \cup W^{1, \tilde{p}}(V_l \setminus Y), \end{aligned} \quad (3.1c)$$

where $\{V_l\}$ are connected components of $V \setminus V_0$, $l = 1, \dots, d$,

$$e_{\lambda, \theta}(z) = e^{\lambda(z_1 + \theta z_2) - \bar{\lambda}(\bar{z}_1 + \bar{\theta} \bar{z}_2)}.$$

Remark 3.1. Proposition 3.1 is a corrected version of Proposition 2.2 from [HM2]. For the case $V = \mathbb{C}$ the results of such a type goes back to [F1], [F2].

Lemma 3.1. Under the conditions of Proposition 3.1, $\forall \lambda \in \mathbb{C} \setminus E_\theta$ function $z \rightarrow \mu_\theta(z, \lambda)$ belonging to $L^{\bar{p}}(Y)$ on Y and to $L^\infty(V \setminus Y)$ on $V \setminus Y$ satisfies (3.1a) iff there exists $C_j = C_{j, \theta}(\lambda)$, $j = 1, \dots, g$, such that

$$\mu_\theta(z, \lambda) = 1 + \frac{i}{2} \int_{\xi \in X} g_{\lambda, \theta}(z, \xi) q \mu_\theta(\xi, \lambda) + i \sum_{j=1}^g C_{j, \theta}(\lambda) g_{\lambda, \theta}(z, a_j) \quad (3.2)$$

and one of two equivalent conditions is valid

$$\mathcal{H}_{\lambda, \theta}(\hat{R}_\theta(\frac{i}{2} q \mu)) + i \sum_{j=1}^g C_{j, \theta}(\lambda) \mathcal{H}_{\lambda, \theta}(\hat{R}_\theta(\delta(z, a_j))) = 0 \quad \text{or} \quad (3.3)$$

$$(\partial + \lambda(dz_1 + \theta dz_2)) \mu_\theta(z, \lambda) \in H_{1,0}(V \setminus (X \cup_{j=1}^g \{a_j\})) \cap L_{1,0}^1(Y \setminus X),$$

where $g_{\lambda, \theta}$ is Faddeev type Green function, \hat{R}_θ , $\mathcal{H}_{\lambda, \theta}$ - operators defined in section 1.

Proof of Lemma 3.1. From Proposition 4 in [He] and from definition of Green function $g_{\lambda, \theta}(z, \xi)$ we deduce that integral equation (3.2) is equivalent to the following differential equation

$$\begin{aligned} \bar{\partial}(\partial + \lambda(dz_1 + \theta dz_2)) \mu &= \frac{i}{2} q \mu + i \sum_{j=1}^g C_{j, \theta} \delta(z, a_j) + \\ \bar{\lambda}(d\bar{z}_1 + \bar{\theta} d\bar{z}_2) &\times [\mathcal{H}_{\lambda, \theta}(\hat{R}_\theta(\frac{i}{2} q \mu)) + i \sum_{j=1}^g C_{j, \theta} \mathcal{H}_{\lambda, \theta}(\hat{R}_\theta(\delta(z, a_j)))]. \end{aligned} \quad (3.4)$$

Equation (3.4) is equivalent to (3.1a) if one of two equivalent conditions (3.3) is valid.

Lemma 3.1 is proved.

Lemma 3.2. Let $\{a_1, \dots, a_g\}$ be generic divisor in $Y \setminus \bar{X}$. Then for any generic $\theta \in \mathbb{C}$ and

$\forall \lambda \in \mathbb{C} \setminus E_\theta : |\lambda| \geq \text{const}(V, \{a_j\}, \theta, \sigma)$, integral equation (3.2), (3.3) is uniquely solvable Fredholm integral equation in the space $\tilde{W}^{1, \bar{p}}(V)$.

Proof of Lemma 3.2. Let $\theta \in \mathbb{C}$ and $\lambda \in \mathbb{C} \setminus E_\theta : |\lambda| \geq \text{const}(V, \{a_j\}, \theta, \sigma)$. From (3.2), (3.3) we obtain integral equation for $\tilde{\mu}_\theta = \mu_\theta - 1$ and $\tilde{C}_{j, \theta}$:

$$\begin{aligned} \tilde{\mu}_\theta(z, \lambda) - \frac{i}{2} \int_{\xi \in V} g_{\lambda, \theta}(z, \xi) q(\xi) \tilde{\mu}_\theta(\xi, \lambda) - i \sum_{j=1}^g \tilde{C}_{j, \theta}(\lambda) g_{\lambda, \theta}(z, a_j) = \\ \frac{i}{2} \int_{\xi \in V} g_{\lambda, \theta}(z, \xi) q(\xi) + i \sum_{j=1}^g C_{j, \theta}^0(\lambda) g_{\lambda, \theta}(z, a_j). \end{aligned} \quad (3.5)$$

Parameters $\tilde{C}_j = \tilde{C}_{j,\theta}(\lambda)$, $j = 1, \dots, g$, are defined by the equations:

$$\begin{aligned} -i \sum_{j=1}^g \tilde{C}_j \int_V \hat{R}_\theta(\delta(\xi, a_j)) \bar{\omega}_k(\xi) e_{\lambda,\theta}(\xi) &= \\ \int_{\xi \in V} e_{\lambda,\theta}(\xi) \hat{R}_\theta\left(\frac{i}{2} q \tilde{\mu}\right) \bar{\omega}_k(\xi), \quad k = 1, 2, \dots, g. \end{aligned} \quad (3.6)$$

We remind that determinant of system (3.6) is exactly $\Delta_\theta(\lambda)$.

Parameters $C_{j,\theta}^0$ are defined by (3.6) with $C_{j,\theta}^0$ in place of $\tilde{C}_{j,\theta}$ and 1 in place of $\tilde{\mu}$. One can see also that $C_{j,\theta}^0(\lambda) = C_{j,\theta}(\lambda) - \tilde{C}_{j,\theta}(\lambda)$.

Let us prove that (3.5), (3.6) determine Fredholm integral equation in the space $\tilde{W}^{1,\tilde{p}}(V)$, $\tilde{p} > 2$.

Propositions 2, 3 of [He] imply that correspondance

$$\tilde{\mu} \mapsto R_{\lambda,\theta} \circ \left(\hat{R}_\theta\left(\frac{i}{2} q \tilde{\mu}\right) + i \sum_{j=1}^g \tilde{C}_{j,\theta} \hat{R}_\theta(\delta(z, a_j)) \right)$$

define linear continuous mapping of $\tilde{W}^{1,\tilde{p}}(V)$ into itself. This mapping is compact because mapping $\tilde{\mu} \rightarrow q\tilde{\mu}$, $\text{supp } q \subset X$, from $\tilde{W}^{1,\tilde{p}}(V)$ into $L_{1,1}^{\tilde{p}}(X)$ is compact, operator $\hat{R}_\theta : L_{1,1}^{\tilde{p}}(X) \rightarrow \tilde{W}_{1,0}^{1,\tilde{p}}(V)$ and operator $R_{\lambda,\theta} : \tilde{W}_{1,0}^{1,\tilde{p}}(V) \rightarrow \tilde{W}^{1,\tilde{p}}(V)$ are bounded.

If for fixed $\lambda \notin E_\theta$ Fredholm equation (3.5), (3.6) is not solvable then corresponding homogeneous equation, when the right-hand side of (3.5) is replaced by zero, admits nontrivial solution $\tilde{\mu}^* = \mu^* - 1$.

By Lemma 3.1 function $\tilde{\mu}^*$ satisfies differential equation (2.2) with C_j replaced by \tilde{C}_j and with property $\tilde{\mu}^*(z) \rightarrow 0$, $z \rightarrow \infty$, $z \in V_1$.

By Proposition 2.1, $\tilde{\mu}^* \equiv 0$ if $\lambda \in \mathbb{C} \setminus E_\theta : |\lambda| \geq \text{const}(V, \{a_j\}, \theta, \sigma)$.

It means that equation (3.2), (3.3) is uniquely solvable Fredholm integral equation for any $\lambda \in \mathbb{C} \setminus E_\theta : |\lambda| \geq \text{const}(V, \{a_j\}, \theta, \sigma)$.

Lemma 3.2 is proved.

Lemma 3.3. *Let $\{a_1, \dots, a_g\}$ be generic divisor on $Y \setminus X$. Let $\lambda \in \mathbb{C} \setminus E_\theta$. Let μ be solution of integral equation (3.2), (3.3). Then relations (3.3) determining parameters $C_j = C_{j,\theta}(\lambda)$ are reduced to the following explicit formulas*

$$2i \sum_{j=1}^g C_{j,\theta} e_{\lambda,\theta}(a_j) \frac{\bar{\omega}_k}{d\bar{z}_1}(a_j) = \int_{z \in X} e_{\lambda,\theta}(z) \left(i \frac{dd^c \sqrt{\sigma}}{\sqrt{\sigma}} + 2\bar{\partial} \ln \sqrt{\sigma} \wedge \partial \ln \sqrt{\sigma} \right) \mu \frac{\bar{\omega}_k}{d\bar{z}_1}(z). \quad (3.7)$$

Proof of Lemma 3.3. By Lemma 3.1 equations (3.2), (3.3) are equivalent to the equation:

$$\bar{\partial}(\partial + \lambda(dz_1 + \theta dz_2))\mu = \frac{i}{2} q \mu + i \sum_{j=1}^g C_{j,\theta} \delta(z, a_j), \quad (3.8)$$

where $\mu = \mu_\theta(z, \lambda) \rightarrow 1$, $z \in V_1$, $z \rightarrow \infty$.

System (2.7) implies the following relation

$$\begin{aligned} \overline{\lim}_{R \rightarrow \infty} \int_{|z_1|=R} \bar{u}_\pm \wedge \bar{\omega}_k + i \int_{z \in V \setminus X} \sum_{j=1}^g (\bar{C}_{j,\theta} \mp C_{j,\theta} e_{\lambda,\theta}(z)) \frac{\delta(z, a_j)}{d\bar{z}_1} \bar{\omega}_k = \\ \mp \int_{z \in X} e^{\lambda(z_1 + \theta z_2) - \bar{\lambda}(\bar{z}_1 + \bar{\theta} \bar{z}_2)} \bar{q}_1 u_\pm dz_1 \wedge \bar{\omega}_k, \end{aligned} \quad (3.9)$$

where $\bar{q}_1 = \frac{\partial \ln \sqrt{\sigma}}{\partial \bar{z}_1}$.

To obtain (3.9) we multiply the both sides of (2.7) by $\wedge \omega_k$, integrate on V and take conjugation.

From Lemma 2.3 and Lemma 3.2 it follows that

$$u_\pm(z) \rightarrow \lambda(1 + \theta \gamma_l) \cdot \lim_{\substack{z \rightarrow \infty \\ z \in V_l}} \mu_\theta(z, \lambda), \quad z \rightarrow \infty, \quad z \in V_l,$$

$$\text{where } \gamma_l = \lim_{\substack{z \rightarrow \infty \\ z \in V_l}} \frac{\partial z_2}{\partial z_1}, \quad \lim_{\substack{z \rightarrow \infty \\ z \in V_1}} \mu_\theta(z, \lambda) = 1.$$

Existence of $\lim_{\substack{z \rightarrow \infty \\ z \in V_l}} \mu_\theta(z, \lambda)$ follows from Lemma 4.1 below. This imply that

$$\overline{\lim}_{R \rightarrow \infty} \left| \int_{|z_1|=R} \bar{u}_\pm \wedge \bar{\omega}_k \right| = \overline{\lim}_{R \rightarrow \infty} \left| \int_{|z_1|=R} \bar{\lambda}(1 + \bar{\theta} \bar{\gamma}_l) \bar{\omega}_k \right| = \lim_{R \rightarrow \infty} |\lambda| O\left(\frac{1}{R}\right) = 0. \quad (3.10)$$

From (3.9), (3.10) and definition of u_\pm we obtain

$$\begin{aligned} 2i \sum_{j=1}^g \int_{z \in V \setminus X} C_j e_{\lambda,\theta}(z) \frac{\delta(z, a_j)}{d\bar{z}_1} \wedge \bar{\omega}_k = \int_{z \in X} e_{\lambda,\theta}(z) \bar{q}_1 (u_+ + u_-) dz_1 \wedge \bar{\omega}_k = \\ 2 \int_{z \in X} e^{-\bar{\lambda}(\bar{z}_1 + \bar{\theta} \bar{z}_2)} \bar{q}_1 f_1 dz_1 \wedge \bar{\omega}_k, \quad \text{where } f_1 = \sqrt{\sigma} \frac{\partial F}{\partial z_1}. \end{aligned}$$

By Lemma 2.3 we have

$$\begin{aligned} 2 \int_{z \in X} e^{-\bar{\lambda}(\bar{z}_1 + \bar{\theta} \bar{z}_2)} \bar{q}_1 f_1 dz_1 \wedge \bar{\omega}_k = \\ 2 \int_{z \in X} e_{\lambda,\theta}(z) \bar{q}_1 \left(\frac{\partial \mu}{\partial z_1} + \lambda \mu + \lambda \theta \frac{\partial z_2}{\partial z_1} \mu - q_1 \mu \right) dz_1 \wedge \bar{\omega}_k. \end{aligned} \quad (3.11)$$

From definition of $\delta(z, a_j)$ we have

$$2i \sum_{j=1}^g \int_{z \in V \setminus X} C_j e_{\lambda,\theta}(z) \frac{\delta(z, a_j)}{d\bar{z}_1} \wedge \bar{\omega}_k = -2i \sum_{j=1}^g C_j e_{\lambda,\theta}(a_j) \frac{\bar{\omega}_k}{d\bar{z}_1}(a_j). \quad (3.12)$$

By integration by part we have

$$\begin{aligned}
& 2 \int_{z \in X} e_{\lambda, \theta}(z) \bar{q}_1 \left(\frac{\partial \mu}{\partial z_1} + \lambda \mu \right) dz_1 \wedge \bar{\omega}_k = 2 \int_X e_{\lambda, \theta}(z) \frac{\partial \ln \sqrt{\sigma}}{\partial \bar{z}_1} \lambda \mu dz_1 \wedge \bar{\omega}_k - \\
& - 2 \int_X e_{\lambda, \theta}(z) \frac{\partial \ln \sqrt{\sigma}}{\partial \bar{z}_1} \left(\lambda \mu + \lambda \theta \frac{\partial z_2}{\partial z_1} \mu \right) dz_1 \wedge \bar{\omega}_k - 2 \int_X e_{\lambda, \theta}(z) \frac{\partial^2 \ln \sqrt{\sigma}}{\partial z_1 \partial \bar{z}_1} \mu dz_1 \wedge \bar{\omega}_k = \quad (3.13) \\
& - 2 \int_X e_{\lambda, \theta}(z) \left(\frac{\partial^2 \ln \sqrt{\sigma}}{\partial z_1 \partial \bar{z}_1} + \frac{\partial \ln \sqrt{\sigma}}{\partial \bar{z}_1} \lambda \theta \frac{\partial z_2}{\partial z_1} \right) \mu dz_1 \wedge \bar{\omega}_k.
\end{aligned}$$

Using (3.11), (3.12), (3.13) we obtain

$$i \sum_{j=1}^g C_{j, \theta} e_{\lambda, \theta}(a_j) \frac{\bar{\omega}_k}{dz_1}(a_j) = \int_{z \in X} e_{\lambda, \theta}(z) \left(\frac{\partial^2 \ln \sqrt{\sigma}}{\partial z_1 \partial \bar{z}_1} + \left| \frac{\partial \ln \sqrt{\sigma}}{\partial \bar{z}_1} \right|^2 \right) \mu dz_1 \wedge \bar{\omega}_k.$$

Lemma 3.3 is proved.

Proof of Proposition 3.1. a) By Lemmas 3.1-3.3 statement (3.1a) of Proposition is valid, i.e. there exists function $z \rightarrow \mu_\theta(z, \lambda)$, $z \in V$ with property (3.1a) $\forall \lambda \in \mathbb{C} \setminus E_\theta : |\lambda| \geq \text{const}(V, \{a_j\}, \theta, \sigma)$.

b) Put $f_0 = \hat{R}_\theta(\frac{i}{2} q \mu)$, $f_1 = \hat{R}_\theta(i \sum_{j=1}^g C_{j, \theta} \delta(z, a_j))$ and $f = f_0 + f_1$. By (3.2) we have $\mu - 1 = R_{\lambda, \theta} f = R_{\lambda, \theta} f_0 + R_{\lambda, \theta} f_1$.

Put

$$L_{0, q}^{p, \tilde{p}}(V) = \{u : u|_Y \in L_{0, q}^p(Y), u|_{V \setminus Y} \in L_{0, q}^{\tilde{p}}(V \setminus Y)\}, \quad 1 \leq p < 2, \tilde{p} > 2, q = 0, 1.$$

By Proposition 3 ii' from [He] we obtain

$$\begin{aligned}
& \|\mu - \mu_\theta(\infty_l, \lambda)\|_{L^{\tilde{p}}(V \setminus Y)} \leq \\
& \text{const}(V, \tilde{p}, \theta) \cdot \min(|\lambda|^{-1/2}, |\lambda|^{-1}) (\|f_0\|_{\tilde{W}_{1,0}^{1, \tilde{p}}(V)} + \sum_{j=1}^g |C_{j, \theta}|) \quad (3.14) \\
& \|\partial \mu\|_{L_{1,0}^{p, \tilde{p}}(V)} \leq \text{const}(V, \tilde{p}, \theta) (\|f_0\|_{\tilde{W}_{1,0}^{1, \tilde{p}}(V)} + \sum_{j=1}^g |C_{j, \theta}|).
\end{aligned}$$

For proving estimates (3.1b) let us now estimate $\{C_{j, \theta}^0\}$.

In order to estimate $\{C_{j, \theta}^0\}$ we must use equations (3.6), where parameters $\{\tilde{C}_{j, \theta}\}$ are replaced by $\{C_{j, \theta}^0\}$ and function $\tilde{\mu}$ is replaced by 1. For modified equations (3.6)

1) we apply Kramer formula for solution of linear system and integration by parts in all integrals of this system, using $e_{\lambda, \theta}(z)(d\bar{z}_1 + \bar{\theta} d\bar{z}_2) = \frac{1}{\lambda} \bar{\theta} e_{\lambda, \theta}(z)$. In addition, we

use: formula (1.2) for $\Delta_\theta(\lambda)$, formula $\bar{\partial}\hat{R}_\theta(\frac{i}{2}q\mu) = \frac{i}{2}q\mu$ and estimate of singular integral, containing $\bar{\partial}(\frac{\bar{\omega}_k}{d\bar{z}_1 + \theta d\bar{z}_2})$. This gives inequality:

$$\sum_j |C_{j,\theta}^0(\lambda)| \leq \frac{\text{const}(V, \{a_j\}, \theta, \sigma)}{|\Delta_\theta(\lambda)|(1+|\lambda|)^g}.$$

ii) The equation (3.5) together with obtained inequality for $\sum |C_{j,\theta}^0(\lambda)|$, estimate of Faddeev type Green function $|g_{\lambda,\theta}(z, \xi)| = O(\frac{1}{|\lambda|^{1-\varepsilon}})$ are used to obtain estimate (3.15) for $\sum |\tilde{C}_{j,\theta}(\lambda)|$ and $|\mu_\theta(\lambda)|$:

$$\begin{aligned} |\lambda|^{-\varepsilon} \|\mu\|_{\tilde{W}^{1,\tilde{p}}(V)} + \sum_j |\tilde{C}_{j,\theta}(\lambda)| &\leq \frac{\text{const}(V, \{a_j\}, \theta, \sigma, \tilde{p}, \varepsilon)}{|\Delta_\theta(\lambda)|(1+|\lambda|)^g} \quad \text{and} \\ \|\mu - \mu(\infty_l, \cdot)\|_{L^{\tilde{p}}(V_l \setminus Y)} &\leq \frac{\text{const}(V, \{a_j\}, \theta, \sigma, \tilde{p}, \varepsilon)}{|\Delta_\theta(\lambda)|(1+|\lambda|)^{g+1-\varepsilon}}, \end{aligned} \quad (3.15)$$

where $\lambda \in \mathbb{C} \setminus E_\theta : |\lambda| \geq \text{const}(V, \{a_j\}, \theta, \sigma, \varepsilon)$, $l = 1, \dots, d$, $\mu_\theta(\infty_1, \lambda) = 1$.

These estimates imply estimates (3.1b).

c) Differentiation of equation (3.2) with respect to $\bar{\lambda}$ gives equality

$$\begin{aligned} \frac{\partial \mu}{\partial \bar{\lambda}} - R_{\lambda,\theta} \circ \left(\hat{R}_\theta \left(\frac{i}{2}q \frac{\partial \mu}{\partial \bar{\lambda}} + i \sum_{j=1}^g \frac{\partial C_{j,\theta}(\lambda)}{\partial \bar{\lambda}} \delta(z, a_j) \right) \right) = \\ (\bar{z}_1 + \bar{\theta} \bar{z}_2)(\mu - 1) - R_{\lambda,\theta} \left((\bar{\xi}_1 + \bar{\theta} \bar{\xi}_2) \hat{R}_\theta \left(\frac{i}{2}q\mu + i \sum_{j=1}^g C_{j,\theta} \delta(z, a_j) \right) \right). \end{aligned} \quad (3.16)$$

Equality (3.16) can be rewritten in the following form

$$\begin{aligned} \frac{\partial \mu}{\partial \bar{\lambda}} = (I - R_{\lambda,\theta} \circ \hat{R}_\theta \left(\frac{i}{2}q \cdot \right))^{-1} [(\bar{z}_1 + \bar{\theta} \bar{z}_2)(\mu - 1) + R_{\lambda,\theta} \circ \hat{R}_\theta \left(i \sum_{j=1}^g \frac{\partial C_{j,\theta}}{\partial \bar{\lambda}} \delta(z, a_j) \right) - \\ R_{\lambda,\theta} \left((\bar{\xi}_1 + \bar{\theta} \bar{\xi}_2) \hat{R}_\theta \left(\frac{i}{2}q\mu + i \sum_{j=1}^g C_{j,\theta}(\lambda) \delta(z, a_j) \right) \right)]. \end{aligned} \quad (3.17)$$

Using Propositions 2, 3 from [He], estimates from part (b) of this proof we obtain from (3.17)

$$\begin{aligned} e_{\lambda,\theta}(z) \frac{\partial \mu}{\partial \bar{\lambda}} \Big|_Y &\in W^{1,p}(Y), \\ e_{\lambda,\theta}(z) \frac{\partial \mu}{\partial \bar{\lambda}} \Big|_{V_l} &\in W^{1,\tilde{p}}(V_l \setminus Y) \cup L^\infty(V_l \setminus Y). \end{aligned}$$

Statement (3.1c) is proved.

Proposition 3.1 is proved.

4. Equation $\frac{\partial \mu(z, \lambda)}{\partial \bar{\lambda}} = b_\theta(\lambda) e_{-\lambda, \theta}(z) \overline{\mu_\theta(z, \lambda)}$. Proof of Theorem 1.1B

Proposition 4.1. *Let conductivity σ , divisor $\{a_1, \dots, a_g\}$ and θ satisfy the conditions of Proposition 2.1. Let function $\psi_\theta(z, \lambda) = e^{\lambda(z_1 + \theta z_2)} \mu_\theta(z, \lambda)$ be the Faddeev type function, associated with σ , θ and divisor $\{a_1, \dots, a_g\}$. Then for $\lambda \in \mathbb{C} \setminus E_\theta : |\lambda| \geq \text{const}(V, \{a_j\}, \theta, \sigma)$*

i) the following $\bar{\partial}$ -equations take place

$$\frac{\partial \mu_\theta(z, \lambda)}{\partial \bar{\lambda}} = b_\theta(\lambda) e_{-\lambda, \theta}(z) \overline{\mu_\theta(z, \lambda)}, \quad \text{if } z \in V \setminus \{a_1, \dots, a_g\}, \quad (4.1)$$

$$\frac{\partial C_{j, \theta}(\lambda)}{\partial \bar{\lambda}} = b_\theta(\lambda) e_{-\lambda, \theta}(a_j) \overline{C_{j, \theta}(\lambda)}, \quad j = 1, \dots, g, \quad \text{where} \quad (4.2)$$

ii) function $b_\theta(\lambda)$ satisfies equations:

$$b_\theta(\lambda) \lim_{\substack{z \rightarrow \infty \\ z \in V_l}} \overline{\mu_\theta(z, \lambda)} = \lim_{\substack{z \rightarrow \infty \\ z \in V_l}} \frac{\bar{z}_1 + \bar{\theta} \bar{z}_2}{\bar{\lambda}} e_{\lambda, \theta}(z) \frac{\partial \mu_\theta(z, \lambda)}{\partial (\bar{z}_1 + \bar{\theta} \bar{z}_2)}, \quad (4.3)$$

$$\bar{\lambda} b_\theta(\lambda) d = -\frac{1}{2\pi i} \int_{z \in bX} e_{\lambda, \theta}(z) \bar{\partial} \mu_\theta(z, \lambda) + i \sum_{j=1}^g C_{j, \theta}(\lambda) e_{\lambda, \theta}(a_j), \quad l = 1, \dots, d$$

and the inequality

$$|\lambda| (1 + |\lambda|)^g |\Delta_\theta(\lambda)| \cdot |b_\theta(\lambda)| \leq \text{const}(V, \{a_j\}, \theta, \sigma) \frac{1}{(|\lambda| + 1)^{1/3}}. \quad (4.4)$$

Remark 4.1. For the case $V = \mathbb{C}$ this statement is obtained in [GN], [N2], [N3]. Proposition 4.1 is a corrected version of Proposition 3.2 of [HM2].

Lemma 4.1. i) *Let function $\mu = \mu_\theta(z, \lambda)$, $z \in V \setminus Y$, $\lambda \in \mathbb{C} \setminus E_\theta : |\lambda| \geq \text{const}(V, \{a_j\}, \theta, \sigma)$ satisfy equation*

$$\bar{\partial}(\partial + \lambda(dz_1 + \theta dz_2))\mu = 0 \quad \text{on } V \setminus Y \quad (4.5)$$

and the property

$$[\mu - \mu_\theta(\infty_l, \lambda)]|_{V_l \setminus Y} \in W^{1, \tilde{p}}(V_l \setminus \bar{Y}), \quad \text{where } \tilde{p} > 2,$$

$$\mu_\theta(\infty_l, \lambda) \stackrel{\text{def}}{=} \lim_{\substack{z \rightarrow \infty \\ z \in V_l}} \mu_\theta(z, \lambda), \quad l = 1, \dots, d.$$

Then

$$A \stackrel{\text{def}}{=} \frac{\partial \mu}{\partial (z_1 + \theta z_2)} + \lambda \mu \in \mathcal{O}(\tilde{V} \setminus \bar{Y}) \quad \text{and}$$

$$A|_{V_l \setminus Y} = \lambda \mu(\infty_l) + \sum_{k=1}^{\infty} A_{k, l} \frac{1}{(z_1 + \theta z_2)^k},$$

$$\begin{aligned}\bar{B} &\stackrel{\text{def}}{=} e_{\lambda,\theta}(z) \frac{\partial\mu}{\partial(\bar{z}_1 + \bar{\theta}\bar{z}_2)} \in \overline{\mathcal{O}(\tilde{V}\setminus\bar{Y})} \quad \text{and} \\ \bar{B}|_{V_i\setminus Y} &= \sum_{k=1}^{\infty} B_{k,l} \frac{1}{(\bar{z}_1 + \bar{\theta}\bar{z}_2)^k}, \quad l = 1, \dots, d,\end{aligned}\tag{4.6}$$

where $\mathcal{O}(\tilde{V}\setminus\bar{Y})$ is the space of holomorphic functions on $(\tilde{V}\setminus\bar{Y})$.

ii) Let

$$M|_{V_i} = \mu_\theta(\infty_l, \lambda) + \sum_{k=1}^{\infty} \frac{a_{k,l}(\lambda)}{(z_1 + \theta z_2)^k} \quad \text{and} \quad \bar{N}|_{V_i} = \sum_{k=1}^{\infty} \frac{b_{k,l}(\lambda)}{(\bar{z}_1 + \bar{\theta}\bar{z}_2)^k}$$

be formal series with coefficients determined by relations

$$\lambda a_{k,l} - (k-1)a_{k-1,l} = A_{k,l}, \quad \bar{\lambda} b_{k,l} - (k-1)b_{k-1,l} = B_{k,l}, \quad l = 1, \dots, d, \quad k = 1, 2, \dots$$

Let

$$M_\nu|_{V_i} = \mu_\theta(\infty_l, \lambda) + \sum_{k=1}^{\nu} \frac{a_{k,l}}{(z_1 + \theta z_2)^k}, \quad \bar{N}_\nu|_{V_i} = \sum_{k=1}^{\nu} \frac{b_{k,l}}{(\bar{z}_1 + \bar{\theta}\bar{z}_2)^k}.\tag{4.7}$$

Then function μ has the asymptotic decomposition

$$\begin{aligned}\mu|_{V_i} &= M|_{V_i} + e_{-\lambda,\theta}(z)\bar{N}|_{V_i}, \quad z_1 \rightarrow \infty, \\ \text{i.e. } \mu|_{V_i} &= M|_{V_i} + e_{-\lambda,\theta}(z)\bar{N}_\nu|_{V_i} + O\left(\frac{1}{|z_1|^{\nu+1}}\right).\end{aligned}$$

Proof of Lemma 4.1.

i) From (4.5) it follows that

$$\partial\bar{\partial}(e^{\lambda(z_1+\theta z_2)}\mu(z, \lambda))|_{V\setminus\bar{Y}} = 0.$$

Thus $\bar{\partial}(e^{\lambda(z_1+\theta z_2)}\mu(z, \lambda)) = e^{\lambda(z_1+\theta z_2)}\bar{\partial}\mu$ is antiholomorphic form on $V\setminus\bar{Y}$ and $\partial\mu + \lambda\mu(dz_1 + \theta dz_2)$ is holomorphic form on $V\setminus\bar{Y}$. From this, condition $\bar{\partial}\mu \in L_{0,1}^{\bar{p}}(V\setminus\bar{Y})$ and the Cauchy theorem it follows that

$$\begin{aligned}e^{\lambda(z_1+\theta z_2)}\bar{\partial}\mu|_{V_i\setminus\bar{Y}} &= e^{\bar{\lambda}(\bar{z}_1+\bar{\theta}\bar{z}_2)}\bar{B}(d\bar{z}_1 + \bar{\theta}d\bar{z}_2)|_{V_i\setminus\bar{Y}} = \\ e^{\bar{\lambda}(\bar{z}_1+\bar{\theta}\bar{z}_2)} \sum_{k=1}^{\infty} \frac{B_{k,l}}{(\bar{z}_1 + \bar{\theta}\bar{z}_2)^k} (d\bar{z}_1 + \bar{\theta}d\bar{z}_2)|_{V_i} \quad \text{and} \\ (\partial\mu + \lambda\mu(dz_1 + \theta dz_2))|_{V_i\setminus\bar{Y}} &= A(dz_1 + \theta dz_2)|_{V_i\setminus\bar{Y}} = \\ (\lambda\mu(\infty_l) + \sum_{k=1}^{\infty} \frac{A_{k,l}}{(z_1 + \theta z_2)^k})(dz_1 + \theta dz_2)|_{V_i\setminus\bar{Y}}.\end{aligned}$$

It gives (4.6).

ii) From (4.6), (4.7) we obtain, first, that

$$\begin{aligned}\bar{\partial}\mu|_{V_i} &= e^{-\lambda(z_1+\theta z_2)}\bar{\partial}(e^{\bar{\lambda}(\bar{z}_1+\bar{\theta}\bar{z}_2)}\bar{N}_\nu)|_{V_i} + O\left(\frac{1}{|\bar{z}_1|^{\nu+1}}\right) \\ \text{then } \mu|_{V_i} &= M_\nu|_{V_i} + e_{-\lambda,\theta}(z)\bar{N}_\nu|_{V_i} + \tilde{O}\left(\frac{1}{|\bar{z}_1|^\nu}\right).\end{aligned}\tag{4.8}$$

Comparison of the last equality for different indexes ν and $\nu + 1$ implies that $\tilde{O}\left(\frac{1}{|\bar{z}_1|^\nu}\right) = O\left(\frac{1}{|\bar{z}_1|^{\nu+1}}\right)$.

It gives statement of Lemma 4.1.

Lemma 4.2. *i) Functions M_ν and N_ν (conjugated to \bar{N}_ν) from decomposition (4.8) have the following properties:*

$$\begin{aligned}\forall z \in \tilde{V} \setminus Y \exists \lim_{\nu \rightarrow \infty} \left(\frac{\partial M_\nu}{\partial(z_1 + \theta z_2)} + \lambda M_\nu \right) &\stackrel{\text{def}}{=} \frac{\partial M}{\partial(z_1 + \theta z_2)} + \lambda M \quad \text{and} \\ \exists \lim_{\nu \rightarrow \infty} \left(\frac{\partial N_\nu}{\partial(z_1 + \theta z_2)} + \lambda N_\nu \right) &\stackrel{\text{def}}{=} \frac{\partial N}{\partial(z_1 + \theta z_2)} + \lambda N.\end{aligned}$$

ii) *Functions $\frac{\partial M}{\partial(z_1 + \theta z_2)} + \lambda M$ and $\frac{\partial N}{\partial(z_1 + \theta z_2)} + \lambda N$ belongs to $\mathcal{O}(\tilde{V} \setminus Y)$ and*

$$\begin{aligned}\frac{\partial \mu}{\partial(\bar{z}_1 + \bar{\theta}\bar{z}_2)} &= e_{-\lambda,\theta}(z) \left(\frac{\partial \bar{N}}{\partial(\bar{z}_1 + \bar{\theta}\bar{z}_2)} + \bar{\lambda}\bar{N} \right), \\ \frac{\partial \mu}{\partial(z_1 + \theta z_2)} + \lambda \mu &= \frac{\partial M}{\partial(z_1 + \theta z_2)} + \lambda M,\end{aligned}\tag{4.9}$$

$$\frac{\partial N}{\partial(z_1 + \theta z_2)} + \lambda N \rightarrow 0, \quad z_1 \rightarrow \infty.\tag{4.10}$$

Proof of Lemma 4.2.

Part i) and equalities (4.9), (4.10) from part ii) follow directly from (4.8).

Properties (4.8), (4.9), (4.10), property $\bar{\partial}\mu \in L_{0,1}^{p,\bar{p}}$ (Proposition 3.1b) and extension property of bounded holomorphic functions through isolated singularities imply that

$$\frac{\partial M}{\partial(z_1 + \theta z_2)} + \lambda M \quad \text{and} \quad \frac{\partial N}{\partial(z_1 + \theta z_2)} + \lambda N$$

belongs to $\mathcal{O}(\tilde{V} \setminus Y)$.

Lemma 4.2 is proved.

Lemma 4.3. *Let $\psi_\theta(z, \lambda) = e^{\lambda(z_1 + \theta z_2)}\mu_\theta(z, \lambda)$ be the Faddeev type function on V , associated with potential $q = \frac{dd^c\sqrt{\sigma}}{\sqrt{\sigma}}$ and divisor $\{a_1, \dots, a_g\}$ on $Y \setminus \bar{X}$. Then*

$\forall \lambda \in \mathbb{C} \setminus E_\theta : |\lambda| \geq \text{const}(V, \{a_j\}, \theta, \sigma)$

$$e_{\lambda, \theta}(z) \frac{\partial \mu}{\partial(\bar{z}_1 + \bar{\theta} \bar{z}_2)} \Big|_{V_i \setminus \bar{Y}} = \sum_{k=1}^{\infty} B_{k,l} (\bar{z}_1 + \bar{\theta} \bar{z}_2)^{-k}, \quad \text{where}$$

$$B_{1,l} = -\frac{1}{2\pi i} \times \int_{\{z \in V_i : |z_1| = r_1\}} e_{\lambda, \theta}(z) \frac{\partial \mu}{\partial(\bar{z}_1 + \bar{\theta} \bar{z}_2)} (d\bar{z}_1 + \bar{\theta} d\bar{z}_2) \quad \forall r_1 : Y \subset \{z \in V : |z_1| < r_1\}. \quad (4.11)$$

Proof of Lemma 4.3.

Estimate of $\partial \mu$ from (3.1b) and the Cauchy theorem, applied to antiholomorphic function $e_{\lambda, \theta}(z) \frac{\partial \mu}{\partial(\bar{z}_1 + \bar{\theta} \bar{z}_2)} \Big|_{V_i \setminus \bar{Y}}$ implies (4.11).

Proof of Proposition 4.1.

Since ψ, μ are Faddeev type functions, we have the equations

$$\bar{\partial}(\partial + \lambda(dz_1 + \theta dz_2))\mu = \frac{i}{2} q\mu + i \sum_{j=1}^{\infty} C_{j, \theta}(\lambda) \delta(z, a_j),$$

$$dd^c \psi = q\psi + 2 \sum_{j=1}^g e^{\lambda(z_1 + \theta z_2)} C_{j, \theta}(\lambda) \delta(z, a_j).$$

Put $\psi_{\bar{\lambda}} = \frac{\partial \psi}{\partial \bar{\lambda}}$ and $\mu_{\bar{\lambda}} = \frac{\partial \mu}{\partial \bar{\lambda}}$.

We obtain

$$dd^c \psi_{\bar{\lambda}} = q\psi_{\bar{\lambda}} + 2 \sum_{j=1}^g e^{\lambda(z_1 + \theta z_2)} \frac{\partial C_{j, \theta}}{\partial \bar{\lambda}}(\lambda) \delta(z, a_j).$$

From Lemma 4.1 we deduce

$$\frac{\partial \mu}{\partial(\bar{z}_1 + \bar{\theta} \bar{z}_2)} \Big|_{V_i \setminus \bar{Y}} = e_{-\lambda, \theta}(z) \frac{B_{1,l}(\lambda)}{\bar{z}_1 + \bar{\theta} \bar{z}_2} + O\left(\frac{1}{|z_1|^2}\right), \quad \text{and}$$

$$\left(\frac{\partial \mu}{\partial(z_1 + \theta z_2)} + \lambda \mu\right) \Big|_{V_i \setminus \bar{Y}} = \lambda \mu(\infty_l) + \frac{A_{1,l}(\lambda)}{z_1 + \theta z_2} + O\left(\frac{1}{|z_1|^2}\right). \quad (4.12)$$

From (4.6), (4.7), (4.8) we deduce

$$\mu \Big|_{V_i \setminus \bar{Y}} = \mu_\theta(\infty_l, \lambda) + \frac{a_l(\lambda)}{z_1 + \theta z_2} + e_{-\lambda, \theta}(z) \frac{b_l(\lambda)}{\bar{z}_1 + \bar{\theta} \bar{z}_2} + O\left(\frac{1}{|z_1|^2}\right), \quad z_1 \rightarrow \infty, \quad (4.13)$$

$$\text{where } \bar{\lambda} b_l(\lambda) \stackrel{\text{def}}{=} \bar{\lambda} b_{1,l}(\lambda) = B_{1,l}, \quad \lambda a_l(\lambda) \stackrel{\text{def}}{=} \lambda a_{1,l}(\lambda) = A_{1,l}, \quad l = 1, \dots, d. \quad (4.14)$$

From (4.13) and (3.1c) we obtain for $l = 1, \dots, d$

$$\begin{aligned}\psi|_{V_l \setminus Y} &= e^{\lambda(z_1 + \theta z_2)} \mu = \\ &e^{\lambda(z_1 + \theta z_2)} \left(\mu_\theta(\infty_l, \lambda) + \frac{a_l(\lambda)}{z_1 + \theta z_2} + e^{\bar{\lambda}(\bar{z}_1 + \bar{\theta} \bar{z}_2) - \lambda(z_1 + \theta z_2)} \frac{b_l(\lambda)}{\bar{z}_1 + \bar{\theta} \bar{z}_2} + O\left(\frac{1}{|z_1|^2}\right) \right), \\ \psi_{\bar{\lambda}}|_{V_l \setminus Y} &= \frac{\partial \psi}{\partial \bar{\lambda}}|_{V_l \setminus Y} = e^{\bar{\lambda}(\bar{z}_1 + \bar{\theta} \bar{z}_2)} \left[(\bar{z}_1 + \bar{\theta} \bar{z}_2) \frac{b_l(\lambda) + e_{\lambda, \theta}(z) \frac{\partial \mu_\theta(\infty_l, \lambda)}{\partial \lambda}}{\bar{z}_1 + \bar{\theta} \bar{z}_2} + O\left(\frac{1}{|z_1|}\right) \right] = \\ &e^{\bar{\lambda}(\bar{z}_1 + \bar{\theta} \bar{z}_2)} \left(b_l(\lambda) + e_{\lambda, \theta}(z) \frac{\partial \mu_\theta(\infty_l, \lambda)}{\partial \bar{\lambda}} + O\left(\frac{1}{|z_1|}\right) \right).\end{aligned}$$

For function $\mu_{\bar{\lambda}} = e^{-\lambda(z_1 + \theta z_2)} \psi_{\bar{\lambda}}$ we obtain

$$\begin{aligned}\bar{\partial}(\partial + \lambda(dz_1 + \theta dz_2))\mu_{\bar{\lambda}} &= \frac{i}{2} q \mu_{\bar{\lambda}} + i \sum_{j=1}^g \frac{\partial C_{j, \theta}}{\partial \bar{\lambda}} \delta(z, a_j) \\ \text{and } \mu_{\bar{\lambda}} &= e_{-\lambda, \theta}(z) \left(b_l(\lambda) + e_{\lambda, \theta}(z) \frac{\partial \mu_\theta(\infty_l, \lambda)}{\partial \bar{\lambda}} + O\left(\frac{1}{|z_1|}\right) \right), \quad z \in V_l.\end{aligned}$$

For z_1 large enough function $e_{-\lambda, \theta}(z) \bar{\mu}_{\bar{\lambda}} \stackrel{\text{def}}{=} \varphi(z, \lambda)$ satisfies equation $\bar{\partial}(\partial + \lambda(dz_1 + \theta dz_2))\varphi = 0$. From this, Lemma 4.1 and property $\overline{\lim}_{z \rightarrow \infty} |\varphi(z, \lambda)|_V < \infty$ we deduce that $\varphi|_{V_l}(z, \lambda) \rightarrow \text{const}_l(\lambda) \stackrel{\text{def}}{=} \varphi(\infty_l, \lambda)$, if $z \in V_l$, $z \rightarrow \infty$, $l = 1, \dots, d$. So in the relations above we have $e_{\lambda, \theta}(z) \mu_{\bar{\lambda}}(\infty_l, \lambda) \equiv 0$, $l = 1, \dots, d$. Functions $e_{-\lambda, \theta}(z) \bar{\mu}_{\bar{\lambda}}$ and μ both satisfy equation $\bar{\partial}(\partial + \lambda(dz_1 + \theta dz_2))\mu = \frac{i}{2} q \mu$ on $V \setminus \{a_1, \dots, a_g\}$. Besides $\mu|_{V_l}(z, \lambda) \rightarrow \overline{\mu(\infty_l, \lambda)}$ and $e_{\lambda, \theta}(z) \mu_{\bar{\lambda}}(z, \lambda) \rightarrow b_l(\lambda)$, if $z \in V_l$, $z \rightarrow \infty$. Applying Proposition 2.1 we obtain

$$e_{\lambda, \theta}(z) \mu_{\bar{\lambda}} = b_l(\lambda) \bar{\mu}_\theta(z, \lambda) (\overline{\mu_\theta(\infty_l, \lambda)})^{-1}, \quad l = 1, \dots, d.$$

This implies equalities (4.1), (4.2), where

$$b_\theta(\lambda) = \frac{b_l(\lambda)}{\overline{\mu_\theta(\infty_l, \lambda)}}, \quad l = 1, \dots, d. \quad (4.15)$$

Asymptotic formula (4.3) follows from (4.11), (4.14) and (4.15). These formulas and Cauchy-Green formula imply also the following important expression for $b_\theta(\lambda)$:

$$\bar{\lambda} b_\theta(\lambda) d = -\frac{1}{2\pi i} \int_{z \in bY} e_{\lambda, \theta}(z) \bar{\partial} \mu = -\frac{1}{2\pi i} \int_{z \in bX} e_{\lambda, \theta}(z) \bar{\partial} \mu + i \sum_{j=1}^g C_{j, \theta} e_{\lambda, \theta}(a_j), \quad (4.16)$$

where

$$\int_{z \in bX} e_{\lambda, \theta}(z) \bar{\partial} \mu = \int_X \frac{1}{2i} e_{\lambda, \theta}(z) q \mu. \quad (4.17)$$

Equality (4.3) follows from (4.16). This equality together with estimate of $\{C_j\}$ from Lemma 2.4 and estimate through integration by parts of $\int_X e_{\lambda,\theta} q \mu$ imply (4.4).

Proposition 4.1 is proved.

5. Reconstruction of function $\psi_\theta|_{bX}$ from Dirichlet-to-Neumann data on bX . Proof of Theorem 1.2A

Let X be domain containing V_0 and relatively compact in V with smooth (of classe $C^{(2)}$) boundry. Let $\sigma \in C^{(2)}(V)$, $\sigma > 0$ on V , $\sigma = 1$ on $V \setminus X$. Let $q = \frac{dd^c \sqrt{\sigma}}{\sqrt{\sigma}}$. Let $u \in C(bX)$ and $\tilde{u} \in W^{1,\tilde{p}}(X)$, $\tilde{p} > 2$, be solution of the Dirichlet problem $d\sigma d^c \tilde{u}|_X = 0$, $\tilde{u}|_{bX} = u$, where $d^c = i(\bar{\partial} - \partial)$, $d = \bar{\partial} + \partial$. Let $\tilde{\psi} = \sqrt{\sigma} \tilde{u}$ and $\psi = \sqrt{\sigma} u$. Then

$$dd^c \tilde{\psi} = \frac{dd^c \sqrt{\sigma}}{\sqrt{\sigma}} \tilde{\psi} = q \tilde{\psi} \text{ on } X, \quad \tilde{\psi}|_{bX} = \psi. \quad (5.1)$$

Let ψ_0 be solution of Dirichlet problem

$$dd^c \psi_0|_X = 0, \quad \psi_0|_{bX} = \psi|_{bX}.$$

Let

$$\hat{\Phi} \psi = \bar{\partial} \tilde{\psi}|_{bX} \quad \text{and} \quad \hat{\Phi}_0 \psi = \bar{\partial} \tilde{\psi}_0|_{bX}. \quad (5.2)$$

Operator $\psi|_{bX} \mapsto \bar{\partial} \tilde{\psi}|_{bX}$ is equivalent to the Dirichlet-to-Neumann operator $u|_{bX} \mapsto \sigma d^c \tilde{u}|_{bX}$.

Proposition 5.1. *Let $\psi = e^{\lambda(z_1 + \theta z_2)} \mu$ be the Faddeev type function associated with potential $q = \frac{dd^c \sqrt{\sigma}}{\sqrt{\sigma}}$ (see Definition 1.4), generic divisor $\{a_1, \dots, a_g\}$ with support in $V \setminus \bar{X}$ and generic $\theta \in \mathbb{C}$. Then $\forall \lambda \in \mathbb{C} \setminus E_\theta : |\lambda| \geq \text{const}(V, \{a_j\}, \theta, \sigma)$ the restriction $\psi|_{bX}$ of ψ on bX can be found from Dirichlet-to-Neumann operator $\psi|_{bX} \rightarrow \sigma d^c \psi|_{bX}$ through the uniquely solvable Fredholm integral equation*

$$\mu_\theta(z, \lambda)|_{bX} + \int_{\xi \in bX} g_{\lambda,\theta}(z, \zeta) m_{-\lambda}(\hat{\Phi} - \hat{\Phi}_0) m_\lambda \mu_\theta(\zeta, \lambda) = 1 + i \sum_{j=1}^g C_{j,\theta}(\lambda) g_{\lambda,\theta}(z, a_j), \quad (5.3)$$

$$i \sum_{j=1}^g (a_{j,1} + \theta a_{j,2})^{-k} C_{j,\theta}(\lambda) + \int_{z \in bX} (z_1 + \theta z_2)^{-k} (\partial + \lambda(dz_1 + \theta dz_2)) \mu = 0, \quad k = 2, \dots, g+1,$$

where $g_{\lambda,\theta}(z, \xi)$ - kernel of operator $R_{\lambda,\theta} \circ \hat{R}_\theta$,

$$m_{-\lambda}(\hat{\Phi} - \hat{\Phi}_0) m_\lambda \mu_\theta(\zeta, \lambda) = \int_{w \in bX} e^{-\lambda(\zeta_1 + \theta \zeta_2)} (\Phi(\zeta, w) - \Phi_0(\zeta, w)) e^{\lambda(w_1 + \theta w_2)} \mu_\theta(w, \lambda), \quad (5.4)$$

$\Phi(\zeta, w)$, $\Phi_0(\zeta, w)$ are kernels of operators $\hat{\Phi}$ and $\hat{\Phi}_0$, $m_{\pm\lambda}$ denote the multiplication operators by $e^{\pm\lambda(z_1 + \theta z_2)}$, values $\{a_{j,1}\}$ of the first coordinate of points $\{a_j\}$ are supposed to be mutually different.

This proposition for the case $V = \mathbb{C}$ is equivalent to the second part of Theorem 1 from [N2].

Lemma 5.1. *Let $\psi = e^{\lambda(z_1 + \theta z_2)}\mu$ be Faddeev type function of Proposition 5.1. Then $\forall z \in V \setminus X$ and $\forall \lambda \in \mathbb{C} \setminus E_\theta : |\lambda| \geq \text{const}(V, \{a_j\}, \theta, \sigma)$ we have equalities*

$$\begin{aligned} \mu_\theta(z, \lambda) &= 1 - \int_{\xi \in bX} g_{\lambda, \theta}(z, \xi) \bar{\partial} \mu_\theta(\xi, \lambda) - \int_{\xi \in bX} \mu_\theta(z, \xi) e^{\lambda(\xi_1 + \theta \xi_2)} \partial (e^{-\lambda(\xi_1 + \theta \xi_2)} g_{\lambda, \theta}(z, \xi)) + \\ &i \sum_{j=1}^g C_{j, \theta}(\lambda) g_{j, \theta}(z, a_j) \end{aligned} \quad (5.5)$$

and

$$- \int_{\xi \in bX} (z_1 + \theta z_2)^{-k} (\partial + \lambda(dz_1 + \theta dz_2)) \mu = \sum_{j=1}^g (a_{j,1} + \theta a_{j,2})^{-k} i C_{j, \theta}(\lambda), \quad k = 2, \dots \quad (5.6)$$

Proof of Lemma 5.1.

The equation

$$\bar{\partial}(\partial + \lambda(dz_1 + \theta dz_2))\mu = \frac{i}{2} q \mu + i \sum_{j=1}^g C_{j, \theta}(\lambda) \delta(z, a_j), \quad (5.7)$$

where $\text{supp } q \subseteq X$ implies that $(1,0)$ -form $f = (\partial + \lambda(dz_1 + \theta dz_2))\mu$ is holomorphic on $(V \setminus (X \cup_{j=1}^g \{a_j\}))$ and $\text{Res}_{a_j}(\partial + \lambda(dz_1 + \theta dz_2))\mu = \frac{i C_j}{2\pi i}$. This and the property (4.12) imply that $\forall \lambda \in \mathbb{C} \setminus E_\theta$ and $\forall k \geq 2$ form $(z_1 + \theta z_2)^{-k} f$ is holomorphic in the neighborhood of $(\tilde{V} \setminus V)$. By residue theorem applied to the form $(z_1 + \theta z_2)^{-k} f$ on $\tilde{V} \setminus X$, we obtain

$$\int_{z \in bX} (z_1 + \theta z_2)^{-k} f(z, \lambda) = -2\pi i \sum_{j=1}^g \text{Res}_{a_j} (z_1 + \theta z_2)^{-k} f(z, \lambda) = -(a_{j,1} + \theta a_{j,2})^{-k} (i C_{j, \theta}(\lambda)),$$

$k = 2, 3, \dots$ Equality (5.6) is proved.

Let us prove now (5.5). Differential equation (5.7), where $\mu|_Y \in L^{\bar{p}}(Y)$, $\mu|_{V \setminus \bar{Y}} \in L^\infty(V \setminus \bar{Y})$, $\mu(z) \rightarrow 1, z \rightarrow \infty, z \in V_1$, is equivalent by Lemma 3.1 to the system of equations

$$\mu(z, \lambda) = 1 + R_{\lambda, \theta} \circ \hat{R}_\theta \left(\frac{i}{2} q \mu + i \sum_{j=1}^g C_j \delta(z, a_j) \right), \quad z \in V, \quad \text{and} \quad (5.8)$$

$$\bar{\partial}(\partial + \lambda(dz_1 + \theta dz_2))\mu = 0, \quad z \in V \setminus (X \cup_{j=1}^g \{a_j\}). \quad (5.9)$$

These equations imply relations (5.6). Besides, we have equality

$$\int_X g_{\lambda, \theta}(z, \xi) \frac{i}{2} q(\xi) \mu(\xi) = \int_X g_{\lambda, \theta}(z, \xi) \bar{\partial}(\partial + \lambda(dz_1 + \theta dz_2))\mu.$$

Using Green-Riemann formula we obtain

$$\begin{aligned} \int_X e^{\lambda((z_1-\xi_1)+\theta(z_2-\xi_2))} g_{\lambda,\theta}(z, \xi) \partial \bar{\partial} \psi &= \int_X \psi \partial \bar{\partial} (e^{\lambda((z_1-\xi_1)+\theta(z_2-\xi_2))} g_{\lambda,\theta}(z, \xi)) + \\ \int_{bX} e^{\lambda((z_1-\xi_1)+\theta(z_2-\xi_2))} g_{\lambda,\theta}(z, \xi) \bar{\partial} \psi &+ \int_{bX} \psi \partial (e^{\lambda((z_1-\xi_1)+\theta(z_2-\xi_2))} g_{\lambda,\theta}(z, \xi)). \end{aligned}$$

For $z \in V \setminus X$ we have

$$\partial \bar{\partial} (e^{\lambda((z_1-\xi_1)+\theta(z_2-\xi_2))} g_{\lambda,\theta}(z, \xi)) = 0.$$

Then

$$- \int_{\xi \in X} g_{\lambda,\theta}(z, \xi) \left(\frac{i}{2} q \mu\right) = \int_{\xi \in bX} g_{\lambda,\theta} \bar{\partial} \mu + \int_{\xi \in bX} e^{\lambda(\xi_1+\theta\xi_2)} \mu \partial (e^{-\lambda(\xi_1+\theta\xi_2)} g_{\lambda,\theta}(z, \xi)). \quad (5.10)$$

From (5.8), (5.10) we deduce statement (5.5) of Lemma 5.1.

Proof of Proposition 5.1.

Let $\psi_0 : \bar{\partial} \psi_0|_X = 0$ and $\psi_0|_{bX} = \psi$. By Green-Riemann formula $\forall z \in V \setminus X$ we have

$$\int_{\xi \in bX} \psi \partial (e^{\lambda((z_1-\xi_1)+\theta(z_2-\xi_2))} g_{\lambda,\theta}(z, \xi)) + \int_{\xi \in bX} e^{\lambda((z_1-\xi_1)+\theta(z_2-\xi_2))} g_{\lambda,\theta}(z, \xi) \bar{\partial} \psi_0 = 0. \quad (5.11)$$

Formulas (5.11) and (5.5), (5.6) imply

$$\begin{aligned} \psi(z, \lambda) &= e^{\lambda(z_1+\theta z_2)} - \int_{bX} e^{\lambda((z_1-\xi_1)+\theta(z_2-\xi_2))} g_{\lambda,\theta}(z, \xi) (\bar{\partial} \psi(\xi) - \bar{\partial} \psi_0(\xi)) + \\ &i \sum_{j=1}^g e^{\lambda(z_1+\theta z_2)} C_j g_{\lambda,\theta}(z, a_j). \end{aligned} \quad (5.12)$$

Formula (5.12), (5.6) are equivalent to (5.3). Integral equation (5.3) is the Fredholm equation in $C(bX)$, because operator $(\hat{\Phi} - \hat{\Phi}_0)$ is compact operator in $C(bX)$. Existence $\forall \lambda \in \mathbb{C} \setminus E_\theta$ of unique Faddeev type function $\psi = e^{\lambda(z_1+\theta z_2)} \mu$, associated with q and divisor $\{a_1, \dots, a_g\}$ imply existence of solution of (5.3) with residue data

$iC_j = \text{Res}_{a_j} (\partial + \lambda(dz_1 + \theta dz_2)) \mu$, $j = 1, \dots, g$. Let us prove uniqueness of solution (5.3) in $C(bX)$ with residue data $\{C_j\}$. Suppose $\mu \in C(bX)$ solves (5.3), (5.6). Consider this μ as Dirichlet data for equation $\bar{\partial}(\partial + \lambda(dz_1 + \theta dz_2)) \mu = \frac{i}{2} q \mu$ on X , solution of which well defines μ on \bar{X} .

Let us also define μ on $V \setminus \bar{X}$ by (5.5). Function $\mu(z, \lambda)$ defined in such a way on V belongs to $C(V \setminus \cup_{j=1}^g \{a_j\})$.

Let us show that μ satisfy (5.7). By Sohotsky-Plemelj jump formula $\forall z^* \in bX$ we have

$$\begin{aligned} \frac{i}{2}\mu(z^*) &= \lim_{\substack{z \rightarrow z^* \\ z \in X}} \left(\int_{bX} g_{\lambda, \theta} \bar{\partial} \mu + \mu e^{\lambda(\xi_1 + \theta \xi_2)} \partial (e^{-\lambda(\xi_1 + \theta \xi_2)} g_{\lambda, \theta}) \right) - \\ &- \lim_{\substack{z \rightarrow z^* \\ z \in V \setminus X}} \left(\int_{bX} g_{\lambda, \theta} \bar{\partial} \mu + \mu e^{\lambda(\xi_1 + \theta \xi_2)} \partial (e^{-\lambda(\xi_1 + \theta \xi_2)} g_{\lambda, \theta}) \right). \end{aligned} \quad (5.13)$$

From (5.5) and (5.13) we deduce equality

$$\mu - \frac{i}{2}\mu = 1 - \int_{\xi \in bX} g_{\lambda, \theta} \bar{\partial} \mu - \int_{\xi \in bX} \mu e^{\lambda(\xi_1 + \theta \xi_2)} \partial (e^{-\lambda(\xi_1 + \theta \xi_2)} g_{\lambda, \theta}) + i \sum_{j=1}^g C_j g_{\lambda, \theta}(z, a_j), \quad z \in X. \quad (5.14)$$

By Green-Riemann formula we have also

$$\begin{aligned} &- \int_{bX} g_{\lambda, \theta} \bar{\partial} \mu - \mu e^{\lambda(\xi_1 + \theta \xi_2)} \partial (e^{-\lambda(\xi_1 + \theta \xi_2)} g_{\lambda, \theta}) + i \sum_{j=1}^g C_j g_{\lambda, \theta}(z, a_j) = \\ &- \int_X \mu (\bar{\partial}(\partial + \lambda(dz_1 + \theta dz_2)) g_{\lambda, \theta}) + \int_X g_{\lambda, \theta} \bar{\partial}(\partial + \lambda(dz_1 + \theta dz_2)) \mu + i \sum_{j=1}^g C_j g_{\lambda, \theta}(z, a_j) = \\ &\begin{cases} \frac{\mu}{2i} + \int_X g_{\lambda, \theta} \bar{\partial}(\partial + \lambda(dz_1 + \theta dz_2)) \mu + i \sum_{j=1}^g C_j g_{\lambda, \theta}(z, a_j), \quad z \in X, \\ \int_X g_{\lambda, \theta} \bar{\partial}(\partial + \lambda(dz_1 + \theta dz_2)) \mu + i \sum_{j=1}^g C_j g_{\lambda, \theta}(z, a_j), \quad z \in V \setminus (X \cup_{j=1}^g \{a_j\}). \end{cases} \end{aligned} \quad (5.15)$$

Equalities (5.5), (5.6), (5.14) and (5.15) imply (3.3) and

$$\begin{aligned} \mu(z) &= 1 + \int_V g_{\lambda, \theta} \bar{\partial}(\partial + \lambda(dz_1 + \theta dz_2)) \mu + i \sum_{j=1}^g C_j g_{\lambda, \theta}(z, a_j) = \\ &1 + R_{\lambda, \theta} \circ \hat{R}_\theta \left(\frac{i}{2} q \mu + i \sum_{j=1}^g C_j g_{\lambda, \theta}(z, a_j) \right), \quad z \in V. \end{aligned}$$

By Lemma 3.1 function $\mu_\theta(z, \lambda)$ is the Faddeev type function associated with q and divisor $\{a_1, \dots, a_g\}$. The uniqueness of solution of (5.3) in $C(bX)$ with residue data $\{C_{j, \theta}\}$ follows now from uniqueness of the Faddeev type function.

6. Reconstruction of conductivity function from Dirichlet-to-Neumann data. Proof of Theorem 1.2B

We will obtain here exact formulas for reconstruction of conductivity function $\sigma \in C^{(3)}(V)$, $\sigma > 0$, $\sigma \equiv 1$ on $V \setminus X$, from Dirichlet-to-Neumann data

$$\psi_\theta|_{bX} \rightarrow \bar{\partial} \psi_\theta|_{bX}$$

for Faddeev type functions $\psi_\theta(z, \lambda) = e^{\lambda(z_1 + \theta z_2)} \mu_\theta(z, \lambda)$, $\theta \in \mathbb{C} \setminus \{\theta_1, \theta_d\}$,
 $\lambda \in \mathbb{C} \setminus E_\theta : |\lambda| \geq \text{const}(V, \{a_j\}, \theta, \sigma)$, $\{a_1, \dots, a_g\} \subset Y \setminus X$.

For simplicity of presentation we consider in detail the case of regular algebraic curves in $\mathbb{C}^2 \subset \mathbb{C}P^2$, only.

Let $\tilde{V} = \{\tilde{z} = (\tilde{z}_0 : \tilde{z}_1 : \tilde{z}_2) \in \mathbb{C}P^2 : \tilde{P}(\tilde{z}) = 0\}$, where $\tilde{P}(\tilde{z})$ homogeneous polynomial of degree N . Let $\mathbb{C}P_\infty^1 = \{\tilde{z} : \mathbb{C}P^2 : \tilde{z}_0 = 0\}$. Put

$$\begin{aligned} \mathbb{C}^2 &= \{\tilde{z} \in \mathbb{C}P^2 : \tilde{z}_0 \neq 0\}, \quad z_1 = \frac{\tilde{z}_1}{\tilde{z}_0}, \quad z_2 = \frac{\tilde{z}_2}{\tilde{z}_0}, \quad P(z) = \tilde{P}(1, z_1, z_2), \\ V &= \{z \in \mathbb{C}^2 : P(z) = 0\} = \tilde{V} \cap \mathbb{C}^2. \end{aligned} \quad (6.1)$$

Without restriction of generality we suppose that \tilde{V} be (regular) curve of degree $N \geq 2$ with property:

$$\begin{aligned} \tilde{V} \cap \mathbb{C}P_\infty^1 &= \{\beta_1, \dots, \beta_d\}, \quad \text{where } \beta_1, \dots, \beta_d \text{ be different points of } \mathbb{C}P_\infty^1, \\ \beta_l &= (0 : \beta_l^1 : \beta_l^2), \quad \frac{\beta_l^2}{\beta_l^1} \in \mathbb{C}, \quad l = 1, \dots, N, \\ \frac{\partial P}{\partial z_2}(z) &\neq 0, \quad \text{if } z \in V : |z_1| \geq r_0 = \text{const}(V). \end{aligned} \quad (6.2)$$

For $\theta \in \mathbb{C}$ let $\{w_m\}$ be points of V , where $(dz_1 + \theta dz_2)|_V(w_m) = 0$. Then for almost all $\theta \in \mathbb{C}$ the following relations are valid

$$\begin{aligned} \theta &= \frac{\partial P}{\partial z_2}(w_m) / \frac{\partial P}{\partial z_1}(w_m), \quad \frac{\partial P}{\partial z_1}(w_m) \neq 0, \\ &[\frac{\partial^2 P}{\partial z_1^2} (\frac{\partial P}{\partial z_2})^2 - 2 \frac{\partial^2 P}{\partial z_1 \partial z_2} (\frac{\partial P}{\partial z_2}) (\frac{\partial P}{\partial z_1}) + \frac{\partial^2 P}{\partial z_2^2} (\frac{\partial P}{\partial z_1})^2](w_m) \neq 0. \end{aligned}$$

Without restriction of generality it is sufficient to give proof under condition that $\theta = 0$, i.e. for points $w_m = (w_{m,1}, w_{m,2}) \in V$ such that

$$\frac{\partial P}{\partial z_1}(w_m) \neq 0, \quad \frac{\partial P}{\partial z_2}(w_m) = 0, \quad \frac{\partial^2 P}{\partial z_2^2}(w_m) \neq 0 \quad (6.3)$$

and also such that $\forall m$ the line $\{z \in \mathbb{C}^2 : z_1 = w_{m,1}\}$ has tangency with X only in the single point w_m , $m = 1, \dots, M$. By Hurwitz-Riemann formula $M = N(N - 1)$. In the neighborhood of point $w_m \in V$ curve V can be represented in the form

$$\begin{aligned} V &= \{(z_1, z_2) \in \mathbb{C}^2 : z_1 = w_{m,1} + \\ &(\frac{\partial P}{\partial z_1}(w_m))^{-1} [-\frac{1}{2} \frac{\partial^2 P}{\partial z_2^2}(w_m)(z_2 - w_{m,2})^2 + O((z_2 - w_{m,2})^3)]\}. \end{aligned} \quad (6.4)$$

The reconstruction formula for $\frac{dd^c \sqrt{\sigma}}{\sqrt{\sigma}}(w_m)$, $m = 1, \dots, M$, will be obtained here by the stationary phase method, using formula (4.17).

Let μ be Faddeev type function (3.1) with properties (3.1a)-(3.1c) and with $\theta = 0$.

Below in this section we will write $\hat{R}_0, R_{\lambda,0}, e_{\lambda,0}, \mu_0, \psi_0, \Delta_0, E_0, C_{j,0}$ as $\hat{R}, R_\lambda, e_\lambda, \mu, \psi, \Delta, E, C_j$.

Let

$$f_0 = F_0 dz_1 = \frac{i}{2} \hat{R}(q\mu), \quad f_1 = F_1 dz_1 = i \sum_{j=1}^g C_j(\lambda) \hat{R}(\delta(\cdot, a_j)),$$

where $\mu = \mu(z, \lambda), z \in V, \lambda \in \mathbb{C} \setminus E: |\lambda| \geq \text{const}(V, \{a_j\}, \sigma)$.

Lemma 6.1. *For $u_0 = R_\lambda f_0$ the following estimate holds:*

$$\|u_0(\cdot, \lambda) - \frac{F_0(\cdot, \lambda)}{\lambda}\|_{L^{9/4}(X)} \leq \frac{\text{const}(V, \tilde{p})}{|\lambda|^{7/5}} \|f_0(\cdot, \lambda)\|_{\tilde{W}_{1,0}^{2,\tilde{p}}(V)}.$$

Proof of Lemma 6.1. By Lemma 2.1 and Proposition 2 from [He] we have $f_0 \in \tilde{W}_{1,0}^{2,\tilde{p}}(V), F_0 \in \tilde{W}^{1,p}(V)$. Using equality $\partial_z e_\lambda(z) = \lambda e_\lambda(z) dz_1$ and integration by parts formula $u_0 = R_\lambda f_0 = e_{-\lambda}(z) \overline{R(e_\lambda f_0)}$ can be transformed into the following

$$\begin{aligned} u_0(z) &= e_{-\lambda}(z) \overline{R_1(e_\lambda f_0)} + e_{-\lambda}(z) \overline{R_0(e_\lambda f_0)} = \\ &= -\frac{e_{-\lambda}(z)}{2\pi i} \frac{1}{\lambda} \int_V \frac{e_\lambda(\xi) \partial F_0 \wedge \overline{d\xi_1} \det\left[\frac{\partial \bar{P}}{\partial \xi}(\xi), \xi - z\right]}{\frac{\partial \bar{P}}{\partial \xi_2}(\xi) \cdot |\xi - z|^2} - \\ &= -\frac{e_{-\lambda}(z)}{2\pi i} \frac{1}{\lambda} \int_V e_\lambda(\xi) F_0 \partial \left(\frac{\det\left[\frac{\partial \bar{P}}{\partial \xi}(\xi), \xi - z\right] \wedge \overline{d\xi_1}}{\frac{\partial \bar{P}}{\partial \xi_2}(\xi) \cdot |\xi - z|^2} \right) + e_{-\lambda}(z) \overline{R_0(e_\lambda f_0)}, \end{aligned} \quad (6.5)$$

where R_1, R_0 operators defined in section 1 (see remark 1.1). From (6.5), using Corollary 1.2 from [He], we deduce

$$\lambda u_0 - F_0 = -e_{-\lambda}(z) \overline{R_1(e_\lambda(\xi) \partial F_0)} - e_{-\lambda}(z) \overline{R_0(e_\lambda(\xi) \partial F_0)} \stackrel{\text{def}}{=} J_1(z) + J_0(z). \quad (6.6)$$

We will estimate further only term $J_1(z)$. Estimate for $J_0(z)$ is similar.

For $J_1(z)$ we have $J_1(z) = J_1^+(z) + J_1^-(z)$, where

$$J_1^\pm(z) = \frac{e_{-\lambda}(z)}{2\pi i} \int_V \frac{e_\lambda(\xi) \chi_\rho^\pm(\xi) \partial F_0(\xi) \wedge \overline{d\xi_2} \det\left[\frac{\partial \bar{P}}{\partial \xi}(\xi), \xi - z\right]}{\frac{\partial \bar{P}}{\partial \xi_1}(\xi) \cdot |\xi - z|^2},$$

χ_ρ^\pm be smooth functions such that $\chi_\rho^+ + \chi_\rho^- \equiv 1$,

$$\chi_\rho^+ = 1, \quad \text{if } \left| \frac{d\xi_1}{d\xi_2} \right| \leq \rho, \quad \text{supp } \chi_\rho^+ \subset \left\{ \xi : \left| \frac{d\xi_1}{d\xi_2} \right| \leq 2\rho \right\} \quad (6.7)$$

and $|d\chi_\rho^\pm| = O\left(\frac{1}{\rho}\right)$.

Let $B_0 = \{z \in V : d\xi_1|_V(z) = 0\}$. Property $\bar{\partial}F_0 = dz_1] \frac{i}{2} q\mu$ implies estimate $\bar{\partial}F_0 = O(\frac{1}{\text{dist}(z, B_0)}) dz_2$. From this, formula for $J_1^+(z)$ and Lemma 3.1 of [He] we obtain estimate for

$$J_1^+ : \|J_1^+\|_{L^{9/4}(X)} = O(\rho^{2/3}) \|f_0\|_{\tilde{W}_{1,0}^{2,\tilde{p}}(V)}. \quad (6.8)$$

In order to estimate $J_1^-(z)$ we integrate by parts in the formula for J_1^- , using $\partial_z e_\lambda(z) = \lambda e_\lambda(z) dz_1$. Then inequalities

$$|\bar{\partial}F_0(z)| = O\left(\frac{1}{\text{dist}(z, B_0)}\right), \quad |\bar{\partial}\partial F_0(z)| = O\left(\frac{1}{\text{dist}(z, B_0)}\right)^2, \quad z \in X \setminus B_0$$

and inequality

$$\left| \int_{\rho \leq |\xi_2| \leq 1} \frac{d\xi_2 \wedge d\bar{\xi}_2}{|\xi_2|^2 (\xi_2 - \bar{z}_2)} \right| + \left| \int_{\rho \leq |\xi_2| \leq 1} \frac{d\xi_2 \wedge d\bar{\xi}_2}{|\xi_2| (\xi_2 - \bar{z}_2)^2} \right| = O\left(\frac{1}{\rho}\right)$$

imply estimate

$$\|J_1^-\|_{L^\infty(X)} = O\left(\frac{1}{|\lambda|\rho}\right) \|f_0\|_{\tilde{W}_{1,0}^{2,\tilde{p}}(V)}. \quad (6.9)$$

From (6.6), (6.8), (6.9) with $\rho = |\lambda|^{-3/5}$ we obtain statement of Lemma 6.1.

Lemma 6.2. *Let $q \in C_{1,1}^{(1)}(V)$, $\text{supp } q \subseteq X$, $f_0 = \frac{i}{2} \hat{R}(q\mu)$, $u_0 = R_\lambda f_0$. Then the following asymptotic estimate is valid*

$$\left| \int_X e_\lambda(z) q(z) u_0(z) \right| = o\left(\frac{1}{|\lambda|}\right), \quad \text{for}$$

$$\lambda \in \mathbb{C} : |\lambda| \geq \text{const}(V, \{a_j\}, \sigma), \quad |\Delta(\lambda)(1 + |\lambda|)^g| \geq \delta > 0, \quad \text{for some sufficient small } \delta.$$

Proof of Lemma 6.2.

From Lemma 6.1, using estimate of μ from (3.1b), we obtain asymptotic relation in the space $L^{\tilde{p}}(V)$, $2 < \tilde{p} < 9/4$:

$$\begin{aligned} u_0(z, \lambda) &= \frac{F_0(z, \lambda)}{\lambda} + O\left(\frac{1}{|\lambda|^{7/5}}\right) = \\ &= \frac{dz_1] \frac{i}{2} \hat{R}(q)}{\lambda} + O\left(\frac{1}{|\lambda|^{7/5}}\right) \quad \text{if } |\Delta_\theta(\lambda)(1 + |\lambda|)^g| \geq \delta > 0. \end{aligned}$$

Putting this relation into $\int_X e_\lambda(z) q(z) u_0(z)$, we obtain

$$\int_X e_\lambda(z) q(z) u_0(z) = \frac{i}{2\lambda} \int_X e_\lambda(z) q(z) (dz_1] \hat{R}(q)) + O\left(\frac{1}{|\lambda|^{7/5}}\right).$$

By Riemann-Lebesgue type theorem

$$\int_X e_\lambda(z)q(z)(dz_1|\hat{R}(q)) = o(1) \text{ if } \lambda \rightarrow \infty, \quad |\Delta(\lambda)|(1+|\lambda|)^g \geq \delta > 0.$$

This implies the statement of Lemma 6.2.

Lemma 6.3. *Let $q \in C_{1,1}^{(1)}(V)$, $\text{supp } q \subset X$. Let w_1, \dots, w_M be the points, where $dz_1|_V(w_m) = 0$. Then the following consequence of stationary phase method is valid:*

$$\int_V e^{i\tau(z_1+\bar{z}_1)}q(z) = \sum_m (1+o(1)) \sum_{m=1}^M -\frac{\pi}{r} \frac{|\frac{\partial P}{\partial z_1}(w_m)|Q_2(w_m)}{|\frac{\partial^2 P}{\partial z_2^2}(w_m)|} e^{i\tau(w_{m,1}+\bar{w}_{m,1})}, \quad (6.10)$$

where $Q_2(w_m) = \frac{q}{2idz_2 \wedge d\bar{z}_2}(w_m)$.

Proof of Lemma 6.3 (see [Fe], Th.2.1).

Lemma 6.4. *Let $q = \frac{dd^c \sqrt{\sigma}}{\sqrt{\sigma}} \in C_{1,1}^{(1)}(X)$, $\text{supp } q \subset X$, $f_1 = i \sum_{j=1}^g C_j(\lambda) \hat{R}(\delta(\cdot, a_j))$, $u_1 = R_\lambda f_1$. Then the following asymptotic estimate is valid*

$$\left| \int_X e_\lambda(z)q(z)u_1(z) \right| = O\left(\frac{1}{|\lambda|^{3/2-\varepsilon}}\right), \text{ for } \lambda \in \mathbb{C} : |\lambda| \geq \text{const}(V, \{a_j\}, \sigma, \varepsilon),$$

$$|\Delta(\lambda)(1+|\lambda|)^g| \geq \delta > 0, \quad \delta \text{ for some sufficiently small } \delta.$$

Proof of Lemma 6.4. Using that $\{a_1, \dots, a_g\}$ be generic divisor, from estimate (3.7) (Lemma 3.3) we obtain inequality

$$\sup_{j,\lambda} |C_j(\lambda)| \leq \text{const}(V, \{a_j\}) \sup_k \left| \int_X e_\lambda(z) \left(i \frac{dd^c \sqrt{\sigma}}{\sqrt{\sigma}} + 2\bar{\partial} \ln \sqrt{\sigma} \wedge \partial \ln \sqrt{\sigma} \right) \mu \frac{\bar{\omega}_k}{d\bar{z}_1}(z) \right|.$$

Let $\varepsilon > 0$ be small enough and $B_\varepsilon = \{z \in X : \left| \frac{dz_1}{dz_2} \right| < \varepsilon\}$. Then

$$\left| \frac{\bar{\omega}_k(z)}{d\bar{z}_1} \right|_X = O\left(\sum_{m=1}^M \frac{1}{|z_2 - w_{m,k}|} \right), \quad z \in X.$$

Let $\chi_\rho^\pm \in C^{(1)}(X)$ be functions with properties (6.7). Using that $\sigma \in C^{(3)}(X)$, $\mu \in \tilde{W}^{1,\bar{p}}(X)$, $\partial_z e_\lambda(z) = \lambda e_\lambda(z) dz_1$ by integration by parts we obtain

$$\left| \int_X \chi_\rho^-(z) e_\lambda(z) \left(i \frac{dd^c \sqrt{\sigma}}{\sqrt{\sigma}} + 2\bar{\partial} \ln \sqrt{\sigma} \wedge \partial \ln \sqrt{\sigma} \right) \mu(z, \lambda) \frac{\bar{\omega}_k}{d\bar{z}_1}(z) \right| \leq$$

$$\frac{\text{const}(V, \sigma)}{\rho \lambda}.$$

We have also directly

$$\left| \int_X \chi_\rho^+(z) e_\lambda(z) \left(i \frac{dd^c \sqrt{\sigma}}{\sqrt{\sigma}} + 2\bar{\partial} \ln \sqrt{\sigma} \wedge \partial \ln \sqrt{\sigma} \right) \mu(z, \lambda) \frac{\bar{\omega}_k}{d\bar{z}_1}(z) \right| \leq \text{const}(V, \sigma) \rho.$$

These estimates with $\rho = \frac{1}{\sqrt{|\lambda|}}$ and estimates for Faddeev type Green function $|R_\lambda \circ \hat{R}(\delta(\cdot, a_j))| = O(\frac{1}{|\lambda|^{1-\varepsilon}})$ from Theorem 4 of [He] imply statement of Lemma 6.4.

Proposition 6.1. *Under conditions (6.1)-(6.4), for $\lambda = i\tau$: $\tau \in \mathbb{R}$, $|\tau^g \Delta(i\tau)| \geq \delta > 0$, δ -small enough, the following formula is valid*

$$\int_{z \in bX} e_{i\tau}(z) \bar{\partial}_z \mu(z, i\tau) = \int_{z \in X} e_{i\tau}(z) \frac{q\mu}{2i} = \frac{1 + o(1)}{\tau} \sum_{m=1}^M \frac{\pi i}{2} \frac{dd^c \sqrt{\sigma}}{\sqrt{\sigma} dd^c |z|^2} \Big|_V(w_m) e^{i\tau(w_{m,1} + \bar{w}_{m,1})} \left| \frac{\partial^2 P}{\partial z_2^2}(w_m) \right|^{-1} \frac{\partial P}{\partial z_1}(w_m). \quad (6.11)$$

Proof of Proposition 6.1 and Theorem 1.2B. From Lemma 3.1 we have equality

$$\mu = 1 + R_\lambda \circ \hat{R}\left(\frac{i}{2} q\mu\right) + R_\lambda \circ \hat{R}\left(i \sum_{j=1}^g C_j \delta(z, a_j)\right) = 1 + u_0 + u_1. \quad (6.12)$$

Let $\delta > 0$ be small enough. Estimates of Lemmas 6.2, 6.4 and (6.12) give asymptotic equality

$$\mu = 1 + o\left(\frac{1}{\lambda}\right) \quad (6.13)$$

under conditions $\lambda \in \mathbb{C}$: $|\lambda| \geq \text{const}(V, \{a_j\}, \sigma)$, $|\Delta(\lambda)(1 + |\lambda|)^g| \geq \delta > 0$.

By Proposition 1.1, $\forall \varepsilon > 0$ we have inequality

$$\underline{\lim}_{\lambda \rightarrow \infty} |\lambda^g \Delta(\lambda)|_\varepsilon = \delta(\varepsilon) > 0, \quad \text{where } |\lambda^g \Delta(\lambda)|_\varepsilon = \sup_{\{\lambda' : |\lambda' - \lambda| \leq \varepsilon\}} |\lambda' \Delta(\lambda')|.$$

So for any $\varepsilon > 0$ and any positive $\delta < \delta(\varepsilon)$ there exists r such that the set $\{\lambda \in \mathbb{C} : |\Delta(\lambda)(1 + |\lambda|)^g| \geq \delta > 0\}$ intersects any disk $\{\lambda' : |\lambda - \lambda'| < \varepsilon\}$, with $|\lambda| \geq r$. This property, Lemma 6.3 and property (6.13) imply Proposition 6.1.

Theorem 1.2B follows from Proposition 1. Indeed, stationary phase method permits differentiation of (6.11) with respect to τ , keeping (in our case) terms of order $\frac{1}{\tau}$. Differentiation of the right-hand side of (6.11) gives for $\theta = 0$ the right-hand side of (1.12).

Theorem 1.2B is proved.

Remark 6.1. To obtain version of Proposition 6.1 with arbitrary generic θ from Proposition 6.1 with $\theta = 0$ it is sufficient to change coordinate system: $\tilde{z}_1 = z_1 + \theta z_2$, $\tilde{z}_2 = z_2$.

Remark 6.2. Proposition 6.1 can be reformulated also as formula for reconstruction of conductivity function from scattering data $b_\theta(i\tau)$ and $C_{j,\theta}(i\tau)$. Indeed, by formula (4.16), we have

$$\int_{bX} e_{i\tau,\theta}(z) \bar{\partial} \mu(z, i\tau) = -2\pi [\tau b_\theta(i\tau) d + \sum_{j=1}^g C_{j,\tau}(i\tau) e_{i\tau,\theta}(a_j)],$$

where d is defined in section 1.

7. Proof of Proposition 1.1

For simplicity of presentation we give proof only for the case when V is algebraic curve in \mathbb{C}^2 . Proposition 1.1 will be obtained here as a corollary of the following statement.

Proposition 7.1. *Let $\theta \in \mathbb{C} \setminus \{\theta_1, \dots, \theta_N\}$, $\delta = \delta(\theta) = \inf_l |\theta - \theta_l|$, $V_0 = \{z \in V : |z_1| \leq r_0(\delta)\}$, $r_0(\delta) = \frac{\text{const}(V)}{\sqrt{\delta}}$. Let $\{b_m\}$ be the points of V , where $(dz_1 + \theta dz_2)|_V(b_m) = 0$, $m = 1, \dots, M$, and $\{a_1, \dots, a_g\}$ be the points of generic divisor in $V \setminus \bar{V}_0$. Then $\forall j, k = 1, \dots, g$ and for $\lambda = i\tau$, where $\tau \in \mathbb{R}$, large enough, such that $|\Delta_\theta(i\tau)| \geq \delta > 0$, the following asymptotic equality is valid*

$$\begin{aligned} \int_V \hat{R}_\theta(\delta(\xi, a_j)) \wedge \bar{\omega}_k(\xi) e_{\lambda,\theta}(\xi) &= -\frac{1}{\lambda} e_{\lambda,\theta}(a_j) \frac{\bar{\omega}_k}{d\bar{\xi}_1 + \theta d\bar{\xi}_2}(a_j) - \\ &- \frac{\pi}{|\lambda|} \sum_{m=1}^M \exp[\lambda(b_{m,1} + \theta b_{m,2}) - \bar{\lambda}(\bar{b}_{m,1} + \theta \bar{b}_{m,2})] K_{j,k}(b_m, a_j) + O\left(\frac{1}{|\lambda|^2}\right), \end{aligned}$$

where

$$\begin{aligned} K_{j,k}(b_m, a_j) &= \\ &= \frac{|\frac{\partial P}{\partial z_1}(b_m)|^3 \hat{R}_\theta(\delta(b_m, a_j)) \wedge \bar{\omega}_k(b_m) (1 + |\theta|^2)}{|\frac{\partial^2 P}{\partial z_1^2} (\frac{\partial P}{\partial z_2})^2 - 2 \frac{\partial^2 P}{\partial z_1 \partial z_2} \frac{\partial P}{\partial z_2} \frac{\partial P}{\partial z_1} + \frac{\partial^2 P}{\partial z_2^2} (\frac{\partial P}{\partial z_1})^2 |dd^c |z|^2|_V(b_m)}. \end{aligned} \quad (7.1)$$

Lemma 7.1. *Let $V \setminus V_0 = \cup_{l=1}^g V_l$ be a curve with properties i)-iv) of section 1. Then $\forall \theta \neq \theta_1, \dots, \theta_d$ any point w , where $(dz_1 + \theta dz_2)|_V(w) = 0$, belongs to $V_0 = \{z \in V : |z_1| \leq r_0(\delta)\}$, where $r_0(\delta) = \text{const}(V)/\sqrt{\delta}$, $\delta = \min_l |\theta - \theta_l|$.*

Proof of Lemma 7.1. For any point $w \in V \setminus V_0$, where $(dz_1 + \theta dz_2)|_V(w) = 0$, definition $\theta_l = -\frac{1}{\gamma_l}$, $l = 1, \dots, d$, and property iii) of Section 1 imply for some $l = l(w)$ equality

$$\begin{aligned} 0 &= (dz_1 + \theta dz_2)|_V(w) = \left[1 + \theta \left(\gamma_l + \frac{\gamma_l^0}{w_1^2} + O\left(\frac{1}{w_1^3}\right)\right)\right] dz_1 = \\ &= \left[1 + \theta \gamma_l + O\left(\frac{\theta}{w_1^2}\right)\right] dz_1 = \gamma_l \left[(\theta - \theta_l) + O\left(\frac{\theta}{w_1^2}\right)\right] dz_1. \end{aligned}$$

This gives equality $\theta \left(1 + O\left(\frac{1}{w_1^2}\right)\right) = \theta_l$. This equality together with inequality $|\theta - \theta_l| \geq \delta$ implies inequality $|w_1| \leq \frac{\text{const}(V)}{\sqrt{\delta}} = r_0(\delta)$.

Lemma 7.1 is proved.

Let further

$$A_{\varepsilon,j} = \{z \in V : |z - a_j| \leq \varepsilon\}, \quad A_\varepsilon = \cup_{j=1}^g A_{\varepsilon,j},$$

$$B_{\varepsilon,m} = \{z \in V : |z - b_m| \leq \varepsilon\}, \quad B_\varepsilon = \cup_{m=1}^M B_{\varepsilon,m}.$$

Lemma 7.2 *Let $r_0(\delta)$, $\delta = \delta(\theta)$ be as in Lemma 7.1. Let χ^{A_ε} , χ^{B_ε} be smooth functions with properties*

$$\chi^{A_\varepsilon}|_{A_\varepsilon} = 1, \quad \chi^{A_\varepsilon}|_{V \setminus A_{2\varepsilon}} = 0, \quad |d\chi^{A_\varepsilon}| = O\left(\frac{1}{\varepsilon}\right),$$

$$\chi^{B_\varepsilon}|_{B_\varepsilon} = 1, \quad \chi^{B_\varepsilon}|_{V \setminus B_{2\varepsilon}} = 0, \quad |d\chi^{B_\varepsilon}| = O\left(\frac{1}{\varepsilon}\right).$$

Then for any $\varepsilon > 0$ small enough we have $B_{2\varepsilon} \cap A_{2\varepsilon} = \{\emptyset\}$ and $\forall j, k = 1, \dots, g$

$$\Delta_{\theta,\varepsilon}^{j,k}(\lambda) \stackrel{\text{def}}{=} \int_{\xi \in V} (1 - \chi^{A_\varepsilon} - \chi^{B_\varepsilon}) \hat{R}(\delta(\xi, a_j)) \wedge \bar{\omega}_k(\xi) e_{\lambda,\theta}(\xi) = O\left(\frac{1}{\lambda^2}\right).$$

Proof of Lemma 7.2. By Lemma 7.1, any point b_m , where $(dz_1 + \theta dz_2)|_V(b_m) = 0$ belongs to $\{z \in V : |z_1| \leq r_0\}$. Under the conditions of Lemma 7.2, any a_j from $\{a_1, \dots, a_g\}$ belongs to $\{z \in V : |z_1| > r_0(\delta)\}$, $\delta = \delta(\theta)$.

Then $B_{2\varepsilon} \cap A_{2\varepsilon} = \{\emptyset\}$ if ε is small enough. From definition of $\Delta_{\theta,\varepsilon}^{j,k}$ and equality $\bar{\partial} \hat{R}_\theta(\delta(\varepsilon, a_j))|_{V \setminus \{a_j\}} = 0$ we obtain

$$\begin{aligned} \Delta_{\theta,\varepsilon}^{j,k}(\lambda) &= \frac{1}{\lambda} \int_V (1 - \chi^{A_\varepsilon} - \chi^{B_\varepsilon}) \hat{R}_\theta(\delta(\xi, a_j)) \wedge \frac{\bar{\omega}_k}{d\bar{\xi}_1 + \theta d\bar{\xi}_2} \bar{\partial} e_{\lambda,\theta}(\xi) = \\ &- \frac{1}{\lambda} \int_V (1 - \chi^{A_\varepsilon} - \chi^{B_\varepsilon}) \hat{R}_\theta(\delta(\xi, a_j)) \wedge \bar{\partial} \left(\frac{\bar{\omega}_k}{d\bar{\xi}_1 + \theta d\bar{\xi}_2} \right) e_{\lambda,\theta}(\xi) - \\ &\frac{1}{\lambda} \int_V \bar{\partial} (\chi^{A_\varepsilon} + \chi^{B_\varepsilon}) \hat{R}_\theta(\delta(\xi, a_j)) \wedge \frac{\bar{\omega}_k(\xi)}{d\bar{\xi}_1 + \theta d\bar{\xi}_2} e_{\lambda,\theta}(\xi) + \\ &\frac{1}{\lambda} \lim_{r \rightarrow \infty} \int_{\{\xi \in V : |\xi_1| = r\}} \hat{R}_\theta(\delta(\xi, a_j)) \wedge \frac{\bar{\omega}_k}{d\bar{\xi}_1 + \theta d\bar{\xi}_2} e_{\lambda,\theta}(\xi). \end{aligned} \tag{7.2}$$

From asymptotic estimates $|\hat{R}_\theta(\delta(\xi, a_j))| = O(|d\xi_1|)$ and $|\bar{\omega}_k| = O\left(\frac{d\bar{\xi}_1}{\xi_1^2}\right)$, $\xi_1 \rightarrow \infty$, and property $\inf_l |\theta - \theta_l| > 0$ we obtain vanishing of the last term of the right-hand side of (7.2).

Property $(d\xi_1 + \theta d\xi_2)|_{V \setminus B_\varepsilon} \neq 0$ permits to integrate other terms of the right-hand side of (7.2) by parts once more and to obtain statement of Lemma 7.2.

Lemma 7.3 For any $k, j \in \{1, \dots, g\}$, $\theta \notin \{\theta_1, \dots, \theta_d\}$ and any $\varepsilon > 0$ we have the asymptotic equality

$$\int_V \chi^{A_{\varepsilon, j}} \hat{R}_\theta(\delta(\xi, a_j)) \wedge \bar{\omega}_k(\xi) e_{\lambda, \theta}(\xi) = -\frac{1}{\lambda} e_{\lambda, \theta}(a_j) \frac{\bar{\omega}_k}{d\xi_1 + \theta d\xi_2}(a_j) + \left(\frac{1}{\lambda^2}\right).$$

Proof of Lemma 7.3. Integration by parts of the left-hand side, equality $\bar{\partial} \hat{R}(\delta(\xi, a_j)) = \delta(\xi, a_j)$ and inequality $(d\xi_1 + \theta d\xi_2)|_{A_{\varepsilon, j}} \neq 0$ imply statement of Lemma 7.3.

Lemma 7.4 Under the conditions of Lemmas 7.1, 7.2, $\forall \delta > 0$, $\theta : \inf_l |\theta - \theta_l| > \delta$, $\forall j, k = 1, \dots, g$,

$$\begin{aligned} & \int_V \chi^{B_\varepsilon} \hat{R}_\theta(\delta(\xi, a_j)) \wedge \bar{\omega}_k(\xi) e_{i\tau, \theta}(\xi) = \\ & - \frac{\pi}{|\lambda|} \sum_{m=1}^M \exp[\lambda(b_{m,1} + \theta b_{m,2}) - \bar{\lambda}(\bar{b}_{m,1} + \bar{\theta} \bar{b}_{m,2})] K_{j,k}(b_m, a_j) + O\left(\frac{1}{|\lambda|^2}\right), \end{aligned}$$

where $\theta = \theta(b_m)$, $m = 1, \dots, M$, and $K_{j,k}(b_m, a_j)$ are defined by (7.1).

Proof of Lemma 7.4. This statement is consequence of the classical result of the stationary phase method [Fe], applied to the left-hand side, taking into account the following equality for $e_{\lambda, \theta}(z)$ in the neighborhood of the stationary points $b_m \in V$, $m = 1, \dots, M$,

$$e_{\lambda, \theta}(z) = \exp[\lambda(b_{m,1} + \theta b_{m,2}) - \bar{\lambda}(\bar{b}_{m,1} + \bar{\theta} \bar{b}_{m,2})] \times \exp[\lambda A(z_2 - b_{m,2})^2 - \bar{\lambda} \bar{A}(\bar{z}_2 - \bar{b}_{m,2})^2],$$

where

$$A = - \frac{\left(\frac{\partial^2 P}{\partial z_1^2} \theta^2 - 2 \frac{\partial^2 P}{\partial z_1 \partial z_2} \theta + \frac{\partial^2 P}{\partial z_2^2}\right)(b_m)(z_2 - b_{m,2})^2 (1 + O(z_2 - b_{m,2}))}{2 \left(\frac{\partial P}{\partial z_1}\right)(b_m)}.$$

We use here z_2, \bar{z}_2 as coordinates of integration.

Lemma 7.4 is proved.

Proof of Proposition 7.1. Proposition 7.1 follows from Lemmas 7.2-7.4.

In the proof of Proposition 1.1 we will apply also the following statement about exponential polynomials discovered by L.Ehrenpreis [E] and reinforced by C.Berenstein and M.Dostal [BD].

Proposition 7.2. ([E], [BD]) Let $Q(\xi)$ be an exponential polynomial

$$Q(\xi) = \sum_{k=1}^N q_k(\xi) e^{\langle \alpha_k, \xi \rangle},$$

where $\{q_k\}$ are polynomials of $\xi = (\xi_1, \dots, \xi_n) \in \mathbb{C}^n$, $\alpha_k = \{\alpha_{k,1}, \dots, \alpha_{k,n}\} \in \mathbb{C}^n$, $k = 1, \dots, N$.

Let $h(\xi) = \max_k \operatorname{Re} \langle \alpha_k, \xi \rangle$. Then $\forall \varepsilon > 0 \exists$ constant $C = C(\varepsilon, Q) > 0$ such that

$$|Q(\xi)|_\varepsilon \stackrel{\text{def}}{=} \sup_{\{\xi' \in \mathbb{C}: |\xi' - \xi| < \varepsilon\}} |Q(\xi')| \geq \frac{1}{C} e^{h(\xi)}.$$

The final part of the proof of Proposition 1.1 consists of the following.

Proposition 7.1 and definition of $\Delta_\theta(\lambda)$ imply asymptotic equality

$$\begin{aligned} |\lambda|^g \Delta_\theta(\lambda) &= \det \left(-\frac{\lambda}{\bar{\lambda}} e^{\lambda, \theta} \frac{\bar{\omega}_k}{d\bar{\xi}_1 + \theta d\xi_2} (a_j) - \right. \\ &\left. \pi \sum_{m=1}^M \exp[\lambda(b_{m,1} + \theta b_{m,2}) - \bar{\lambda}(\bar{b}_{m,1} + \bar{\theta} \bar{b}_{m,2})] K_{j,k}(b_m, a_j) \right) + O\left(\frac{1}{|\lambda|}\right), \end{aligned} \quad (7.3)$$

where $j, k = 1, \dots, g$.

The determinant of the right-hand side of (7.3) is an exponential polynomial $Q(\lambda, \bar{\lambda})$ of the form

$$Q(\lambda, \bar{\lambda}) = \sum_{k=1}^N q_k(\lambda, \bar{\lambda}) e^{\lambda \alpha_k - \bar{\lambda} \bar{\alpha}_k}, \quad (7.4)$$

where $\lambda \in \mathbb{C}$, $\alpha_k \in \mathbb{C}$, $k = 1, \dots, N$. Coefficient $q_k(\lambda, \bar{\lambda})$ of exponential polynomial $Q(\lambda, \bar{\lambda})$ and complex frequencies $\{\alpha_k\}$ depend on V , $\{a_j\}$, θ , $\{b_m\}$. Applying Proposition 7.2 to the exponential polynomial (7.4) we obtain uniformly for $\lambda \in \mathbb{C}$ estimate

$$|Q(\lambda, \bar{\lambda})|_\varepsilon \geq \frac{1}{C(\varepsilon, Q)} e^{\max_k \operatorname{Re}(\lambda \alpha_k - \bar{\lambda} \bar{\alpha}_k)} = \frac{1}{C(\varepsilon, Q)}. \quad (7.5)$$

The both inequalities of Proposition 1.1 follow from (7.3)-(7.5).

References

- [BC1] Beals R., Coifmann R., Multidimensional inverse scattering and nonlinear partial differential equations, Proc. Symp.Pure Math. **43** (1985), A.M.S. Providence, Rhode Island, 45-70
- [BC2] Beals R., Coifmann R., The spectral problem for the Davey- Stewartson and Ishomori hierarchies, In: "Nonlinear Evolution Equations: Integrability and Spectral methods", Proc.Workshop, Como, Italy 1988, Proc. Nonlinear Sci., 15-23 (1990)
- [BD] Berenstein C., Dostal M., Some remarks on convolution equations, Annales de l'Institut Fourier, **23**, 55-73 (1973)
- [BLMP] Boiti M., Leon J., Manna M., Pempinelli F., On a spectral transform of a KDV-like equation related to the Schrödinger operator in the plane, Inverse problems **3**, 25-36 (1987)
- [Bu] Bukhgeim A.L., Recovering a potential from the Cauchy data in the two-dimensional case, J.Inv.Ill-posed Problems, **16**, (2008)
- [C] Calderon A.P., On an inverse boundary problem. In: Seminar on Numerical Analysis and its Applications to Continuum Physics, Soc. Brasileira de Matematica, Rio de Janeiro, pp. 61-73 (1980)

- [D] Druskin V.L., The unique solution of the inverse problem in electrical surveying and electrical well logging for piecewise-constant conductivity, *Physics of the Solid Earth* **18**(1), 51-53 (1982)
- [DKN] Dubrovin B.A., Krichever I.M., Novikov S.P., The Schrödinger equation in a periodic field and Riemann surfaces, *Dokl.Akad.Nauk SSSR* **229**, 15-18 (1976) (in Russian), *Sov.Math.Dokl.*, **17**, 947-951, (1976)
- [E] Ehrenpreis L., Solutions of some problems of division II, *Amer. J. Math.* **77**, 286-292 (1955)
- [F1] Faddeev L.D., Increasing solutions of the Schrödinger equation, *Dokl.Akad.Nauk SSSR*, **165**, 514-517 (1965) (in Russian), *Sov.Phys.Dokl.* **10**, 1033-1035 (1966)
- [F2] Faddeev L.D., The inverse problem in the quantum theory of scattering II, *Curr.Probl. Math.* **3**, 93-180 (1974) (in Russian), *J.Sov.Math.* **5**, 334-396 (1976)
- [Fe] Fedorjuk M.V., *Asymptotic: integrals and series*, M.Nauka (1987) (in Russian)
- [Ga] Garsia A.M., An imbedding of closed Riemann surfaces in euclidean space, *Comm. Math.Helv.* **35**, 93-110 (1961)
- [Ge] Gelfand I.M., Some problems of functional analysis and algebra, In: *Proc.Int.Congr. Math.*, Amsterdam, pp.253-276 (1954)
- [GH] Griffiths Ph., Harris J., *Principles of algebraic geometry*, John Wiley, 1978
- [GN] Grinevich P.G., Novikov S.P., Two-dimensional inverse scattering problem for negative energies and generalized analytic functions, *Funktsional Anal i Prilozhen.* **22**(1), 23-33 (1988)
- [GT] Guillarmou C., Tzou L., Calderon inverse problem for the Schrödinger operator on Riemann surfaces, *arXiv:0904.3804* (2009) v.1
- [Ha] Hartshorne R., *Algebraic geometry*, Springer-Verlag, (1977)
- [He] Henkin G.M., Cauchy-Pompeiu type formulas for $\bar{\partial}$ on affine algebraic Riemann surfaces and some applications, *arXiv:0804.3761*, (2008) v.1 (2010) v.2
- [HP] Henkin G.M., Polyakov P.L., Homotopy formulas for the $\bar{\partial}$ -operator on $\mathbb{C}P^n$ and the Radon-Penrose transform, *Math.USSR Izvestiya* **28**, 555-587 (1987)
- [HM1] Henkin G.M., Michel V., On the explicit reconstruction of a Riemann surface from its Dirichlet-to-Neumann operator, *GAFA, Geom.Funct.Anal.* **17**, 116-155 (2007)
- [HM2] Henkin G.M., Michel V., Inverse conductivity problem on Riemann surfaces, *J.Geom. Anal.* **18**, 1033-1052 (2008)
- [Ho] Hodge W., *The theory and applications of harmonic integrals*, Cambridge Univ.Press, 1952
- [Hö] Hörmander L., *The analysis of linear partial differential operators I*, Springer 1990
- [KV] Kohn R., Vogelius M., Determining conductivity by boundary measurements II, *Comm.Pure Appl.Math.* **38**, 644-667 (1985)
- [Na] Nachman A., Global uniqueness for a two-dimensional inverse boundary problem, *Ann. of Math.* **143**, 71-96 (1996)
- [N1] Novikov R., Reconstruction of a two-dimensional Schrödinger operator from the scattering amplitude at fixed energy, *Funktsional Anal i Prilozhen.* **20**(3), 90-91 (1986) (in Russian)
- [N2] Novikov R., Multidimensional inverse spectral problem for the equation $-\Delta\psi+(v(x)-Eu(x))\psi=0$, *Funktsional Anal i Prilozhen.* **22**(4), 11-22 (1988) (in Russian)

- [N3] Novikov R., The inverse scattering problem on a fixed energy level for the two-dimensional Schrödinger operator, *J.Funct.Anal.***103** (2), 409-463 (1992)
- [NV] Novikov S.P., Veselov A.P., Two-dimensional Schrödinger operators in periodic fields, *Current Problems in Math.* **23**, 3-32 (1983) (in Russian)
- [Ro] Rodin Y., Generalized analytic functions on Riemann surfaces, *Lecture Notes Math.*, **1288**, Springer (1987)
- [Ru] Ruedy R.A., Embeddings of open Riemann surfaces, *Comm.Math. Helv.* **46**, 214-225 (1971)
- [SU] Sylvester I., Uhlmann G., A uniqueness theorem for an inverse boundary value problem in electrical prospection, *Comm.Pure Appl.Math.* **39**, 91-112 (1986)
- [Ts] Tsai T.Y., The Schrödinger operator in the plane, *Inverse Problems* **9**, 763-787 (1993)
- [V] Vekua I.N., Generalized analytic functions, Pergamon, (1962)