

AN EXTENSION OF PONTRYAGIN'S PRINCIPLE FOR STATE-CONSTRAINED OPTIMAL CONTROL OF SEMILINEAR ELLIPTIC EQUATIONS AND VARIATIONAL INEQUALITIES*

FRÉDÉRIC BONNANS[†] AND EDUARDO CASAS[‡]

Abstract. This paper deals with state-constrained optimal control problems governed by semilinear elliptic equations or variational inequalities. By using Ekeland's principle, we derive a minimum principle of Pontryagin's type under some stability conditions of the optimal cost with respect to the state constraints.

Key words. Pontryagin's principle, boundary control, semilinear elliptic operators, optimality conditions, state constraints

AMS subject classifications. 49K20, 35J65, 35J85

1. Introduction. There exists a vast literature devoted to Pontryagin's principle for optimal control problems governed by ordinary differential equations or evolution partial differential equations, but very few papers have considered the case of elliptic equations. A simple case corresponding to a linear equation was studied by Lions [19]. More recently, the authors have derived Pontryagin's principle for semilinear monotone elliptic equations in [7]. Here we extend the results of the last work by letting the existence of pointwise state constraints generalizing some preliminary results of Bonnans [3], [4]; in [8] we have considered, assuming continuity of the data, the case of boundary as well as distributed control and we obtain a "symmetric" formulation of the optimality system involving boundary and interior hamiltonians. See Bonnans and Casas [5] for a different approach to the optimality conditions of state-constrained control problems.

The difficulty of deriving the optimality conditions for control problems associated with variational inequalities is well known; see the works of Mignot [20], Mignot and Puel [21], Barbu [2]. Zheng-Xu He [18] obtained the optimality conditions for state-constrained problems governed by variational inequalities and Bonnans and Tiba [9] proved Pontryagin's principle for control problems of semilinear elliptic variational inequalities. Here we will derive a principle of Pontryagin's type for state-constrained control problems of semilinear elliptic variational inequalities.

In this article we prove Pontryagin's principle as follows: with the aid of Ekeland's principle, we introduce a family of control problems without state constraints for which some approximate solutions converge towards the optimal control of the initial problem; we derive the optimality conditions for the problems of this family by using some results on problems without state constraints that generalize those of Bonnans and Casas [7] and Bonnans and Tiba [9] and finally we pass to the limit. In order to apply Ekeland's principle we need assume some stability conditions of the optimal cost with respect to small perturbations of the feasible state set. We distinguish two different stability conditions, called weak and strong respectively. Under a weak stability condition we derive the optimality conditions in a non qualified form, while the strong stability allows us to prove a qualified Pontryagin's principle. The

*The research of the second author was partially supported by Dirección General de Investigación Científica y Técnica (Madrid)

[†]INRIA, Domaine de Voluceau, BP 105, Rocquencourt, 78153 Le Chesnay, France

[‡]Departamento de Matemática Aplicada y Ciencias de la Computación, E.T.S.I. de Caminos, C. y P., Universidad de Cantabria, 39071 Santander, Spain.

weak stability condition has been used by Casas [13] to prove the convergence of the numerical approximations of state-constrained control problems.

The paper is organized as follows: in the next section we formulate the control problem associated to a monotone semilinear elliptic equation and in Section 3 the statements of the weak and strong Pontryagin's principles are presented; in Section 4 we give some technical results used in sections 5 and 6 to prove the theorems stated in the third section; finally Section 7 is devoted to the control of variational inequalities.

2. Setting of the problem. Let Ω be an open and bounded subset of R^n , $n \geq 1$, with a Lipschitz boundary Γ . Given a non empty bounded set $K \subset R^m$, $m \geq 1$, and $f : \Omega \times R \times K \longrightarrow R$ we consider the following boundary value problem

$$(2.1) \quad \begin{cases} Ay = f(x, y(x), u(x)) & \text{in } \Omega \\ y = 0 & \text{on } \Gamma, \end{cases}$$

where

$$(2.2) \quad \begin{cases} Ay = - \sum_{i,j=1}^n \partial_{x_j} (a_{ij}(x) \partial_{x_i} y(x)), \\ a_{ij} \in C^{0,1}(\overline{\Omega}) \text{ and} \\ \exists \Lambda > 0 \text{ such that } \sum_{i,j=1}^n a_{ij}(x) \xi_i \xi_j \geq \Lambda |\xi|^2 \quad \forall \xi \in R^n, \forall x \in \Omega. \end{cases}$$

We recall that $C^{0,\alpha}(\overline{\Omega})$, with $\alpha \in (0, 1]$, is the space of all continue functions in $\overline{\Omega}$ that satisfy the Hölder condition:

$$\sup_{x_1, x_2 \in \overline{\Omega}} \frac{|u(x_2) - u(x_1)|}{|x_2 - x_1|^\alpha} < +\infty.$$

Given two measurable functions $L : \Omega \times R \times K \longrightarrow R$ and $g : \overline{\Omega} \times R \longrightarrow R$, for every $\delta > 0$ we formulate the control problem

$$(P_\delta) \quad \begin{cases} \min J(y, u) = \int_{\Omega} L(x, y(x), u(x)) dx \\ (y, u) \text{ satisfies (2.1), } u(x) \in K \text{ a.e. } x \in \Omega \text{ and } g(x, y(x)) \leq \delta \quad \forall x \in \Omega. \end{cases}$$

We will make the following assumptions on the functions defining the problem (P_δ) : $g \in C(\overline{\Omega} \times R)$, g , L and f are continuously differentiable with respect to the second variable for every $(x, u) \in \Omega \times K$ and there exist functions $M_1 \in L^s(\Omega)$, $s > n/2$ and $s \geq 2$, $M_2 \in L^1(\Omega)$ and η increasing monotone verifying for every $(x, y, u) \in \Omega \times R \times K$

$$(2.3) \quad \begin{cases} |f(x, 0, u)| + \left| \frac{\partial f}{\partial y}(x, y, u) \right| \leq M_1(x) + \eta(|y|), \\ |L(x, 0, u)| + \left| \frac{\partial L}{\partial y}(x, y, u) \right| \leq M_2(x) + \eta(|y|), \\ \frac{\partial g}{\partial y} \in C(\overline{\Omega} \times R), \quad \frac{\partial f}{\partial y}(x, y, u) \leq 0. \end{cases}$$

We will say that a control $u : \Omega \rightarrow R^m$ is feasible if $u(x) \in K$ a.e. $x \in \Omega$ and the mapping $(x, y) \rightarrow (f(x, y, u(x)), L(x, y, u(x)))$ is measurable in $\Omega \times R$. The set of feasible controls is denoted by \mathcal{K} . In this set we define the distance, called Ekeland's distance,

$$d(u, v) = m(\{x \in \Omega : u(x) \neq v(x)\}),$$

where m denotes the Lebesgue measure. Adapting the proof of Ekeland [16] to our case it is easy to check that (\mathcal{K}, d) is a complete metric space (the only difference is that we have to check the feasibility of the limit of a Cauchy sequence, which is immediate from the definition and the fact that a limit of measurable functions is measurable).

Under the previous hypotheses and thanks to the boundedness of K , we can deduce the following theorem

THEOREM 2.1. *There exist constants $C_1 > 0$ and $\alpha \in (0, 1)$ such that for every $u \in \mathcal{K}$ the equation (2.1) has a unique solution $y_u \in H_0^1(\Omega) \cap C^{0,\alpha}(\overline{\Omega})$ satisfying*

$$(2.4) \quad \|y_u\|_{H_0^1(\Omega)} + \|y_u\|_{C^{0,\alpha}(\overline{\Omega})} \leq C_1.$$

Furthermore the mapping $u \in (\mathcal{K}, d) \rightarrow y_u \in H_0^1(\Omega) \cap C^{0,\alpha}(\overline{\Omega})$ is continuous.

Before proving this theorem we state the following lemma

LEMMA 2.2. *There exist $\alpha \in (0, 1)$, C_2 and C_3 such that for every $a, b \in L^s(\Omega)$, $a(x) \geq 0$, the problem*

$$(2.5) \quad \begin{cases} Ay + ay = b & \text{in } \Omega, \\ y = 0 & \text{on } \Gamma, \end{cases}$$

has a unique solution $y \in H_0^1(\Omega) \cap C^{0,\alpha}(\overline{\Omega})$ verifying

$$(2.6) \quad \|y\|_{H_0^1(\Omega)} + \|y\|_{L^\infty(\Omega)} \leq C_2 \|b\|_{L^s(\Omega)},$$

$$(2.7) \quad \|y\|_{C^{0,\alpha}(\overline{\Omega})} \leq C_3 \|b\|_{L^s(\Omega)} (1 + \|a\|_{L^s(\Omega)}).$$

This result follows from classical estimations in the spaces $C^{0,\alpha}(\overline{\Omega})$ (Gilbarg and Trudinger [17] or Stampacchia [22]); see Bonnans and Casas [7] for details. Now we prove Theorem 2.1.

Proof. The first part of the theorem is also proved in Bonnans and Casas [7]. Let us prove the continuity of $u \rightarrow y_u$. Let $\{u_k\}_{k=1}^\infty \subset \mathcal{K}$ be a sequence converging to $u \in \mathcal{K}$, i.e. $d(u_k, u) \rightarrow 0$. Denote by y_k and y the states corresponding to u_k and u respectively. From (2.3) and (2.4) we deduce the existence of $M > 0$ such that

$$b_k(x) = f(x, y_k(x), u(x)) - f(x, y_k(x), u_k(x))$$

satisfies

$$\|b_k\|_{L^s(\Omega)} =$$

$$\left(\int_{\{x : u_k(x) \neq u(x)\}} |f(x, y_k(x), u(x)) - f(x, y_k(x), u_k(x))|^s dx \right)^{1/s} \rightarrow 0.$$

Now applying the mean value theorem we get for some function $\theta_k : \overline{\Omega} \rightarrow (0, 1)$

$$A(y - y_k) - \frac{\partial f}{\partial y}(x, y + \theta_k(y - y_k), u(x))(y - y_k) = b_k.$$

Therefore, from Lemma 2.2 and using (2.3) we obtain

$$\|y - y_k\|_{H_0^1(\Omega)} + \|y - y_k\|_{C^{0,\alpha}(\overline{\Omega})} \leq M' \|b_k\|_{L^s(\Omega)} \rightarrow 0,$$

which completes the proof. \square

REMARK 2.3. *In order to use Lemma 2.2 in the above proof we have to check that*

$$x \longrightarrow \frac{\partial f}{\partial y}(x, y(x) + \theta_k(x)(y(x) - y_k(x)), u(x))$$

is a measurable function. Although $\theta_k(x)$ itself might be non measurable, this is true because by definition of θ_k as

$$\frac{\partial f}{\partial y}(x, y(x) + \theta_k(x)(y(x) - y_k(x)), u(x)) = \begin{cases} \frac{f(x, y(x), u(x)) - f(x, y_k(x), u(x))}{y(x) - y_k(x)} & \text{if } y(x) \neq y_k(x), \\ \frac{\partial f}{\partial y}(x, y(x), u(x)) & \text{if } y(x) = y_k(x). \end{cases}$$

We finish this section by proving a lemma that will be used several times in this paper. First let us introduce some notation. In the sequel $M(\Omega)$ will denote the space of real regular Borel measures in Ω , which is identified with the dual space of $C_0(\Omega)$, the space formed by the real continuous functions defined in $\overline{\Omega}$ and vanishing on Γ . Let A^* denote the formal adjoint operator of A :

$$A^*y = - \sum_{i,j=1}^n \partial_{x_j} (a_{ji}(x) \partial_{x_i} y(x)).$$

LEMMA 2.4. *For every function $a \in L^s(\Omega)$, with $a(x) \geq 0$ a.e. $x \in \Omega$, and every Borel measure $\mu \in M(\Omega)$ there exists a unique solution in $W_0^{1,\sigma}(\Omega)$, for all $\sigma < n/(n-1)$, of problem*

$$\begin{cases} A^*p + ap = \mu & \text{in } \Omega, \\ y = 0 & \text{on } \Gamma. \end{cases}$$

Moreover there exists a constant $M > 0$ independent of a such that

$$(2.8) \quad \|p\|_{W_0^{1,\sigma}(\Omega)} \leq M \|\mu\|_{M(\Omega)}.$$

Proof. The existence and uniqueness in $W_0^{1,\sigma}(\Omega)$ of p solution of the above Dirichlet problem is well known; see Stampacchia [22] or Casas [12]. Let us prove (2.8). Let

t be the conjugate of σ , $1/t + 1/\sigma = 1$, thus $t > n$. For every $\psi \in W^{-1,t}(\Omega) = (W_0^{1,s}(\Omega))'$, the equation

$$\begin{cases} A\varphi + a\varphi = \psi & \text{in } \Omega, \\ \varphi = 0 & \text{on } \Gamma \end{cases}$$

has a unique solution in $H_0^1(\Omega) \cap C_0(\Omega)$ and, proceeding as in [7, Lemma 3.2], there exists $M > 0$ independent of a such that

$$\|\varphi\|_\infty \leq M\|\psi\|_{W^{-1,t}(\Omega)}.$$

Hence

$$\left| \int_\Omega p\psi dx \right| = \left| \int_\Omega p(A\varphi + a\varphi) dx \right| = \left| \int_\Omega \varphi d\mu \right| \leq$$

$$\|\varphi\|_\infty \|\mu\|_{M(\Omega)} \leq M\|\psi\|_{W^{-1,t}(\Omega)} \|\mu\|_{M(\Omega)},$$

which proves the desired inequality. \square

3. The weak and strong Pontryagin's minimum principle. In this section we present the statements of the weak and strong Pontryagin's principles. First let us introduce some notations and definitions.

DEFINITION 3.1. *We will say that problem (P_δ) is weakly stable on the right if*

$$(3.9) \quad \liminf_{\delta' \searrow \delta} (P_{\delta'}) = \inf(P_\delta),$$

and weakly stable on the left if

$$(3.10) \quad \liminf_{\delta' \nearrow \delta} (P_{\delta'}) = \inf(P_\delta).$$

(P_δ) is said strongly stable on the right (resp. left) if there exist $\epsilon > 0$ and $r > 0$ such that:

$$(3.11) \quad \inf(P_\delta) - \inf(P_{\delta'}) \leq r(\delta' - \delta) \quad \forall \delta' \in [\delta, \delta + \epsilon],$$

respectively

$$(3.12) \quad \inf(P_{\delta'}) - \inf(P_\delta) \leq r(\delta - \delta') \quad \forall \delta' \in [\delta - \epsilon, \delta].$$

If (P_δ) is weakly (resp. strongly) stable on the left and on the right, it will be called weakly (resp. strongly) stable.

Sufficient conditions for the weak stability were given by Casas [14] under additional regularity hypotheses on the functions L and f . In particular, if they are continuous with respect to the third variable, L is convex with respect to the same variable, K is convex and closed and (P_{δ_0}) has a feasible pair (y, u) , then (P_δ) is stable on the right for every $\delta > \delta_0$. In spite of these results, in general it is difficult to establish the stability of a problem, mainly the strong stability. However most of problems (P_δ) are weak and strongly stable, more precisely:

PROPOSITION 3.2. *Let us denote by δ_0 a real number such that (P_{δ_0}) has at least one feasible pair (y, u) . Then for every $\delta \geq \delta_0$, except at most a countable number of*

them (resp. a set of zero measure), the problem (P_δ) is weakly (resp. strongly) stable.

Proof. If we define $\phi : [\delta_0, +\infty) \rightarrow \mathbb{R}$ by $\phi(\delta) = \inf(P_\delta)$, then ϕ is a decreasing monotone function and therefore ϕ is continuous (resp. differentiable) at each point except at most a countable number of them (resp. a set of zero measure). Finally it is obvious that the continuity (resp. differentiability) of ϕ at δ implies the weak (resp. strong) stability of (P_δ) . \square

Given a number $\alpha \geq 0$, we define the Hamiltonian associated to (P_δ) by

$$H_\alpha(x, y, u, p) = \alpha L(x, y, u) + pf(x, y, u).$$

If $\alpha = 1$, we simply write H instead of H_1 . Now we can formulate the following theorems:

THEOREM 3.3 (Weak Pontryagin's Principle). *Let \bar{u} be a solution of (P_δ) in (\mathcal{K}, d) , with \bar{y} its associated state. If (P_δ) is weakly stable on the right, then there exist $\bar{\alpha} \geq 0$, $\bar{p} \in W_0^{1,\sigma}(\Omega)$ for every $\sigma < \frac{n}{n-1}$ and $\bar{\mu} \in M(\Omega)$ such that*

$$(3.13) \quad \bar{\alpha} + \|\bar{\mu}\|_{M(\Omega)} > 0,$$

$$(3.14) \quad \begin{cases} A^* \bar{p} = \frac{\partial f}{\partial y}(x, \bar{y}(x), \bar{u}(x)) + \bar{\alpha} \frac{\partial L}{\partial y}(x, \bar{y}(x), \bar{u}(x)) + \frac{\partial g}{\partial y}(x, \bar{y}(x)) \bar{\mu} \text{ in } \Omega, \\ \bar{p} = 0 \text{ on } \Gamma, \end{cases}$$

$$(3.15) \quad \int_{\Omega} (z(x) - g(x, \bar{y}(x))) d\bar{\mu}(x) \leq 0 \quad \forall z \in C_0(\Omega) \text{ with } z(x) \leq \delta \quad \forall x \in \Omega,$$

and for every $v \in K$

$$(3.16) \quad H_{\bar{\alpha}}(x, \bar{y}(x), \bar{u}(x), \bar{p}(x)) \leq H_{\bar{\alpha}}(x, \bar{y}(x), v, \bar{p}(x)) \quad \text{a.e. } x \in \Omega.$$

Moreover if there exists a Lebesgue measurable set $\Omega_0 \subset \Omega$, with $m(\Omega_0) = m(\Omega)$, of such a kind that one of the two following conditions is satisfied

H1) For each $y \in C_0(\Omega)$ and $\forall v \in K$ the set of Lebesgue points of the functions $x \rightarrow f(x, y(x), v)$ and $x \rightarrow L(x, y(x), v)$ contains Ω_0 ,

H2) The functions L and f are continuous with respect to the third variable for every $x \in \Omega_0$,

then

$$(3.17) \quad H_{\bar{\alpha}}(x, \bar{y}(x), \bar{u}(x), \bar{p}(x)) = \min_{v \in K} H_{\bar{\alpha}}(x, \bar{y}(x), v, \bar{p}(x)) \quad \text{a.e. } x \in \Omega.$$

THEOREM 3.4 (Strong Pontryagin's Principle). *Under the assumptions of Theorem 3.3 and assuming that (P_δ) is strongly stable on the right, then there exist $\bar{p} \in W_0^{1,\sigma}(\Omega)$ and $\bar{\mu} \in M(\Omega)$ satisfying (3.14)–(3.16), or (3.17) if the conditions **H1**) or **H2**) hold, with $\bar{\alpha} = 1$.*

A first version of these theorems (with stronger hypotheses) was given by Bonnans in [3] and [4]. Since we will use penalization techniques to prove these theorems, the stability on the right is the proper condition to obtain the desired result. However the Slater condition, which is a stability condition on the left, is the usual hypothesis to derive the optimality conditions (different of Pontryagin's principle) in a qualified form; see Bonnans and Casas [6]. Weak stability on the left also was the assumption in [13] to prove the convergence of the numerical approximations.

4. Hamiltonian formulation of the cost variation. In this section we generalize some results of [7] that we will use later. Let us denote by $h : \bar{\Omega} \times R \rightarrow R$ and $\phi : R \rightarrow R$ two functions satisfying the condition that ϕ is of class C^1 and h is continuous, differentiable with respect to the second variable and $\frac{\partial h}{\partial y} \in C(\bar{\Omega} \times R)$. Now we consider the functional

$$\hat{J}(y, u) = \int_{\Omega} L(x, y(x), u(x)) dx + \phi \left(\int_{\Omega} h(x, y(x)) dx \right).$$

We are interested in studying this type of functionals because they play an important role in the proof of Pontryagin's principle, the second term being particularized later to some penalization of state constraints. As in the previous section

$$H(x, y, u, p) = L(x, y, u) + pf(x, y, u).$$

In the first part of this section we will assume that the following regularity condition holds:

$$(4.18) \quad \left| \frac{\partial L}{\partial y}(x, y, u) \right| \leq M_3(x) + \eta(|y|) \quad \forall (x, y, u) \in \Omega \times R \times K,$$

with $M_3 \in L^s(\Omega)$.

Let $u, v \in \mathcal{K}$ be two controls and y_u and y_v the associated states. From the mean value theorem it follows that there exist the intermediate states \tilde{y} , \hat{y} and \tilde{y} satisfying

$$\begin{aligned} \phi \left(\int_{\Omega} h(x, y_v(x)) dx \right) &= \phi \left(\int_{\Omega} h(x, y_u(x)) dx \right) + \\ \phi' \left(\int_{\Omega} h(x, \tilde{y}(x)) dx \right) &\int_{\Omega} \frac{\partial h}{\partial y}(x, \tilde{y}(x))(y_v(x) - y_u(x)) dx, \end{aligned}$$

$$f(\cdot, y_v, v) = f(\cdot, y_u, v) + \frac{\partial f}{\partial y}(\cdot, \hat{y}, v)(y_v - y_u),$$

$$L(\cdot, y_v, v) = L(\cdot, y_u, v) + \frac{\partial L}{\partial y}(\cdot, \tilde{y}, v)(y_v - y_u),$$

with $\tilde{y}(x), \hat{y}(x), \tilde{y}(x) \in [y_u(x), y_v(x)] \forall x \in \bar{\Omega}$. Since y_u and y_v are bounded, it follows that \tilde{y}, \hat{y} and \tilde{y} also are bounded. Now we define the intermediate adjoint state $p_{u,v}$ as solution of

$$(4.19) \quad \begin{cases} A^* p_{u,v} = \frac{\partial f}{\partial y}(\cdot, \hat{y}, v) p_{u,v} + \frac{\partial L}{\partial y}(\cdot, \tilde{y}, v) + \phi' \left(\int_{\Omega} h(x, \tilde{y}) dx \right) \frac{\partial h}{\partial y}(x, \tilde{y}) & \text{in } \Omega \\ p_{u,v} = 0 & \text{on } \Gamma. \end{cases}$$

Note that if $u = v$, then $\tilde{y}, \hat{y} = \tilde{y} = y_u$ and $p_{u,v} = p_u$ is the adjoint state associated to u . Let us verify that (4.19) is well posed

LEMMA 4.1. *If (4.18) holds, then equation (4.19) has a unique solution $p_{u,v} \in H_0^1(\Omega) \cap C^{0,\alpha}(\bar{\Omega})$ that moreover satisfies*

$$(4.20) \quad \|p_{u,v}\|_{H_0^1(\Omega)} + \|p_{u,v}\|_{C^{0,\alpha}(\bar{\Omega})} \leq C_4 \quad \forall u, v \in \mathcal{K}.$$

Proof. It is a straightforward consequence of Lemma 2.2 and the inequalities (2.3). \square

Now we have the following Hamiltonian formulation of the cost variation.

PROPOSITION 4.2. *Assume that (4.18) is satisfied and let $u, v \in \mathcal{K}$ and $p_{u,v}$ be the intermediate adjoint state associated. Then*

$$\hat{J}(y_v, v) = \hat{J}(y_u, u) + \int_{\Omega} [H(x, y_u(x), v(x), p_{u,v}(x)) - H(x, y_u(x), u(x), p_{u,v}(x))] dx.$$

Proof. We have

$$\begin{aligned} \hat{J}(y_v, v) - \hat{J}(y_u, u) &= \int_{\Omega} [L(x, y_u(x), v(x)) - L(x, y_u(x), u(x))] dx + \\ &\int_{\Omega} [L(x, y_v(x), v(x)) - L(x, y_u(x), v(x))] dx + \\ &\phi \left(\int_{\Omega} h(x, y_v(x)) dx \right) - \phi \left(\int_{\Omega} h(x, y_u(x)) dx \right). \end{aligned}$$

From (4.19) we deduce

$$\begin{aligned} &\int_{\Omega} [L(x, y_v(x), v(x)) - L(x, y_u(x), v(x))] dx + \\ &\phi \left(\int_{\Omega} h(x, y_v(x)) dx \right) - \phi \left(\int_{\Omega} h(x, y_u(x)) dx \right) = \\ &\int_{\Omega} \frac{\partial L}{\partial y}(x, \tilde{y}(x), v(x))(y_v(x) - y_u(x)) dx + \\ &\phi' \left(\int_{\Omega} h(x, \tilde{y}(x)) dx \right) \int_{\Omega} \frac{\partial h}{\partial y}(x, \tilde{y}(x))(y_v(x) - y_u(x)) dx = \\ &\int_{\Omega} [A^* p_{u,v} - \frac{\partial f}{\partial y}(x, \hat{y}(x), v(x)) p_{u,v}](y_v - y_u) dx = \\ &\int_{\Omega} A(y_v - y_u) p_{u,v} dx - \int_{\Omega} \frac{\partial f}{\partial y}(x, \hat{y}(x), v(x)) p_{u,v} (y_v - y_u) dx = \\ &\int_{\Omega} [f(x, y_v(x), v(x)) - f(x, y_u(x), u(x))] p_{u,v} dx + \\ &\int_{\Omega} [f(x, y_u(x), v(x)) - f(x, y_v(x), v(x))] p_{u,v} dx = \end{aligned}$$

$$\int_{\Omega} [f(x, y_u(x), v(x)) - f(x, y_u(x), u(x))] p_{u,v} dx,$$

which proves the proposition. \square

PROPOSITION 4.3. *Assume that (4.18) holds and let $\{v_k\}_{k=1}^{\infty} \subset \mathcal{K}$ be a sequence converging to u in the topology defined by Ekeland's distance. Then the states and the adjoint states associated $y_k = y_{v_k}$ and $p_k = p_{u,v_k}$ converge to y_u and p_u respectively in $H_0^1(\Omega) \cap C^{0,\alpha}(\bar{\Omega})$.*

Proof. The convergence $y_k \rightarrow y_u$ follows from Theorem 2.1. The convergence of $\{p_k\}$ follows from the continuity of $v \in \mathcal{K} \rightarrow p_{u,v} \in H_0^1(\Omega) \cap C^{0,\alpha}(\bar{\Omega})$ which can be proved arguing in a similar way to the proof of Theorem 2.1. \square

Given a point $x_0 \in \Omega$, we will denote

$$\omega_k(x_0) = \{x \in \Omega : \|x - x_0\| \leq 1/k\}$$

and $m_k(x_0) = m(\omega_k(x_0))^{-1}$. We will say that a sequence $\{v_k\}$ in \mathcal{K} is a spike perturbation of $u \in \mathcal{K}$ around x_0 associated to $v \in K$ if

$$v_k(x) = \begin{cases} v & \text{if } x \in \omega_k(x_0), \\ u(x) & \text{otherwise.} \end{cases}$$

PROPOSITION 4.4. *Assume that (4.18) holds and let v_k be a spike perturbation of u around x_0 associated to $v \in K$, and let y_k be the associated state. Then for every $u \in \mathcal{K}$ there exists a set $\Omega(u, v) \subset \Omega$, with $m(\Omega(u, v)) = m(\Omega)$, such that*

$$\lim_{k \rightarrow \infty} m_k(x_0) [\hat{J}(y_k, v_k) - \hat{J}(y_u, u)] =$$

$$H(x_0, y_u(x_0), v, p_u(x_0)) - H(x_0, y_u(x_0), u(x_0), p_u(x_0)) \quad \forall x_0 \in \Omega(u, v).$$

Proof. From Proposition 4.2 we have

$$\hat{J}(y_k, v_k) - \hat{J}(y_u, u) = \int_{\omega_k(x_0)} [H(x, y_u(x), v, p_k(x)) - H(x, y_u(x), u(x), p_k(x))] dx,$$

where $p_k = p_{u,v_k}$ converges to p_u in $H_0^1(\Omega) \cap C^{0,\alpha}(\bar{\Omega})$ as stated in Proposition 4.3. Then

$$\hat{J}(y_k, v_k) - \hat{J}(y_u, u) = \int_{\omega_k(x_0)} [H(x, y_u(x), v, p_u(x)) - H(x, y_u(x), u(x), p_u(x))] dx +$$

$$\int_{\omega_k(x_0)} f(x, y_u(x), v)(p_k(x) - p_u(x)) dx + \int_{\omega_k(x_0)} f(x, y_u(x), u(x))(p_u(x) - p_k(x)) dx.$$

Let $\Omega(u, v)$ the intersection of the Lebesgue points of the following mappings

$$\begin{aligned} x &\longrightarrow f(x, y_u(x), v), \\ x &\longrightarrow f(x, y_u(x), u(x)), \\ x &\longrightarrow H(x, y_u(x), v, p_u(x)), \\ x &\longrightarrow H(x, y_u(x), u(x), p_u(x)). \end{aligned}$$

Then $m(\Omega(u, v)) = m(\Omega)$. Using now the uniform convergence of $p_k \rightarrow p_u$, it follows for every $x_0 \in \Omega(u, v)$

$$m_k(x_0) \left| \int_{\omega_k(x_0)} f(x, y_u(x), v)(p_u(x) - p_k(x)) dx \right| \leq$$

$$m_k(x_0) \int_{\omega_k(x_0)} |f(x, y_u(x), v)| dx \|p_u - p_k\|_{L^\infty(\Omega)} \rightarrow 0.$$

Analogously

$$m_k(x_0) \left| \int_{\omega_k(x_0)} f(x, y_u(x), u(x))(p_u(x) - p_k(x)) dx \right| \leq$$

$$m_k(x_0) \int_{\omega_k(x_0)} |f(x, y_u(x), u(x))| dx \|p_u - p_k\|_{L^\infty(\Omega)} \rightarrow 0.$$

Therefore we deduce

$$\lim_{k \rightarrow \infty} m_k(x_0) [\hat{J}(y_k, v_k) - \hat{J}(y_u, u)] =$$

$$\lim_{k \rightarrow \infty} m_k(x_0) \int_{\omega_k(x_0)} [H(x, y_u(x), v, p_u(x)) - H(x, y_u(x), u(x), p_u(x))] dx =$$

$$H(x_0, y_u(x_0), v, p_u(x_0)) - H(x_0, y_u(x_0), u(x_0), p_u(x_0)),$$

which concludes the proof. \square

The last proposition allows us to deduce easily Pontryagin's principle for control problems without state constraints. In fact it is enough to suppose that $\bar{u} \in \mathcal{K}$ is a stationary point to derive a minimum principle.

DEFINITION 4.5. *We say that \bar{u} is a stationary point of the control problem*

$$(P) \begin{cases} \min \hat{J}(y, u) \\ (y, u) \text{ satisfies (2.1) and } u(x) \in K \text{ a.e. } x \in \Omega \end{cases}$$

if

$$\liminf_{d(u, \bar{u}) \rightarrow 0} \frac{\hat{J}(y, u) - \hat{J}(\bar{y}, \bar{u})}{d(u, \bar{u})} \geq 0.$$

Obviously, every local solution in (\mathcal{K}, d) is a stationary point. Now we can prove the following proposition

PROPOSITION 4.6. *Let us suppose that (4.18) holds and let \bar{u} be a stationary point of (P). Then for every $v \in K$*

$$H(x, \bar{y}(x), \bar{u}(x), \bar{p}(x)) \leq H(x, \bar{y}(x), v, \bar{p}(x)) \quad \text{a.e. } x \in \Omega,$$

where \bar{y} and \bar{p} are the state and adjoint state associated to \bar{u} . Moreover, if condition **H1**) or **H2**) is verified, then \bar{u} satisfies Pontryagin's principle:

$$H(x, \bar{y}(x), \bar{u}(x), \bar{p}(x)) = \min_{v \in K} H(x, \bar{y}(x), v, \bar{p}(x)) \quad \text{a.e. } x \in \Omega.$$

Proof. The first part of the proof is an immediate consequence of Proposition 4.4, it is enough to remark that $d(v_k, \bar{u}) \leq m_k(x_0)^{-1}$. To derive Pontryagin's principle under the condition **H1**) we use the fact that the set $\Omega(\bar{u}, v)$, defined in the proof of Proposition 4.4, contains the intersection of Ω_0 and the set of Lebesgue points of the functions:

$$\begin{aligned} x &\longrightarrow f(x, \bar{y}(x), \bar{u}(x)), \\ x &\longrightarrow H(x, \bar{y}(x), \bar{u}(x), \bar{p}(x)). \end{aligned}$$

Indeed the continuity of \bar{p} and the condition **H1**) imply that Ω_0 is a subset of the Lebesgue point set of the functions:

$$\begin{aligned} x &\longrightarrow f(x, \bar{y}(x), v), \\ x &\longrightarrow H(x, \bar{y}(x), v, \bar{p}(x)). \end{aligned}$$

Therefore

$$H(x, \bar{y}(x), \bar{u}(x), \bar{p}(x)) \leq H(x, \bar{y}(x), v, \bar{p}(x)) \quad \forall x \in \Omega_0 \text{ and } \forall v \in K.$$

Then

$$H(x, \bar{y}(x), \bar{u}(x), \bar{p}(x)) = \min_{v \in K} H(x, \bar{y}(x), v, \bar{p}(x)) \quad \forall x \in \Omega_0.$$

In the case **H2**), let us take a sequence $\{v_k\}_{k=1}^\infty$ dense in K and $\Omega_1 = \cap_k \Omega(\bar{u}, v_k)$. Then

$$H(x, \bar{y}(x), \bar{u}(x), \bar{p}(x)) \leq H(x, \bar{y}(x), v_k, \bar{p}(x)) \quad \forall x \in \Omega_1 \text{ and } \forall k.$$

Finally the continuity of the Hamiltonian with respect to the control and the last inequality imply Pontryagin's principle in the points $x \in \Omega_0 \cap \Omega_1$. \square

We now get rid of the regularity hypothesis (4.18).

PROPOSITION 4.7. *Let \bar{u} be a local solution of (P). Then the conclusions of Proposition 4.6 remain true without hypothesis (4.18).*

To prove this proposition we will use Ekeland's principle:

THEOREM 4.8 (Ekeland [16]). *Let (E, d) be a complete metric space, $F : E \longrightarrow R \cup \{+\infty\}$ a lower semicontinuous function and let $e_\epsilon \in E$ satisfy*

$$F(e_\epsilon) \leq \inf_{e \in E} F(e) + \epsilon^2.$$

Then there exists an element $\bar{e}_\epsilon \in E$ such that

$$F(\bar{e}_\epsilon) \leq F(e_\epsilon), \quad d(\bar{e}_\epsilon, e_\epsilon) \leq \epsilon$$

and

$$F(\bar{e}_\epsilon) \leq F(e) + \epsilon d(e, \bar{e}_\epsilon) \quad \forall e \in E.$$

Now we proceed to prove Proposition 4.7.

Proof. The idea is to regularize L and to check that \bar{u} is an approximate solution of the regularized problem. Using Ekeland's principle we get some optimality conditions as in Proposition 4.7 and finally we pass to the limit in these optimality conditions and get the desired result.

The regularization is as follows. By $Proj_\epsilon$ we denote the projection onto the segment $[-1/\epsilon, +1/\epsilon]$, i.e.

$$Proj_\epsilon(t) = \max\{-1/\epsilon, \min\{t, +1/\epsilon\}\}.$$

We define

$$\varphi_\epsilon(x, t, u) = Proj_\epsilon \left(\frac{\partial L}{\partial y}(x, t, u) \right)$$

and

$$L_\epsilon(x, t, u) = L(x, 0, u) + \int_0^t \varphi_\epsilon(x, t, u) dt.$$

We can now state the problem

$$(P^\epsilon) \begin{cases} \min \hat{J}_\epsilon(y, u) = \int_\Omega L_\epsilon(x, y(x), u(x)) dx + \phi \left(\int_\Omega h(x, y(x)) dx \right) \\ (y, u) \text{ satisfies (2.1), } u(x) \in K \text{ a.e. } x \in \Omega. \end{cases}$$

We claim that $\inf(P^\epsilon) \rightarrow \inf(P)$ when $\epsilon \searrow 0$. To prove this it is enough to check that

$$(4.21) \quad |\hat{J}_\epsilon(y, u) - \hat{J}(y, u)| \leq r_\epsilon \quad \forall u \in K \text{ and } (y, u) \text{ satisfying (2.1),}$$

with $r_\epsilon \searrow 0$ when $\epsilon \searrow 0$ and r_ϵ independent of u . Indeed, if $u_\epsilon \in \mathcal{K}$ satisfies that $\hat{J}_\epsilon(y_\epsilon, u_\epsilon) \leq \inf(P_\epsilon) + \epsilon$, y_ϵ being the state associated with u_ϵ , then by (4.21)

$$\liminf_{\epsilon \searrow 0} \inf(P_\epsilon) \geq \liminf_{\epsilon \searrow 0} (\hat{J}_\epsilon(y_\epsilon, u_\epsilon) - \epsilon) \geq \liminf_{\epsilon \searrow 0} (\hat{J}_\epsilon(\bar{y}, \bar{u}) - r_\epsilon) = \inf(P)$$

and also

$$\limsup_{\epsilon \searrow 0} \inf(P_\epsilon) \leq \limsup_{\epsilon \searrow 0} \hat{J}_\epsilon(\bar{y}, \bar{u}) \leq \limsup_{\epsilon \searrow 0} [\hat{J}(\bar{y}, \bar{u}) + r_\epsilon] = \inf(P),$$

which proves that $\inf(P_\epsilon) \rightarrow \inf(P)$, as desired. Let us now check that (4.21) holds. Indeed

$$L(x, y, u) - L_\epsilon(x, y, u) = \int_0^y \left[\frac{\partial L}{\partial y}(x, t, u) - \varphi_\epsilon(x, t, u) \right] dt.$$

Let $M > 0$ be such that $|y(x)| \leq M$ whenever (y, u) is solution of (2.1) and $u \in \mathcal{K}$. Then

$$|J_\epsilon(y, u) - J(y, u)| \leq \int_\Omega \int_{-M}^{+M} \left| \varphi_\epsilon(x, t, u(x)) - \frac{\partial L}{\partial y}(x, t, u(x)) \right| dt dx =$$

$$\int_{-M}^{+M} \int_{\Omega} \left| \varphi_{\epsilon}(x, t, u(x)) - \frac{\partial L}{\partial y}(x, t, u(x)) \right| dx dt \leq$$

$$2M \int_{\Omega} \sup_{|t| \leq M, v \in K} \left| \varphi_{\epsilon}(x, t, u(x)) - \frac{\partial L}{\partial y}(x, t, u(x)) \right| dx.$$

Put

$$\Omega_{\epsilon} = \left\{ x \in \Omega : \sup_{|t| \leq M, v \in K} \left| \frac{\partial L}{\partial y}(x, t, u(x)) \right| \geq 1/\epsilon \right\}.$$

As

$$\left| \frac{\partial L}{\partial y}(x, t, u(x)) \right| \leq M_2 \in L^1(\Omega),$$

it follows that $m(\Omega_{\epsilon}) \searrow 0$ as $\epsilon \searrow 0$, and

$$\left| \hat{J}_{\epsilon}(y, u) - \hat{J}(y, u) \right| \leq r_{\epsilon} = 2M \int_{\Omega_{\epsilon}} M_2(x) dx \rightarrow 0.$$

As a consequence of the previous results, we can get a family of real numbers $\{\delta_{\epsilon}\}_{\epsilon > 0}$, with $\delta_{\epsilon} \searrow 0$ when $\epsilon \searrow 0$, such that

$$\hat{J}_{\epsilon}(\bar{y}, \bar{u}) \leq \inf(P_{\epsilon}) + \delta_{\epsilon}^2.$$

Therefore we can apply Ekeland's principle with $F(u) = \hat{J}_{\epsilon}(y_u, u)$ defined in the complete metric space (\mathcal{K}, d) , and deduce the existence of a control $u_{\epsilon} \in \mathcal{K}$ such that $d(\bar{u}, u_{\epsilon}) \leq \delta_{\epsilon}$ and

$$(4.22) \quad \hat{J}_{\epsilon}(y_{\epsilon}, u_{\epsilon}) \leq \hat{J}_{\epsilon}(y_u, u) + \delta_{\epsilon} d(u, u_{\epsilon}) \quad \forall u \in \mathcal{K},$$

where y_{ϵ} is the state associated with u_{ϵ} . To apply Proposition 4.6 we must put the cost given by the right hand side of inequality (4.22) into the framework of this proposition. For it we introduce the function $\chi_{\epsilon} : \Omega \times K \rightarrow R$ by

$$\chi_{\epsilon}(x, v) = \begin{cases} 0 & \text{if } v = u_{\epsilon}(x) \\ 1 & \text{otherwise.} \end{cases}$$

Then $(y_{\epsilon}, u_{\epsilon})$ is the solution of the problem

$$(Q^{\epsilon}) \begin{cases} \min \hat{J}_{\epsilon}(y, u) = \hat{J}_{\epsilon}(y, u) + \delta_{\epsilon} \int_{\Omega} \chi_{\epsilon}(x, u(x)) \\ (y, u) \text{ satisfies (2.1), } u(x) \in K \text{ a.e. } x \in \Omega. \end{cases}$$

Then Proposition 4.6 implies that for every $v \in K$

$$H^{\epsilon}(x, y_{\epsilon}(x), u_{\epsilon}(x), p_{\epsilon}(x)) \leq H^{\epsilon}(x, y_{\epsilon}(x), v, p_{\epsilon}(x)) + \delta_{\epsilon} \quad \text{a.e. } x \in \Omega,$$

where p_{ϵ} is the adjoint state

$$(4.23) \quad \begin{cases} A^* p_{\epsilon} = \frac{\partial f}{\partial y}(x, y_{\epsilon}, u_{\epsilon}) p_{\epsilon} + \frac{\partial L_{\epsilon}}{\partial y}(x, y_{\epsilon}, u_{\epsilon}) \text{ in } \Omega, \\ p_{\epsilon} = 0 \text{ on } \Gamma, \end{cases}$$

and

$$(4.24) \quad H^\epsilon(x, y, u, p) = L_\epsilon(x, y, u) + pf(x, y, u) + \delta_\epsilon \chi_\epsilon(x, u).$$

From Theorem 2.1 it follows that $y_\epsilon \rightarrow \bar{y}$ in $H_0^1(\Omega) \cap C^{0,\alpha}(\bar{\Omega})$. Then, thanks to hypothesis (2.3) and the definition of L_ϵ , we have

$$f(x, y_\epsilon, u_\epsilon) \rightarrow f(x, \bar{y}, \bar{u}) \quad \text{and} \quad \frac{\partial f}{\partial y}(x, y_\epsilon, u_\epsilon) \rightarrow \frac{\partial f}{\partial y}(x, \bar{y}, \bar{u}) \quad \text{in } L^s(\Omega),$$

$$L_\epsilon(x, y_\epsilon, u_\epsilon) \rightarrow L(x, \bar{y}, \bar{u}) \quad \text{and} \quad \frac{\partial L_\epsilon}{\partial y}(x, y_\epsilon, u_\epsilon) \rightarrow \frac{\partial L}{\partial y}(x, \bar{y}, \bar{u}) \quad \text{in } L^1(\Omega).$$

With the aid of these relations and Lemma 2.4 we can pass to the limit in (4.23) and (4.24) and to deduce the first conclusion of Proposition 4.6. To prove the second conclusion, i.e. the Pontryagin's principles, we argue as follows. Under condition **H2**), the argument used in the proof of Proposition 4.6 can be repeated here without modifications. If condition **H1**) is satisfied, then, thanks to Proposition 4.6, we can take a sequence $\epsilon_j \searrow 0$ and $\delta_j = \delta_{\epsilon_j} \searrow 0$ such that

$$H^{\epsilon_j}(x, y_{\epsilon_j}(x), u_{\epsilon_j}(x), p_{\epsilon_j}(x)) \leq \min_{v \in K} H^{\epsilon_j}(x, y_{\epsilon_j}(x), v, p_{\epsilon_j}(x)) + \delta_j \quad \text{a.e. } x \in \Omega_j,$$

with $m(\Omega_j) = m(\Omega)$. Now we pass to the limit and get

$$H(x, \bar{y}(x), \bar{u}(x), \bar{p}(x)) = \min_{v \in K} H(x, \bar{y}(x), v, \bar{p}(x)) \quad \text{a.e. } x \in \tilde{\Omega} = \bigcap_{j=1}^{\infty} \Omega_j,$$

which concludes the proof because $m(\tilde{\Omega}) = m(\Omega)$. \square

THEOREM 4.9. *The statement of Proposition 4.6 is still valid without hypothesis (4.18).*

Proof. From the definition of stationary point we deduce that for every $\epsilon > 0$ there exists $r_\epsilon > 0$ such that

$$\frac{\hat{J}(y, u) - \hat{J}(\bar{y}, \bar{u})}{d(u, \bar{u})} \geq -\epsilon \quad \forall u \in B_{r_\epsilon}(\bar{u}),$$

where $B_{r_\epsilon}(\bar{u})$ is the open ball of (\mathcal{K}, d) of radius r_ϵ and center at \bar{u} . Hence

$$\hat{J}(\bar{y}, \bar{u}) \leq \hat{J}(y, u) + \epsilon \int_{\Omega} \chi(x, u(x)) dx \quad \forall u \in B_{r_\epsilon}(\bar{u}),$$

with

$$\chi(x, v) = \begin{cases} 0 & \text{if } v = \bar{u}(x) \\ 1 & \text{otherwise.} \end{cases}$$

Then it is enough to apply Proposition 4.7 to the problem

$$\begin{cases} \min \hat{J}(y, u) + \epsilon \int_{\Omega} \chi(x, u(x)) dx \\ u \in B_{r_\epsilon}(\bar{u}), u(x) \in K, \end{cases}$$

and pass to the limit when $\epsilon \searrow 0$, to deduce the desired result. \square

The hypotheses made about K , L and f do not allow to assure the existence of a solution of control problem (P) . Here we will prove a principle of Pontryagin's type for ϵ -solutions.

DEFINITION 4.10. *A control $u \in \mathcal{K}$ is called an ϵ -solution of (P) if $\hat{J}(y_u, u) \leq \inf(P) + \epsilon$.*

THEOREM 4.11. *For every $\epsilon > 0$ there exists at least one ϵ^2 -solution of (P) in \mathcal{K} . Furthermore for every ϵ^2 -solution of (P) , \bar{u}_ϵ , there exists another ϵ^2 -solution u_ϵ such that $d(u_\epsilon, \bar{u}_\epsilon) \leq \epsilon$ and for every $v \in K$*

$$H(x, y_\epsilon(x), u_\epsilon(x), p_\epsilon(x)) \leq H(x, y_\epsilon(x), v, p_\epsilon(x)) + \epsilon \quad \text{a.e. } x \in \Omega,$$

where $y_\epsilon = y_{u_\epsilon}$ and $p_\epsilon = p_{u_\epsilon}$. Moreover, if there exists a Lebesgue measurable set $\Omega_0 \subset \Omega$, with $m(\Omega_0) = m(\Omega)$, in such a way that **H1**) or **H2**) holds, then \bar{u} satisfies Pontryagin's principle:

$$H(x, y_\epsilon(x), u_\epsilon(x), p_\epsilon(x)) = \min_{v \in K} H(x, y_\epsilon(x), v, p_\epsilon(x)) + \epsilon \quad \text{a.e. } x \in \Omega.$$

Proof. Thanks to the hypothesis (2.3), we have that $\inf(P) \in R$. Therefore there exists at least one ϵ^2 -solution of (P) . Let \bar{u}_ϵ be one of them. Then we can apply Theorem 4.8, with $F(u) = \hat{J}(y_u, u)$ defined in the metric space (\mathcal{K}, d) , and deduce the existence of a control $u_\epsilon \in \mathcal{K}$, ϵ^2 -solution of (P) , such that $d(u_\epsilon, \bar{u}_\epsilon) \leq \epsilon$ and

$$(4.25) \quad \hat{J}(y_{u_\epsilon}, u_\epsilon) \leq \hat{J}(y_u, u) + \epsilon d(u, u_\epsilon) \quad \forall u \in \mathcal{K}.$$

Now we put the cost into the framework of Proposition 4.7, using the function χ_ϵ as in its proof, and apply it to get the result. \square

5. Proof of the weak minimum principle. Let \bar{u} be a solution of (P_δ) and \bar{y} its associated state. For every $\gamma > 0$ we define the problems

$$(Q_\gamma) \begin{cases} \min J_\gamma(y, u) = \int_\Omega \left[L(x, y(x), u(x)) + \frac{1}{2\gamma} ((g(x, y(x)) - \delta)^+)^2 \right] dx \\ (y, u) \text{ satisfies (2.1) and } u(x) \in K \text{ a.e. } x \in \Omega. \end{cases}$$

The first issue to remark is the following

PROPOSITION 5.1. *Let (P_δ) be weakly stable on the right. Then*

$$\liminf_{\gamma \searrow 0} \inf(Q_\gamma) = \inf(P_\delta).$$

Proof. Let $\{u_\gamma\}$ be a family of γ -solutions of problems (Q_γ) and $\{y_\gamma\}$ the associated states:

$$J(y_\gamma, u_\gamma) \leq \inf(Q_\gamma) + \gamma.$$

From the definition of (Q_γ) it follows that $(g(x, y_\gamma(x)) - \delta)^+ \rightarrow 0$ in $L^2(\Omega)$, which, together with (2.4) and the compactness of the inclusion $H_0^1(\Omega) \cap C^{0,\alpha}(\bar{\Omega}) \subset C_0(\Omega)$, implies the convergence $(g(x, y_\gamma(x)) - \delta)^+ \rightarrow 0$ in $C_0(\Omega)$. Therefore

$$\delta_\gamma = \|(g(x, y_\gamma(x)) - \delta)^+\|_{L^\infty(\Omega)} + \delta \rightarrow \delta \text{ if } \gamma \searrow 0.$$

As (y_γ, u_γ) is feasible pair for (P_{δ_γ}) we deduce that

$$\inf(P_{\delta_\gamma}) \leq J(y_\gamma, u_\gamma) \leq \inf(Q_\gamma) + \gamma.$$

Then, using the weak stability of (P_δ) on the right, we obtain

$$\inf(P_\delta) = \lim_{\gamma \searrow 0} \inf(P_{\delta_\gamma}) \leq \lim_{\gamma \searrow 0} \{\inf(Q_\gamma) + \gamma\} = \lim_{\gamma \searrow 0} \inf(Q_\gamma) \leq \inf(P_\delta),$$

with the last inequality due to the fact that (y_u, u) is feasible for (Q_γ) whenever it is feasible for (P_δ) , with the same cost. \square

Proof of Theorem 3.3 Thanks to Proposition 5.1 we deduce that \bar{u} is a ϵ_γ^2 -solution of (Q_γ) , with $\epsilon_\gamma \searrow 0$ when $\gamma \searrow 0$. Applying Theorem 4.11 we obtain the existence of a control $u_\gamma \in \mathcal{K}$, ϵ_γ^2 -solution of (Q_γ) , with $d(u_\gamma, \bar{u}) \leq \epsilon_\gamma$ and such that for every $v \in K$

$$H(x, y_\gamma(x), u_\gamma(x), p_\gamma(x)) \leq H(x, y_\gamma(x), v, p_\gamma(x)) + \epsilon_\gamma \quad \text{a.e. } x \in \Omega,$$

where $y_\gamma = y_{u_\gamma}$ and p_γ is the adjoint state:

$$\begin{cases} A^* p_\gamma = \frac{\partial f}{\partial y}(x, y_\gamma, u_\gamma) p_\gamma + \frac{\partial L}{\partial y}(x, y_\gamma, u_\gamma) + \frac{1}{\gamma}(g(x, y_\gamma) - \delta)^+ \frac{\partial g}{\partial y}(x, y_\gamma) & \text{in } \Omega, \\ p_\gamma = 0 & \text{on } \Gamma. \end{cases}$$

Defining

$$\begin{aligned} \bar{\alpha}_\gamma &= \left(1 + \left\| \frac{1}{\gamma}(g(x, y_\gamma) - \delta)^+ \right\|_{M(\Omega)}\right)^{-1} \\ \bar{\mu}_\gamma &= \frac{\bar{\alpha}_\gamma}{\gamma}(g(x, y_\gamma) - \delta)^+, \quad \text{and} \quad \bar{p}_\gamma = \bar{\alpha}_\gamma p_\gamma, \end{aligned}$$

we get

$$(5.26) \quad \bar{\alpha}_\gamma + \|\bar{\mu}_\gamma\|_{M(\Omega)} = 1,$$

$$(5.27) \quad \begin{cases} A^* \bar{p}_\gamma = \frac{\partial f}{\partial y}(x, y_\gamma, u_\gamma) \bar{p}_\gamma + \bar{\alpha}_\gamma \frac{\partial L}{\partial y}(x, y_\gamma, u_\gamma) + \bar{\mu}_\gamma \frac{\partial g}{\partial y}(x, y_\gamma) & \text{in } \Omega, \\ \bar{p}_\gamma = 0 & \text{on } \Gamma, \end{cases}$$

and for every $v \in K$

$$(5.28) \quad H_{\bar{\alpha}_\gamma}(x, y_\gamma(x), u_\gamma(x), \bar{p}_\gamma(x)) \leq H_{\bar{\alpha}_\gamma}(x, y_\gamma(x), v, \bar{p}_\gamma(x)) + \epsilon_\gamma \quad \text{a.e. } x \in \Omega.$$

If $\gamma \searrow 0$, then $d(u_\gamma, \bar{u}) \leq \epsilon_\gamma \rightarrow 0$, therefore from Theorem 2.1 we get $y_\gamma \rightarrow \bar{y}$ in $H_0^1(\Omega) \cap C^{0,\alpha}(\bar{\Omega})$. Using now (2.3), we deduce the convergences

$$\begin{aligned} f(x, y_\gamma, u_\gamma) &\rightarrow f(x, \bar{y}, \bar{u}), & \frac{\partial f}{\partial y}(x, y_\gamma, u_\gamma) &\rightarrow \frac{\partial f}{\partial y}(x, \bar{y}, \bar{u}) \text{ in } L^s(\Omega), \\ L(x, y_\gamma, u_\gamma) &\rightarrow L(x, \bar{y}, \bar{u}), & \frac{\partial L}{\partial y}(x, y_\gamma, u_\gamma) &\rightarrow \frac{\partial L}{\partial y}(x, \bar{y}, \bar{u}) \text{ in } L^s(\Omega). \end{aligned}$$

Applying Lemma 2.4, we obtain the following estimation for \bar{p}_γ

$$\|\bar{p}_\gamma\|_{W_0^{1,\sigma}(\Omega)} \leq M \left\| \bar{\alpha}_\gamma \frac{\partial L}{\partial y}(x, y_\gamma, u_\gamma) + \bar{\mu}_\gamma \frac{\partial g}{\partial y}(x, y_\gamma(x)) \right\|_{M(\Omega)} \leq M' < +\infty$$

for every $\sigma < n/(n-1)$. Therefore, remembering (5.26), we can extract subsequences, denoted in the same way, such that $\bar{\mu}_\gamma \rightarrow \bar{\mu}$ in $M(\Omega)$ \ast weakly and $\bar{p}_\gamma \rightarrow \bar{p}$ weakly in $W_0^{1,\sigma}(\Omega)$. From the Rellich's theorem (Adams [1]) it follows $\bar{p}_\gamma \rightarrow \bar{p}$ strongly in $L^q(\Omega)$ for each $q < n/(n-2)$. Then

$$\frac{\partial f}{\partial y}(x, y_\gamma, u_\gamma) \bar{p}_\gamma \rightarrow \frac{\partial f}{\partial y}(x, \bar{y}, \bar{u}) \bar{p} \text{ in } L^1(\Omega).$$

We can pass to the limit in (5.27) and derive (3.14). Relation (3.16) follows from (5.28). Relation (3.15) is obtained as follows: for every $z \in C_0(\Omega)$ with $z(x) \leq \delta$ for all $x \in \Omega$

$$\int_{\Omega} (z(x) - g(x, \bar{y}(x))) d\bar{\mu}(x) = \lim_{\gamma \rightarrow 0} \frac{\bar{\alpha}_\gamma}{\gamma} \int_{\Omega} (z(x) - g(x, y_\gamma(x)))(g(x, y_\gamma(x)) - \delta)^+ dx \leq 0.$$

It follows that

$$\langle \bar{\mu}, g(\cdot, \bar{y}) \rangle = \max\{\langle \bar{\mu}, z \rangle : z \in C_0(\Omega), z(x) \leq \delta\},$$

hence $\bar{\mu}$ is nonnegative and the value of the max is $\delta \|\bar{\mu}\|_{M(\Omega)}$.

To obtain (3.13) it is enough to remember (5.26) and remark that

$$\|\bar{\mu}\|_{M(\Omega)} = \frac{1}{\delta} \langle \bar{\mu}, g(\cdot, \bar{y}) \rangle = \lim_{\gamma \rightarrow 0} \frac{1}{\delta} \langle \bar{\mu}_\gamma, g(\cdot, y_\gamma) \rangle = \lim_{\gamma \rightarrow 0} \|\bar{\mu}_\gamma\|_{M(\Omega)}.$$

Finally we must prove (3.17). If we assume that **H1**) is satisfied, then, thanks to Theorem 4.11, (5.28) can be written

$$H_{\bar{\alpha}_\gamma}(x, y_\gamma(x), u_\gamma(x), \bar{p}_\gamma(x)) = \min_{v \in K} H_{\bar{\alpha}_\gamma}(x, y_\gamma(x), v, \bar{p}_\gamma(x)) + \epsilon_\gamma \text{ a.e. } x \in \Omega.$$

Taking a subsequence $\{\gamma_j\}_{j=1}^\infty$, we pass to the limit as above and get (3.17).

If **H2**) is satisfied, we can argue as in the proof of Theorem 4.9 to conclude (3.17).

□

6. Proof of the strong minimum principle. In this section we establish the existence of a certain link between the stability of the cost with respect to small perturbations of the feasible state set and the viability of the exact penalization procedure of the state constraints. In the context of the abstract optimization Burke [11], generalizing an idea of Clarke [15], proved an equivalence result between stability and exact penalization. Since we are assuming the hypotheses of Theorem 3.4, we have that (P_δ) is strongly stable on the right and (\bar{y}, \bar{u}) is a solution of this problem. Now we consider the exact penalization of state constraints:

PROPOSITION 6.1. *If $r > 0$ satisfies (3.11), then \bar{u} is a local solution in (\mathcal{K}, d) of the penalized control problem*

$$\begin{cases} \min J_r(u) = \int_{\Omega} L(x, y_u(x), u(x)) dx + r \|(g(x, y_u) - \delta)^+\|_\infty \\ u \in \mathcal{K}. \end{cases}$$

Proof. From (3.11) it follows

$$\inf(P_\delta) = \inf \{J(y_u, u) + r(\delta' - \delta) : u \in \mathcal{K}, g(x, y_u(x)) \leq \delta', \delta' \in [\delta, \delta + \epsilon]\}.$$

Minimizing first with respect to δ' for fixed u we find

$$\inf(P_\delta) = \inf \left\{ J(y_u, u) + r \|(g(x, y_u) - \delta)^+\|_\infty : u \in \mathcal{K}, g(x, y_u(x)) \leq \delta + \epsilon \right\}.$$

Since the mapping $u \in (\mathcal{K}, d) \longrightarrow y_u \in C_0(\Omega)$ is continuous, we deduce the existence of a ball $B_\lambda(\bar{u})$, $\lambda > 0$, such that

$$\|g(x, y_u)\|_\infty < \delta + \epsilon \quad \forall u \in B_\lambda(\bar{u}),$$

which, together with the previous identity, proves that \bar{u} is a local solution of the penalized control problem. \square

Take $\lambda > 0$ as in the proof of the previous proposition and $r > 0$ verifying (3.11). We introduce the problem

$$(Q_r) \left\{ \begin{array}{l} \min J_r(u) = \int_\Omega L(x, y_u(x), u(x)) dx + r \|(g(x, y_u) - \delta)^+\|_\infty \\ u \in B_\lambda(\bar{u}). \end{array} \right.$$

Then \bar{u} is a solution of this problem. We passed from a state-constrained control problem to another control problem without state constraints. The difficulty in this new problem is that the penalization term is not differentiable. To overcome this difficulty we define

$$J_{r,q}(u) = \int_\Omega L(x, y_u(x), u(x)) dx + r \left(q^{-q} + \int_\Omega [(g(x, y_u(x)) - \delta)^+]^q dx \right)^{1/q}$$

and

$$(Q_{r,q}) \left\{ \begin{array}{l} \min J_{r,q}(u) \\ u \in B_\lambda(\bar{u}), \end{array} \right.$$

with $q > 1$. Note that $(Q_{r,q})$ has a differentiable cost and moreover it represents an approximation of (Q_r) :

PROPOSITION 6.2. *The following identity holds*

$$\inf(Q_r) = \lim_{q \rightarrow \infty} \inf(Q_{r,q}).$$

Proof. From the convergence $\|z\|_{L^q(\Omega)} \rightarrow \|z\|_\infty$ for every $z \in L^\infty(\Omega)$ and the inequalities

$$\|(g(x, y_u) - \delta)^+\|_{L^q(\Omega)} \leq \left(q^{-q} + \int_\Omega [(g(x, y_u(x)) - \delta)^+]^q dx \right)^{1/q} \leq$$

$$\frac{1}{q} + \|(g(x, y_u) - \delta)^+\|_{L^q(\Omega)}$$

we deduce that $J_{r,q}(u) \rightarrow J_r(u)$ when $q \rightarrow +\infty$. Therefore for every $u \in B_\lambda(\bar{u})$

$$\limsup_{q \rightarrow +\infty} \inf(Q_{r,q}) \leq \limsup_{q \rightarrow +\infty} J_{r,q}(u) = J_r(u),$$

hence

$$(6.29) \quad \limsup_{q \rightarrow +\infty} \inf(Q_{r,q}) \leq \inf(Q_r).$$

Now we prove the converse inequality. Let us take $\epsilon > 0$ arbitrary and let $C_1 > 0$ be the constant given in Theorem 2.1. Since $g : \bar{\Omega} \times R \rightarrow R$ is continuous, it follows the existence of a constant $\rho \in (0, \epsilon)$ such that $\forall x, x' \in \bar{\Omega}$

$$(6.30) \quad |g(x, t) - g(x', t)| < \epsilon \quad \text{if} \quad |x - x'| \leq \rho \quad \text{and} \quad |t| \leq C_1.$$

Moreover we assume ρ small enough in such a way that $m(\{x : |x| \leq \rho\}) < 1$. Now we define $\Omega_\rho(x_0) = \Omega \cap B_\rho(x_0)$. Since the boundary of Γ is Lipschitz, there exists a number $\beta \in (0, 1)$ verifying

$$m(\Omega_\rho(x_0)) \geq \beta m(\{x : |x| \leq \rho\}) \quad \forall x_0 \in \bar{\Omega}.$$

On the other hand, from the continuity of $\frac{\partial g}{\partial y}$ we deduce the existence of another constant $M > 0$ such that

$$(6.31) \quad |g(x, t)| + \left| \frac{\partial g}{\partial y}(x, t) \right| \leq M \quad \forall (x, t) \in \bar{\Omega} \times [-C_1, +C_1].$$

Pick $u \in B_\lambda(\bar{u})$. If $\|(g(x, y_u) - \delta)^+\|_\infty = 0$, then $J_{r,q}(u) = J_r(u)$. Let us suppose that $\|(g(x, y_u) - \delta)^+\|_\infty > 0$ and take $x_0 \in \bar{\Omega}$ verifying

$$(g(x_0, y_u(x_0)) - \delta)^+ = \|(g(x, y_u) - \delta)^+\|_\infty.$$

Then for each $x \in \Omega_\rho(x_0)$ we get with the aid of (2.4), (6.30) and (6.31)

$$|g(x, y_u(x)) - g(x_0, y_u(x_0))| \leq$$

$$|g(x, y_u(x)) - g(x_0, y_u(x))| + |g(x_0, y_u(x)) - g(x_0, y_u(x_0))| \leq$$

$$\epsilon + M|y_u(x) - y_u(x_0)| \leq \epsilon + MC_1\rho^\alpha \leq M'\epsilon,$$

hence

$$(g(x, y_u(x)) - \delta)^+ \geq (g(x_0, y_u(x_0)) - \delta - M'\epsilon)^+ \quad \forall x \in \Omega_\rho(x_0).$$

Therefore, we obtain

$$\begin{aligned} \|(g(x, y_u) - \delta)^+\|_{L^q(\Omega)} &\geq \left(\int_{\Omega_\rho(x_0)} [g(x, y_u(x)) - \delta]^q dx \right)^{1/q} \geq \\ m(\Omega_\rho(x_0))^{1/q} (g(x_0, y_u(x_0)) - \delta - M'\epsilon)^+ &\geq \|(g(x, y_u) - \delta)^+\|_\infty + \\ (m(\Omega_\rho(x_0))^{1/q} - 1) \|(g(x, y_u) - \delta)^+\|_\infty - M'\epsilon m(\Omega_\rho(x_0))^{1/q} &\geq \\ \|(g(x, y_u) - \delta)^+\|_\infty + M(m(\Omega_\rho(x_0))^{1/q} - 1) - M'\epsilon. \end{aligned}$$

Choosing $q_\epsilon > 1$ such that

$$1 - m(\Omega_\rho(x_0))^{1/q} \leq 1 - [\beta m(\{x : |x| \leq \rho\})]^{1/q} < \epsilon \quad \forall q > q_\epsilon,$$

it follows

$$\|(g(x, y_u) - \delta)^+\|_{L^q(\Omega)} \geq \|(g(x, y_u) - \delta)^+\|_\infty - (M + M')\epsilon \quad \forall q > q_\epsilon.$$

We have proved that

$$J_{r,q}(u) \geq J_r(u) - (M + M')\epsilon$$

for each $u \in B_\lambda(\bar{u})$ and all $q \geq q_\epsilon$, hence

$$\liminf_{q \rightarrow +\infty} \inf(Q_{r,q}) \geq \inf(Q_r) - (M + M')\epsilon$$

for $\epsilon > 0$ arbitrary, consequently

$$(6.32) \quad \liminf_{q \rightarrow +\infty} \inf(Q_{r,q}) \geq \inf(Q_r).$$

So the proposition follows from (6.29) and (6.32). \square

Proof of Theorem 3.4 Thanks to Proposition 6.2 we deduce that \bar{u} is an ϵ_q^2 -solution of $(Q_{r,q})$, with $\epsilon_q \rightarrow 0$ as $q \rightarrow \infty$. Then Theorem 4.11 states the existence of a control $u_q \in \mathcal{K}$, with $d(u_q, \bar{u}) \leq \epsilon_q$, satisfying for every $v \in K$

$$(6.33) \quad H(x, y_q(x), u_q(x), p_q(x)) \leq H(x, y_q(x), v, p_q(x)) + \epsilon_q \quad \text{a.e. } x \in \Omega,$$

where $y_q = y_{u_q}$ and p_q is the adjoint state:

$$(6.34) \quad \begin{cases} A^* p_q = \frac{\partial f}{\partial y}(x, y_q, u_q) p_q + \frac{\partial L}{\partial y}(x, y_q, u_q) + \mu_q \frac{\partial g}{\partial y}(x, y_q) & \text{in } \Omega, \\ p_q = 0 & \text{on } \Gamma, \end{cases}$$

with

$$\mu_q = r \left(q^{-q} + \int_\Omega [(g(x, y_q(x)) - \delta)^+]^q dx \right)^{1/q-1} [(g(x, y_q) - \delta)^+]^{q-1}.$$

Now we must pass to the limit. From $d(u_q, \bar{u}) \leq \epsilon_q \rightarrow 0$ and Theorem 2.1 we obtain that $y_q \rightarrow \bar{y}$ in $H_0^1(\Omega) \cap C^{0,\alpha}(\bar{\Omega})$. On the other hand, from the definition of μ_q we get

$$\|\mu_q\|_{M(\Omega)} = \|\mu_q\|_{L^1(\Omega)} \leq r \left(\int_\Omega [(g(x, y_q(x)) - \delta)^+]^q dx \right)^{1/q-1} \int_\Omega [(g(x, y_q(x)) - \delta)^+]^{q-1} dx =$$

$$r \|(g(x, y_q) - \delta)^+\|_{L^q(\Omega)}^{1-q} \|(g(x, y_q) - \delta)^+\|_{L^{q-1}(\Omega)}^{q-1}.$$

Applying Hölder's inequality with exponents $q/(q-1)$ and q it follows

$$\|z\|_{L^{q-1}(\Omega)} \leq m(\Omega)^{1/q} \|z\|_{L^q(\Omega)} \quad \forall z \in L^q(\Omega),$$

which together with the previous relation leads to

$$\|\mu_q\|_{M(\Omega)} \leq m(\Omega)^{1/q} r \|(g(x, y_q) - \delta)^+\|_{L^q(\Omega)} \leq$$

$$m(\Omega)^{2/q} r \|(g(x, y_q) - \delta)^+\|_\infty \leq M < +\infty \quad \forall q > 1.$$

As in the proof of Theorem 3.3, the boundedness of $\{\mu_q\}_{q \geq 1}$, the convergence of $\{(y_q, u_q)\}_{q \geq 1}$ and assumptions (2.3) imply the boundedness of $\{p_q\}_{q \geq 1}$ in $W_0^{1,\sigma}(\Omega)$ for every $\sigma < n/(n-1)$. Therefore we can extract subsequences $\{p_{q_k}\}$ and $\{\mu_{q_k}\}$, with $q_k \rightarrow +\infty$, converging to \bar{p} and $\bar{\mu}$ in $W_0^{1,\sigma}(\Omega)$ weakly and $M(\Omega)$ *-weakly respectively. Now it is easy to pass to the limit in (6.33) and (6.34) and to obtain (3.16) and (3.14). As in the proof of Theorem 3.3 we derive (3.15) from the definition of μ_q .

Finally, as stated in Theorem 4.11, under the conditions **H1)** or **H2)** given in Theorem 3.3, the relation (6.33) becomes

$$H(x, y_q(x), u_q(x), p_q(x)) = \min_{v \in K} H(x, y_q(x), v, p_q(x)) + \epsilon_q \quad \text{a.e. } x \in \Omega.$$

Therefore, passing to the limit in this inequality, we get (3.17). \square

7. Pontryagin's Principle in the Control of Variational Inequalities. In this section we will consider the following control system:

$$(7.35) \quad \begin{cases} Ay + \beta(y) = f(x, y(x), u(x)) & \text{in } \Omega, \\ y = 0 & \text{on } \Gamma, \end{cases}$$

where A and f are as in Section 2 and β is a maximal monotone graph in $R \times R$ (see Brezis [10] and Barbu [2]) such that $\text{dom}(\beta) \ni 0$. The control problem is

$$(P_\delta) \quad \begin{cases} \min J(y, u) = \int_{\Omega} L(x, y(x), u(x)) dx \\ (y, u) \text{ satisfies (7.35), } u(x) \in K \text{ a.e. } x \in \Omega \text{ and } g(x, y(x)) \leq \delta \quad \forall x \in \Omega. \end{cases}$$

We keep the assumptions stated in Section 2 on the data of this problem. Then we have the following result about the state equation analogous to Theorem 2.1:

THEOREM 7.1. *There exist constants $C_5 > 0$ and $\alpha \in (0, 1)$ such that for every $u \in \mathcal{K}$ (7.35) has a unique solution $y_u \in H_0^1(\Omega) \cap C^{0,\alpha}(\bar{\Omega})$ satisfying*

$$(7.36) \quad \|y_u\|_{H_0^1(\Omega)} + \|y_u\|_{C^{0,\alpha}(\bar{\Omega})} \leq C_5.$$

Furthermore the mapping $u \in (\mathcal{K}, d) \longrightarrow y_u \in H_0^1(\Omega) \cap C^{0,\alpha}(\bar{\Omega})$ is continuous.

Proof. We may assume that $\beta(0) \ni 0$. If $\text{dom}(\beta) = R$ and β is Lipschitz and of class C^1 , the result is consequence of Theorem 2.1. When β is a general maximal monotone graph in $R \times R$ it is enough to apply the standard procedure that consists in approximating (via Yosida's approximation and convolution with a smoothing kernel: see [2]) with a Lipschitz C^1 monotone function β_ϵ . In this way we obtain solutions $y_\epsilon \in H_0^1(\Omega) \cap C^{0,\alpha}(\bar{\Omega})$ of

$$\begin{cases} Ay + \beta_\epsilon(y) = f(x, y(x), u(x)) & \text{in } \Omega, \\ y = 0 & \text{on } \Gamma. \end{cases}$$

In order to pass to the limit and derive (7.36) we need a uniform estimate of y_ϵ in $H_0^1(\Omega) \cap C^{0,\alpha}(\bar{\Omega})$. Using the mean value theorem we can write

$$\begin{aligned} \beta_\epsilon(y_\epsilon) &= \beta'_\epsilon(\tilde{y}_\epsilon)y_\epsilon, \\ f(x, y_\epsilon(x), u(x)) &= f(x, 0, u(x)) + \frac{\partial f}{\partial y}(x, \tilde{y}_\epsilon, u(x))y_\epsilon, \end{aligned}$$

with $|\hat{y}_\epsilon(x)| \leq |y_\epsilon(x)|$ and $|\tilde{y}_\epsilon(x)| \leq |y_\epsilon(x)|$ for all $x \in \Omega$. Hence

$$\begin{cases} Ay_\epsilon + \left(\beta'_\epsilon(\hat{y}_\epsilon)(x) - \frac{\partial f}{\partial y}(x, \tilde{y}_\epsilon, u(x)) \right) y_\epsilon = f(x, 0, u(x)) & \text{in } \Omega, \\ y = 0 & \text{on } \Gamma. \end{cases}$$

Now applying Lemma 2.2 to the above equation we get a uniform estimate of y_ϵ in $H_0^1(\Omega) \cap L^\infty(\Omega)$. Then the hypotheses (2.3) on f imply that $\{f(x, y_\epsilon(x), u(x))\}_{\epsilon>0}$ is uniform bounded in $L^s(\Omega)$. On the other hand, arguing as in [9, Appendix] we also deduce the boundedness of $\{\beta_\epsilon(y_\epsilon)\}_{\epsilon>0}$ in $L^s(\Omega)$. Therefore $\{Ay_\epsilon\}_{\epsilon>0}$ is uniform bounded in $L^s(\Omega)$; applying Lemma 2.2 again we obtain the desired result. \square

The aim of this section is to prove the Pontryagin's principle for control problem (P_δ) . For this purpose we need the following approximation scheme. First let us observe that Proposition 6.1 is still valid and consequently there exists a number $\lambda > 0$ such that \bar{u} is a solution of the problem

$$(Q_r) \begin{cases} \min J_r(u) = \int_{\Omega} L(x, y_u(x), u(x)) dx + r \|(g(x, y_u) - \delta)^+\|_{\infty}, \\ u \in B_\lambda(\bar{u}). \end{cases}$$

Here y_u denotes the solution of (7.35) corresponding to the control u .

The next step consists in approximating (Q_r) by a new control problem with a differentiable cost functional and a state equation with a C^1 monotone term $\beta_q(y)$. Let us begin with the last question. Following Bonnans and Tiba [9] we will say that a maximal monotone graph in $R \times R$ β_q , with $q > 1$, is an $(1/q)$ -uniform approximation to β if β_q satisfies the following two conditions:

1. $\beta(t + 1/q) \geq \beta_q(t) \geq \beta(t - 1/q)$, $\forall t \in R$,
2. $\text{dom}(\beta_q) \supset \text{dom}(\beta)$.

Here we view β and β_q as multivalued operators extended on R with value $-\infty$ on the left of their domains and $+\infty$ on the right of their domains, and the inequality for sets means

$$\xi \geq \eta \geq \nu, \quad \forall \xi \in \beta(t + 1/q), \eta \in \beta_q(t), \nu \in \beta(t - 1/q).$$

A constructive procedure for $(1/q)$ -uniform approximations of class C^1 was given in [9], and the following result was proved:

PROPOSITION 7.2. *Let $u \in \mathcal{K}$. Then the problem*

$$\begin{cases} Ay + \beta_q(y) = f(x, y(x), u(x)) & \text{in } \Omega, \\ y = 0 & \text{on } \Gamma, \end{cases}$$

has a unique solution $y_{q,u} \in H_0^1(\Omega) \cap C^\alpha(\bar{\Omega})$ and $\|y_{q,u} - y_u\|_{\infty} \leq 1/q$.

Now we consider the following approximation of (Q_r) :

$$(Q_{r,q}) \begin{cases} \min J_{r,q}(u) \\ u \in B_\lambda(\bar{u}), \end{cases}$$

where

$$J_{r,q}(u) = \int_{\Omega} L(x, y_{q,u}(x), u(x)) dx + r \left(q^{-q} + \int_{\Omega} [(g(x, y_{q,u}) - \delta)^+]^q dx \right)^{1/q}.$$

PROPOSITION 7.3. *The following identity holds*

$$\inf(Q_r) = \lim_{q \rightarrow \infty} \inf(Q_{r,q}).$$

Proof. With the aid of Proposition 7.2 we get

$$\begin{aligned} & \left| \|(g(x, y_{q,u}) - \delta)^+\|_{L^q(\Omega)} - \|(g(x, y_{q,u}) - \delta)^+\|_\infty \right| \leq \\ & \left| \|(g(x, y_u) - \delta)^+\|_{L^q(\Omega)} - \|(g(x, y_u) - \delta)^+\|_\infty \right| + \\ & \left| \|(g(x, y_{q,u}) - \delta)^+\|_{L^q(\Omega)} - \|(g(x, y_u) - \delta)^+\|_{L^q(\Omega)} \right| + \\ & \left| \|(g(x, y_{q,u}) - \delta)^+\|_\infty - \|(g(x, y_u) - \delta)^+\|_\infty \right| \leq \\ & \left| \|(g(x, y_u) - \delta)^+\|_{L^q(\Omega)} - \|(g(x, y_u) - \delta)^+\|_\infty \right| + \\ & \|g(x, y_{q,u}) - g(x, y_u)\|_{L^q(\Omega)} + \|g(x, y_{q,u}) - g(x, y_u)\|_\infty \leq \\ & \left| \|(g(x, y_{q,u}) - \delta)^+\|_{L^q(\Omega)} - \|(g(x, y_u) - \delta)^+\|_\infty \right| + \frac{m(\Omega)^{1/q} + 1}{q} \rightarrow 0 \text{ as } q \rightarrow \infty. \end{aligned}$$

Therefore we can argue as in the proof of Proposition 6.2 to deduce that

$$(7.37) \quad \limsup_{q \rightarrow \infty} \inf(Q_{r,q}) \leq \inf(Q_r).$$

Let us prove the converse inequality. Let $C_5 > 0$ be the constant given in Theorem 7.1. From the properties of g we obtain

$$(7.38) \quad |g(x, t)| + \left| \frac{\partial g}{\partial y}(x, t) \right| \leq M \quad \forall (x, t) \in \bar{\Omega} \times [-C_5, +C_5]$$

for some constant $M > 0$. Applying the mean value theorem and using the hypotheses (2.3), (7.38) and Proposition 7.2 we get for some constant M' and all $u \in B_\lambda(\bar{u})$:

$$\begin{aligned} & \int_{\Omega} L(x, y_{q,u}(x), u(x)) dx + r \left(q^{-q} + \int_{\Omega} [(g(x, y_{q,u}) - \delta)^+]^q dx \right)^{1/q} \geq \\ & \int_{\Omega} L(x, y_u(x), u(x)) dx + r \left(q^{-q} + \int_{\Omega} [(g(x, y_u) - \delta)^+]^q dx \right)^{1/q} - \frac{M'}{q}. \end{aligned}$$

Therefore

$$\liminf_{q \rightarrow \infty} \inf(Q_{r,q}) \geq \liminf_{q \rightarrow \infty} \inf \left\{ \int_{\Omega} L(x, y_u(x), u(x)) dx + \right.$$

$$r \left(q^{-q} + \int_{\Omega} [(g(x, y_u) - \delta)^+]^q dx \right)^{1/q} : u \in B_{\lambda}(\bar{u}) \Big\}.$$

The proof is concluded by noting that the second term of this inequality is greater than or equal to $\inf(Q_r)$, which is proved exactly in the same way than in Proposition 6.2. \square

Now we are ready to state the extension of Pontryagin's principle.

THEOREM 7.4. *Let \bar{u} be a solution of (P_{δ}) in (K, d) , with \bar{y} its associated state. If (P_{δ}) is strongly stable on the right, then there exist $\chi \in W^{-1, \sigma}(\Omega)$, $\bar{p} \in W_0^{1, \sigma}(\Omega)$ for every $\sigma < \frac{n}{n-1}$ and $\bar{\mu} \in M(\Omega)$ such that χ is a limit point in $W^{-1, \sigma}(\Omega)$ weak of $\{\beta'_q(y_q)p_q\}$ and*

$$(7.39) \begin{cases} A^* \bar{p} + \chi = \frac{\partial f}{\partial y}(x, \bar{y}(x), \bar{u}(x)) + \frac{\partial L}{\partial y}(x, \bar{y}(x), \bar{u}(x)) + \frac{\partial g}{\partial y}(x, \bar{y}(x)) \bar{\mu} \text{ in } \Omega, \\ \bar{p} = 0 \text{ on } \Gamma, \end{cases}$$

$$(7.40) \quad \int_{\Omega} (z(x) - g(x, \bar{y}(x))) d\bar{\mu}(x) \leq 0 \quad \forall z \in C_0(\Omega) \text{ with } z(x) \leq \delta \quad \forall x \in \Omega,$$

and for every $v \in K$

$$(7.41) \quad H(x, \bar{y}(x), \bar{u}(x), \bar{p}(x)) \leq H(x, \bar{y}(x), v, \bar{p}(x)) \quad \text{a.e. } x \in \Omega.$$

Moreover if the conditions **H1)** or **H2)** of Theorem 3.3 hold, then

$$(7.42) \quad H(x, \bar{y}(x), \bar{u}(x), \bar{p}(x)) = \min_{v \in K} H(x, \bar{y}(x), v, \bar{p}(x)) \quad \text{a.e. } x \in \Omega.$$

Proof. This theorem can be proved in the same manner than Theorem 3.4: applying Proposition 7.2 we deduce that \bar{u} is a solution of (Q_r) and then Theorem 4.11 provides a minimum principle for an ϵ_q^2 -solution u_q . The adjoint state corresponding to u_q satisfies the equation

$$\begin{cases} A^* p_q + \beta'_q(y_q)p_q = \frac{\partial f}{\partial y}(x, y_q, u_q)p_q + \frac{\partial L}{\partial y}(x, y_q, u_q) + \mu_q \frac{\partial g}{\partial y}(x, y_q) \text{ in } \Omega, \\ p_q = 0 \text{ on } \Gamma, \end{cases}$$

with

$$\mu_q = r \left(q^{-q} + \int_{\Omega} [(g(x, y_q(x)) - \delta)^+]^q dx \right)^{1/q-1} [(g(x, y_q) - \delta)^+]^{q-1}.$$

The passage to the limit is carried out as in the proof of Theorem 3.4 with the only modification due to the term $\beta'_q(y_q)p_q$. That $\{\mu_q\}$ is bounded in $L^1(\Omega)$ can be proved as in Section 7, therefore the boundedness of $\{p_q\}$ in $W_0^{1, \sigma}(\Omega)$ is a consequence of Lemma 2.4. Finally from the adjoint state equation it follows that $\{\beta'_q(y_q)p_q\}$ is bounded in $W^{-1, \sigma}(\Omega)$, for all $\sigma < n/(n-1)$. Then there exists a subsequence, denoted in the same way, and an element $\chi \in W^{-1, \sigma}(\Omega)$ such that $\beta'_q(y_q)p_q \rightarrow \chi$ weakly in $W^{-1, \sigma}(\Omega)$ when $q \rightarrow \infty$. \square

REMARK 7.5. *Additional information on χ can be derived from Theorem 7.4 for particular choices of β . For instance if β is Lipschitz near $y_0 \in R$ and $x_0 \in \Omega$ is such that $y(x_0) = y_0$, then $\chi(x) \in \partial_c \beta(y(x))$ with $\partial_c \beta$ the Clarke gradient [14] of β , for x close to x_0 . See e.g. [9] for other illustrations.*

REFERENCES

- [1] R. ADAMS, *Sobolev Spaces*, Academic Press, New York, 1975.
- [2] V. BARBU, *Optimal control of variational inequalities*, vol. 100 of Lecture Notes in Mathematics, Pitman, London, 1984.
- [3] J.F. BONNANS, *El principio de Pontryagin para el control de sistemas elípticos con restricciones sobre el estado: el método de penalización*, in Jornadas Hispano-Francesas sobre Control de Sistemas Distribuidos, A. Valle, ed., Málaga, 1990, 1991, pp. 13–19.
- [4] ———, *Pontryagin's principle for the optimal control of semilinear elliptic systems with state constraints*, in 30th IEEE Conference on Control and Decision, Brighton, England, 1991, pp. 1976–1979.
- [5] J.F. BONNANS AND E. CASAS, *Contrôle de systèmes elliptiques semilinéaires comportant des contraintes sur l'état*, in Nonlinear Partial Differential Equations and Their Applications. Collège de France Seminar, H. Brezis and J. Lions, eds., vol. 8, Longman Scientific & Technical, New York, 1988, pp. 69–86.
- [6] ———, *Optimal control of semilinear multistate systems with state constraints*, SIAM J. on Control & Optim., 27 (1989), pp. 446–455.
- [7] ———, *Un principe de Pontryagin pour le contrôle des systèmes elliptiques*, J. Differential Equations, 90 (1991), pp. 288–303.
- [8] ———, *A boundary Pontryagin's principle for the optimal control of state constrained elliptic systems*, in Proc. French-Romanian Conference on Optimization, Optimal control and Partial Differential Equations, Int. Ser. Num. Math. 107, V. Barbu, J.F. Bonnans and D. Tiba eds., Birkhäuser, 241–249, 1992.
- [9] J.F. BONNANS AND D. TIBA, *Pontryagin's principle in the control of semilinear elliptic variational equations*, J. Appl. Math. and Optim., 23 (1991), pp. 299–312.
- [10] H. BREZIS, *Problèmes unilatéraux*, J. Math. Pures Appl., 51 (1972), pp. 1–68.
- [11] J. BURKE, *Calmness and exact penalization*, SIAM J. on Control & Optim., 29 (1991), pp. 493–497.
- [12] E. CASAS, *Control of an elliptic problem with pointwise state constraints*, SIAM J. on Control & Optim., 24 (1986), pp. 1309–1318.
- [13] ———, *Finite element approximations for some optimal control problems with pointwise state constraints*, in IMACS '91 13th World Congress on Computation and Applied Mathematics, R. Vichnevetsky and J. Miller, eds., vol. 3, Dublin, 1991, Criterion Press, pp. 1165–1166.
- [14] ———, *Análisis de la convergencia en la aproximación numérica de problemas de control con restricciones sobre el estado*, in Actas del XII C.E.D.Y.A./II Congreso de Matemática Aplicada, Oviedo, 1991, pp. 301–306.
- [15] F. CLARKE, *A new approach to Lagrange multipliers*, Math. Op. Res., 1 (1976), pp. 165–174.
- [16] I. EKELAND, *Nonconvex minimization problems*, Bull. Amer. Math. Soc., 1 (1979), pp. 76–91.
- [17] D. GILBARG AND N. TRUDINGER, *Elliptic Partial Differential Equations of Second Order*, Springer-Verlag, Berlin-Heidelberg-New York, 1977.
- [18] Z.-X. HE, *State constrained control problems governed by variational inequalities*, SIAM J. Control and Optim., 25 (1987), pp. 1119–1143.
- [19] J.L. LIONS, *Contrôle Optimal de Systèmes Gouvernés par des Equations aux Dérivées Partielles*, Dunod, Paris, 1968.
- [20] F. MIGNOT, *Contrôle dans les inéquations variationnelles elliptiques*, Functional Analysis, 22 (1974), pp. 130–185.
- [21] F. MIGNOT AND J. PUEL, *Optimal control in some variational inequalities*, SIAM J. on Control and Optim., 22 (1984), pp. 466–476.
- [22] G. STAMPACCHIA, *Le problème de Dirichlet pour les équations elliptiques du second ordre à coefficients discontinus*, Ann. Inst. Fourier (Grenoble), 15 (1965), pp. 189–258.