



THE ORIGIN OF SPECIES

BY MEANS OF NATURAL SELECTION,

OR THE


PRESERVATION OF FAVOURED RACES IN THE STRUGGLE FOR LIFE.

By CHARLES DARWIN, M.A.,


FELLOW OF THE ROYAL, GEOLOGICAL, LINNÆAN, ETC., SOCIETIES;
AUTHOR OF 'JOURNAL OF RESEARCHES DURING H. M. S. BEAGLE'S VOYAGE
ROUND THE WORLD.'


LONDON:
JOHN MURRAY, ALBEMARLE STREET.
1859.

The right of Translation is reserved.



THE 'UNIFORMITARIANISM' OF GRADUALISM






"...natural selection is daily and hourly scrutinising [...] at the improvement of each organic being [...]. We see nothing of these slow changes in progress, until the hand of time has marked the long lapse of ages."

"The **amount of organic change** in the fossils [...] probably serves as a **fair measure of the lapse of actual time**."

C. R. Darwin (1859)

'ABOMINABLE MYSTERIES'

to 90. Frank comes back in to 486 beginning of them wants for I ha pic red Bells' apig. It is july bit. The repid dentyment, and an as we can just of to white within ment gentified time within the first is an obsailable mysters. Certains it is he was a find the first of link cold time of de high level; but with it is uprivately



"The **rapid** development as far as we can judge of all the higher plants within recent geological times is an **abominable mystery**".


C. R. Darwin to J. Hooker (1879)

~300,000 extant Angiosperms (younger) vs ~1000 extant Gymnosperms (older).

'GHOST CONTINENTS'

"Nothing is more extraordinary [...] than the apparently very sudden or abrupt development of the higher plants. I have sometimes speculated whether there did not exist somewhere during long ages an extremely isolated continent, perhaps near the South Pole."


C. R. Darwin to J. Hooker (1881)




'ON THE IMPERFECTION OF THE GEOLOGICAL RECORD'

"If then the **geological record be as imperfect as I believe it to be** [...] the main objections to the theory of natural selection are greatly diminished or disappear."

C. R. Darwin (1859)






Tempo and Mode in Evolution

George Gaylord Simpson

A Columbia Classic in Evolution


With a new introduction by George Gaylord Simpson



PALAEONTOLOGY'S BRAVADO


"They [experimental biologists] may reveal what happens to a hundred rats in the course of ten years under fixed and simple conditions, but not what happened to a billion rats in the course of ten million years under the fluctuating conditions of earth history.


Obviously, the latter problem is much more important."




G.G. Simpson (1944)


'QUANTUM EVOLUTION'






Fast diversification result from colonisation of new adaptive zones: the opening of ecological opportunities (e.g., key innovations) that materialise at the clade level.





[&]quot;Much of life's diversity originated from adaptive radiations".

BIODIVERSITY IS THE RESULT OF A DIVERSIFICATION PROCESS





Birth-death models

 λ (Speciation rate)

 μ (Extinction rate)


Few major discrete shifts following 'ecological opportunities' (e.g., key innovations, geographical colonization)






A **less predictable** interplay of species ecologies with their environment:

Contingent diversification dynamics.





 $\Phi(\lambda(t), \mu(t), t)$ Probability of a lineage at time t going extinct by the present.

$$\Phi(\lambda_0, \mu_0, t + \delta t) = \mu_0 \delta t + (1 - \mu_0 \delta t) \times \lambda_0 \delta t \times \left[\iint \frac{1}{2\pi\sigma_\lambda \sigma_\mu \delta t} \exp\left(\frac{(\ln(\lambda_1) - \ln(\lambda_0))^2}{2\sigma_\lambda \sqrt{\delta t}} + \frac{(\ln(\mu_1) - \ln(\mu_0))^2}{2\sigma_\mu \sqrt{\delta t}}\right) \Phi(\lambda_1, \mu_1, t) d\lambda_1 d\mu_1 \right]^2 + (3) \\
(1 - \mu_0 \delta t) \times (1 - \lambda_0 \delta t) \times \int \frac{1}{2\pi\sigma_\lambda \sigma_\mu \delta t} \exp\left(\frac{(\ln(\lambda_1) - \ln(\lambda_0))^2}{2\sigma_\lambda \sqrt{\delta t}} + \frac{(\ln(\mu_1) - \ln(\mu_0))^2}{2\sigma_\mu \sqrt{\delta t}}\right) \Phi(\lambda_1, \mu_1, t) d\lambda_1 d\mu_1.$$

BAYESIAN DATA AUGMENTATION

The BDD model remains (in)famously intractable.

Bayesian data augmentation to estimate the posterior distribution.

$$p(\theta | o)$$


$$p(\theta | o) = \int_{A} p(\theta | o, a) p(a | o) da$$

 $\theta = \text{parameters}$

o = observed data


a = augmented data

BAYESIAN DATA AUGMENTATION OF CONSTANT BIRTH-DEATH




What are the past rates of speciation and extinction?

BAYESIAN DATA AUGMENTATION OF CONSTANT BIRTH-DEATH



BAYESIAN DATA AUGMENTATION OF DIFFUSION




$$x_1 \sim N(x_a, \sigma^2 t)$$

 $x_2 \sim N(x_a, \sigma^2 t)$


 σ^2 = diffusion rate


BAYESIAN DATA AUGMENTATION OF DIFFUSION




$$\int_{X} \prod_{i=1}^{n-1} x_{i+1} \sim N\left(x_{i}, \sigma^{2} \frac{t}{n-1}\right) dx$$

$$\sigma^2$$
 = diffusion rate







COMBINING PALEO- AND NEONTOLOGICAL INFORMATION





PALEO-EXTINCTION CURVES





SPECIATION RATE HETEROGENEITY

















SUPPORT FOR BDD OVER CLADE-WIDE SHIFTS







Viktor Senderov










In **extant-only phylogenies**, diversity seems to always increase or stabilise: used as evidence for **`adaptive radiations'** and **`ecological limits'**.













CLADE-LEVEL PROCESSES


Diversity stagnates and decreases through **filling (diversity-dependence)** or **closing** of **`adaptive zones'**.


















B One complete (data augmented)
FBDD tree



 ${f C}$ Latent speciation rate $\lambda(t)$









$$dln(\lambda_i(t)) = \alpha_{\lambda}dt + \sigma_{\lambda}dW(t)$$


$$dln(\mu_i(t)) = \alpha_\mu dt + \sigma_\mu dW(t)$$

27 groups encompassing ~5500 extinct and ~3000 extant species.




Quintero et al. (2025) NatEcoEvo








UNDERLYING DIVERSIFICATION: TUATARA




UNDERLYING DIVERSIFICATION: NOTOSUCHIA










effect on speciation effect on extinction













In J. R. R. Tolkein's Middle Earth []. R. R. Tolkien, *The Lord of the Rings* (Mariner Books, Boston, 2012)], the Ents lost their wives and thus had no means of regenerating their race, hence the term the Entwives effect.













N. Eldredge & S.J. Gould

Speciation as the main mechanism of evolutionary change.

AGE-DEPENDENT SPECIATION AND EXTINCTION

Lineages that do not speciate:

Become **stagnant** given the **homogenising and stabilising** effects of gene flow.




Become bested by spectating (changing) lineages and an unstable environment, **becoming maladapted**.



Decrease their speciation and increase their extinction rates.

'ZARATHUSTRAN HYPOTHESIS'



The maintenance of life does not merely follow from conservation, but rather from constant renovation, movement and expansion.

"And life itself confided the secret to me: behold, it said, I am that which **must** always overcome itself."

Thus Spoke Zarathustra (1883)

I. **Reject a simple model of diversification** explained by clade-level processes, such as adaptive radiations or diversity equilibria.

II. Suggest a more **nuanced**, **dynamic and unpredictable** diversification process with **species as individuals:** species traits and their specific environmental context periodically propitiate suitable conditions for speciation pulses.

III. **Stagnation** brought about by **a lack of speciation** causes most lineages to lag behind a continuously reshaping (biotic and abiotic) environment.

IV. The **few successful individual lineages** that depart from this general tendency **disproportionally engender biodiversity**, yet sometimes fail to compensate a general **insidious loss of macroevolutionary fitness**, leading to the **recurrent decline and fall of clades**.





Tapestree.jl
Unraveling the evolutionary tapestry: Tapestree is a Julia package of phylogenetic analyses of diversification, trait and biogeographic dynamics.

