

Around the modelling of negative materials

Lucas Chesnel¹

Coll. with A.-S. Bonnet-Ben Dhia², P. Ciarlet², C. Carvalho²,
X. Claeys³, S.A. Nazarov⁴.

¹Inverse Problems Research Group, Aalto University, Finland

²POems team, Ensta, Palaiseau, France

³LJLL, Paris VI, France

⁴FMM, St. Petersburg State University, Russia



Introduction: general framework

- ▶ Scattering by a **metal** in electromagnetism in **time-harmonic** regime at **optical frequency**.
- ▶ For **metals** at optical frequency, $\Re \varepsilon(\omega) < 0$ and $\Im m \varepsilon(\omega) \ll |\Re \varepsilon(\omega)|$.
⇒ We neglect losses and study the ideal case $\varepsilon(\omega) \in (-\infty; 0)$.

Positive material

$$\varepsilon > 0$$

and $\mu > 0$

Negative metal

$$\varepsilon < 0$$

and $\mu > 0$


Introduction: general framework

- ▶ Scattering by a **metal** in electromagnetism in **time-harmonic** regime at **optical frequency**.
- ▶ For **metals** at optical frequency, $\Re \varepsilon(\omega) < 0$ and $\Im m \varepsilon(\omega) \ll |\Re \varepsilon(\omega)|$.
⇒ We neglect losses and study the ideal case $\varepsilon(\omega) \in (-\infty; 0)$.

Positive material

$$\varepsilon > 0$$

and $\mu > 0$



Negative metal



$$\varepsilon < 0$$

and $\mu > 0$

- ▶ Waves called **Surface Plasmon Polaritons** can propagate **at the interface** between a dielectric and a negative metal.

Introduction: applications

- ▶ **Surface Plasmons Polaritons** can propagate information. Physicists hope to exploit them to reduce the size of **computer chips**.




Figures from O'Connor *et al.*, *Appl. Phys. Lett.* 95, 171112 (2009)

- ▶ In this context, physicists use **singular geometries** to **focus energy**. It allows to stock information.

Introduction: in this talk

- We study a scalar model problem set in a **bounded** domain $\Omega \subset \mathbb{R}^2$:

$$(\mathcal{P}) \quad \left| \begin{array}{l} \text{Find } u \in H_0^1(\Omega) \text{ s.t.:} \\ -\operatorname{div}(\sigma \nabla u) = f \text{ in } \Omega. \end{array} \right.$$




Introduction: in this talk

- We study a scalar model problem set in a **bounded** domain $\Omega \subset \mathbb{R}^2$:

$$(\mathcal{P}) \quad \left| \begin{array}{l} \text{Find } u \in H_0^1(\Omega) \text{ s.t.:} \\ -\operatorname{div}(\sigma \nabla u) = f \text{ in } \Omega. \end{array} \right.$$

- $H_0^1(\Omega) = \{v \in L^2(\Omega) \mid \nabla v \in L^2(\Omega); v|_{\partial\Omega} = 0\}$
- f is the source term in $H^{-1}(\Omega)$




Introduction: in this talk

- We study a scalar model problem set in a **bounded** domain $\Omega \subset \mathbb{R}^2$:

$$(\mathcal{P}) \quad \left| \begin{array}{l} \text{Find } u \in H_0^1(\Omega) \text{ s.t.:} \\ -\operatorname{div}(\sigma \nabla u) = f \text{ in } \Omega. \end{array} \right.$$

- $H_0^1(\Omega) = \{v \in L^2(\Omega) \mid \nabla v \in L^2(\Omega); v|_{\partial\Omega} = 0\}$
- f is the source term in $H^{-1}(\Omega)$




Introduction: in this talk


- We study a scalar model problem set in a **bounded** domain $\Omega \subset \mathbb{R}^2$:

$$(\mathcal{P}) \quad \left| \begin{array}{l} \text{Find } u \in H_0^1(\Omega) \text{ s.t.:} \\ -\operatorname{div}(\sigma \nabla u) = f \text{ in } \Omega. \end{array} \right.$$

- $H_0^1(\Omega) = \{v \in L^2(\Omega) \mid \nabla v \in L^2(\Omega); v|_{\partial\Omega} = 0\}$
- f is the source term in $H^{-1}(\Omega)$



- We slightly **round** the **interface** Σ :



$$(\mathcal{P}^\delta) \quad \left| \begin{array}{l} \text{Find } u^\delta \in H_0^1(\Omega) \text{ s.t.:} \\ -\operatorname{div}(\sigma^\delta \nabla u^\delta) = f \text{ in } \Omega. \end{array} \right.$$


- δ denotes the **radius of curvature** of the rounded interface at the origin.

Introduction: in this talk


- ▶ We study a scalar model problem set in a **bounded** domain $\Omega \subset \mathbb{R}^2$:

$$(\mathcal{P}) \quad \left| \begin{array}{l} \text{Find } u \in H_0^1(\Omega) \text{ s.t.:} \\ -\operatorname{div}(\sigma \nabla u) = f \text{ in } \Omega. \end{array} \right.$$

- $H_0^1(\Omega) = \{v \in L^2(\Omega) \mid \nabla v \in L^2(\Omega); v|_{\partial\Omega} = 0\}$
- f is the source term in $H^{-1}(\Omega)$



- ▶ We slightly **round** the **interface** Σ :



$$(\mathcal{P}^\delta) \quad \left| \begin{array}{l} \text{Find } u^\delta \in H_0^1(\Omega) \text{ s.t.:} \\ -\operatorname{div}(\sigma^\delta \nabla u^\delta) = f \text{ in } \Omega. \end{array} \right.$$

- δ denotes the **radius of curvature** of the rounded interface at the origin.



What is the **behaviour** of the **sequence** $(u^\delta)_\delta$ when δ tends to zero?

Outline of the talk

1 Numerical experiments

To get an **intuition**, we **discretize** (\mathcal{P}^δ) and observe what happens when $\delta \rightarrow 0$.

Outline of the talk

1 Numerical experiments

To get an **intuition**, we **discretize** (\mathcal{P}^δ) and observe what happens when $\delta \rightarrow 0$.

2 Properties of the limit problem

We present the properties of the **limit problem** in the geometry with the **real corner** ($\delta = 0$). Since σ changes sign, **original phenomena** appear.

Outline of the talk

1 Numerical experiments

To get an **intuition**, we **discretize** (\mathcal{P}^δ) and observe what happens when $\delta \rightarrow 0$.

2 Properties of the limit problem

We present the properties of the **limit problem** in the geometry with the **real corner** ($\delta = 0$). Since σ changes sign, **original phenomena** appear.

3 Asymptotic analysis

We prove a curious **instability** phenomenon: for certain configurations, (\mathcal{P}^δ) **critically depends** on δ .


1 Numerical experiments

2 Properties of the limit problem

3 Asymptotic analysis


Setting

- ▶ For the numerical experiments, we **round the corner** in a particular way




Setting

- ▶ For the numerical experiments, we **round the corner** in a particular way




Setting

- ▶ For the numerical experiments, we **round the corner** in a particular way



Setting


- ▶ For the numerical experiments, we **round the corner** in a particular way



δ is the **rounding**
parameter

Setting


- For the numerical experiments, we **round the corner** in a particular way (in this domain, we can **separate variables**).



δ is the **rounding parameter**

Setting

- ▶ For the numerical experiments, we **round the corner** in a particular way (in this domain, we can **separate variables**).




δ is the **rounding parameter**

- ▶ Our goal is to study the behaviour of the solution, *if it is well-defined*, of

$$(\mathcal{P}^\delta) \quad \left| \quad \begin{array}{l} \text{Find } u^\delta \in H_0^1(\Omega^\delta) \text{ such that:} \\ -\operatorname{div}(\sigma^\delta \nabla u^\delta) = f \quad \text{in } \Omega^\delta. \end{array} \right.$$

Setting

- ▶ For the numerical experiments, we **round the corner** in a particular way (in this domain, we can **separate variables**).



δ is the **rounding parameter**

- ▶ Our goal is to study the behaviour of the solution, *if it is well-defined*, of

$$(\mathcal{P}^\delta) \quad \left| \begin{array}{l} \text{Find } u^\delta \in H_0^1(\Omega^\delta) \text{ such that:} \\ -\operatorname{div}(\sigma^\delta \nabla u^\delta) = f \quad \text{in } \Omega^\delta. \end{array} \right.$$

- ▶ We approximate u^δ , *assuming it is well-defined*, by a **usual P1 Finite Element Method**. We compute the solution u_h^δ of the discretized problem with *FreeFem++* (**F. Hecht**).

We display the behaviour of u_h^δ as $\delta \rightarrow 0$ with *Paraview* and *Matlab*.

Numerical experiments 1/2


$\sigma_1 = 1$ and $\sigma_2 = 1$ (positive materials)

Numerical experiments 1/2

$\sigma_1 = 1$ and $\sigma_2 = 1$ (positive materials)

(...)

u_h^δ w.r.t. δ



$\|\nabla u_h^\delta\|_{\Omega^\delta}$ w.r.t. $1 - \delta$

- ▶ For **positive materials**, it is well-known that $(u^\delta)_\delta$ converges to u , the solution in the limit geometry.
- ▶ The **rate of convergence** depends on the **regularity** of u .
- ▶ To avoid to mesh Ω^δ , we can **approximate u^δ** by u_h .

Numerical experiments 2/2

... and what about for a **sign-changing** σ ???


$$\sigma_1 = 1 \text{ and } \sigma_2 = -0.9999$$

Numerical experiments 2/2

... and what about for a **sign-changing** σ ???

$$\sigma_1 = 1 \text{ and } \sigma_2 = -0.9999$$

(...)



u_h^δ w.r.t. δ

$\|\nabla u_h^\delta\|_{\Omega^\delta}$ w.r.t. $1 - \delta$

- For this configuration, u^δ seems to **depend critically** on δ .

In this talk, our goal is to **explain** this behaviour.

- 1 Numerical experiments
- 2 Properties of the limit problem
- 3 Asymptotic analysis

Mathematical difficulty

- Classical case $\sigma > 0$ everywhere:

$$a(u, u) = \int_{\Omega} \sigma |\nabla u|^2 \geq \min(\sigma) \|u\|_{H_0^1(\Omega)}^2 \quad \text{coercivity}$$

Lax-Milgram theorem \Rightarrow (\mathcal{P}) well-posed.

Mathematical difficulty

- Classical case $\sigma > 0$ everywhere:

$$a(u, u) = \int_{\Omega} \sigma |\nabla u|^2 \geq \min(\sigma) \|u\|_{H_0^1(\Omega)}^2 \quad \text{coercivity}$$

Lax-Milgram theorem \Rightarrow (\mathcal{P}) well-posed.

----- VS. -----

- The case σ changes sign:

~~$$a(u, u) = \int_{\Omega} \sigma |\nabla u|^2 \geq C \|u\|_{H_0^1(\Omega)}^2 \quad \text{loss of coercivity}$$~~

Mathematical difficulty

- Classical case $\sigma > 0$ everywhere:

$$a(u, u) = \int_{\Omega} \sigma |\nabla u|^2 \geq \min(\sigma) \|u\|_{H_0^1(\Omega)}^2 \quad \text{coercivity}$$

Lax-Milgram theorem \Rightarrow (\mathcal{P}) well-posed.

----- VS. -----

- The case σ changes sign:

~~$$a(u, u) = \int_{\Omega} \sigma |\nabla u|^2 \geq C \|u\|_{H_0^1(\Omega)}^2 \quad \text{loss of coercivity}$$~~


- ▶ When $\sigma_2 = -\sigma_1$, (\mathcal{P}) is always ill-posed (Costabel-Stephan 85). For a symmetric domain (w.r.t. Σ) we can build a kernel of infinite dimension.

Problems with a sign changing coefficient

$$(\mathcal{P}) \quad \left| \begin{array}{l} \text{Find } u \in H_0^1(\Omega) \text{ such that:} \\ -\operatorname{div}(\sigma \nabla u) = f \quad \text{in } \Omega. \end{array} \right.$$


► We have the following properties (see Costabel and Stephan 85, Dauge and Texier 97, Bonnet-Ben Dhia *et al.* 99,10,12,13):

Smooth interface Σ



✓ (\mathcal{P}) well-posed in the Fredholm sense iff $\kappa_\sigma = \sigma_2/\sigma_1 \neq -1$.

Interface Σ with a corner




✓ (\mathcal{P}) well-posed in the Fredholm sense iff $\kappa_\sigma \notin I_c = [-\ell; -1/\ell]$, $\ell = (2\pi - \vartheta)/\vartheta$.

Problems with a sign changing coefficient

$$(\mathcal{P}) \quad \left| \begin{array}{l} \text{Find } u \in H_0^1(\Omega) \text{ such that:} \\ -\operatorname{div}(\sigma \nabla u) = f \quad \text{in } \Omega. \end{array} \right.$$


► We have the following properties (see Costabel and Stephan 85, Dauge and Texier 97, Bonnet-Ben Dhia *et al.* 99,10,12,13):

Smooth interface Σ



✓ (\mathcal{P}) well-posed in the Fredholm sense iff $\kappa_\sigma = \sigma_2/\sigma_1 \neq -1$.

Interface Σ with a corner



✓ (\mathcal{P}) well-posed in the Fredholm sense iff $\kappa_\sigma \notin I_c = [-\ell; -1/\ell]$, $\ell = (2\pi - \vartheta)/\vartheta$.




Well-posedness depends on the smoothness of Σ and on σ .

The problematic of the rounded corner

$$(\mathcal{P}) \quad \left| \quad \begin{array}{l} \text{Find } u \in H_0^1(\Omega) \text{ such that:} \\ -\operatorname{div}(\sigma \nabla u) = f \quad \text{in } \Omega. \end{array} \right.$$


- When the interface has a **corner**, (\mathcal{P}) is well-posed in the Fredholm sense iff $\kappa_\sigma \notin I_c$ (the critical interval).




The problematic of the rounded corner

$$(\mathcal{P}) \quad \left| \begin{array}{l} \text{Find } u \in H_0^1(\Omega) \text{ such that:} \\ -\operatorname{div}(\sigma \nabla u) = f \quad \text{in } \Omega. \end{array} \right.$$

- When the interface has a **corner**, (\mathcal{P}) is well-posed in the Fredholm sense iff $\kappa_\sigma \notin I_c$ (the critical interval).




- When the interface is **smooth**, (\mathcal{P}) is well-posed in the Fredholm sense iff $\kappa_\sigma \neq -1$.



The problematic of the rounded corner



$$(\mathcal{P}) \quad \left| \begin{array}{l} \text{Find } u \in H_0^1(\Omega) \text{ such that:} \\ -\operatorname{div}(\sigma \nabla u) = f \quad \text{in } \Omega. \end{array} \right.$$

- When the interface has a **corner**, (\mathcal{P}) is well-posed in the Fredholm sense iff $\kappa_\sigma \notin I_c$ (the critical interval).



- When the interface is **smooth**, (\mathcal{P}) is well-posed in the Fredholm sense iff $\kappa_\sigma \neq -1$.


What happens for a **slightly rounded corner** when $\kappa_\sigma \in I_c \setminus \{-1\}$?




The problematic of the rounded corner

$$(\mathcal{P}) \quad \left| \begin{array}{l} \text{Find } u \in H_0^1(\Omega) \text{ such that:} \\ -\operatorname{div}(\sigma \nabla u) = f \quad \text{in } \Omega. \end{array} \right.$$


- When the interface has a **corner**, (\mathcal{P}) is well-posed in the Fredholm sense iff $\kappa_\sigma \notin I_c$ (the critical interval).



- When the interface is **smooth**, (\mathcal{P}) is well-posed in the Fredholm sense iff $\kappa_\sigma \neq -1$.



What happens for a **slightly rounded corner** when $\kappa_\sigma \in I_c \setminus \{-1\}$?




- We need to clarify the properties of (\mathcal{P}) when the **interface** has a **corner** in the case $\kappa_\sigma \in I_c \setminus \{-1\}$.

Properties of the limit problem inside the critical interval

$$(\mathcal{P}) \quad \left| \begin{array}{l} \text{Find } u \in H_0^1(\Omega) \text{ such that:} \\ -\operatorname{div}(\sigma \nabla u) = f \quad \text{in } \Omega. \end{array} \right.$$


- To simplify the presentation, we work on a particular configuration.



Properties of the limit problem inside the critical interval

$$(\mathcal{P}) \quad \left| \begin{array}{l} \text{Find } u \in H_0^1(\Omega) \text{ such that:} \\ -\operatorname{div}(\sigma \nabla u) = f \quad \text{in } \Omega. \end{array} \right.$$


- ▶ To simplify the presentation, we work on a particular configuration.



Properties of the limit problem inside the critical interval

$$(\mathcal{P}) \quad \left| \begin{array}{l} \text{Find } u \in H_0^1(\Omega) \text{ such that:} \\ -\operatorname{div}(\sigma \nabla u) = f \quad \text{in } \Omega. \end{array} \right.$$


- ▶ To simplify the presentation, we work on a particular configuration.



Properties of the limit problem inside the critical interval

$$(\mathcal{P}) \quad \left| \begin{array}{l} \text{Find } u \in H_0^1(\Omega) \text{ such that:} \\ -\operatorname{div}(\sigma \nabla u) = f \quad \text{in } \Omega. \end{array} \right.$$

- ▶ To simplify the presentation, we work on a particular configuration.




- ▶ Using the **variational method** of the **T-coercivity**, we prove the

PROPOSITION. The problem (\mathcal{P}) is well-posed as soon as the **contrast** $\kappa_\sigma = \sigma_2/\sigma_1$ satisfies $\kappa_\sigma \notin I_c = [-1; -1/3]$.

Properties of the limit problem inside the critical interval

$$(\mathcal{P}) \quad \left| \begin{array}{l} \text{Find } u \in H_0^1(\Omega) \text{ such that:} \\ -\operatorname{div}(\sigma \nabla u) = f \quad \text{in } \Omega. \end{array} \right.$$

- ▶ To simplify the presentation, we work on a particular configuration.




- ▶ Using the **variational method** of the **T-coercivity**, we prove the

PROPOSITION. The problem (\mathcal{P}) is well-posed as soon as the **contrast** $\kappa_\sigma = \sigma_2/\sigma_1$ satisfies $\kappa_\sigma \notin I_c = [-1; -1/3]$.

What happens when $\kappa_\sigma \in (-1; -1/3]$?

Analogy with a waveguide problem

- Bounded sector Ω




- Equation:

$$\underbrace{-\operatorname{div}(\sigma \nabla u)}_{-r^{-2}(\sigma(r\partial_r)^2 + \partial_\theta \sigma \partial_\theta)u} = f$$

Analogy with a waveguide problem

- Bounded sector Ω



- Equation:

$$\underbrace{-\operatorname{div}(\sigma \nabla u)}_{-r^{-2}(\sigma(r\partial_r)^2 + \partial_\theta \sigma \partial_\theta)u} = f$$



- **Singularities** in the sector

$$s(r, \theta) = r^\lambda \varphi(\theta)$$

Analogy with a waveguide problem

We compute the singularities $s(r, \theta) = r^\lambda \varphi(\theta)$ and we observe two cases:



► **Outside the critical interval**





Analogy with a waveguide problem

We compute the singularities $s(r, \theta) = r^\lambda \varphi(\theta)$ and we observe two cases:

Outside the critical interval





Inside the critical interval





Analogy with a waveguide problem

We compute the singularities $s(r, \theta) = r^\lambda \varphi(\theta)$ and we observe two cases:

Outside the critical interval




Inside the critical interval



How to deal with the **propagative singularities** inside the critical interval?

Analogy with a waveguide problem

- Bounded sector Ω



- Equation:


$$\underbrace{-\operatorname{div}(\sigma \nabla u)}_{-r^{-2}(\sigma(r\partial_r)^2 + \partial_\theta \sigma \partial_\theta)u} = f$$

- **Singularities** in the sector



$$s(r, \theta) = r^\lambda \varphi(\theta)$$

Analogy with a waveguide problem

- Bounded sector Ω



- Half-strip \mathcal{B}



- Equation:


$$\underbrace{-\operatorname{div}(\sigma \nabla u)}_{-r^{-2}(\sigma(r\partial_r)^2 + \partial_\theta \sigma \partial_\theta)u} = f$$

- Singularities in the sector

$$s(r, \theta) = r^\lambda \varphi(\theta)$$

Analogy with a waveguide problem

- Bounded sector Ω





- Equation:

$$\underbrace{-\operatorname{div}(\sigma \nabla u)}_{-r^{-2}(\sigma(r\partial_r)^2 + \partial_\theta \sigma \partial_\theta)u} = f$$

- Singularities in the sector

$$s(r, \theta) = r^\lambda \varphi(\theta)$$

- Half-strip \mathcal{B}




- Equation:

$$\underbrace{-\operatorname{div}(\sigma \nabla u)}_{-(\sigma \partial_z^2 + \partial_\theta \sigma \partial_\theta)u} = e^{-2z} f$$

Analogy with a waveguide problem

- Bounded sector Ω





- Equation:

$$\underbrace{-\operatorname{div}(\sigma \nabla u)}_{-r^{-2}(\sigma(r\partial_r)^2 + \partial_\theta \sigma \partial_\theta)u} = f$$

- Singularities in the sector

$$s(r, \theta) = r^\lambda \varphi(\theta)$$

- Half-strip \mathcal{B}



- Equation:


$$\underbrace{-\operatorname{div}(\sigma \nabla u)}_{-(\sigma \partial_z^2 + \partial_\theta \sigma \partial_\theta)u} = e^{-2z} f$$

- Modes in the strip

$$m(z, \theta) = e^{-\lambda z} \varphi(\theta)$$

Analogy with a waveguide problem

- Bounded sector Ω



- Equation:

- Singularities** in the sector


$$s(r, \theta) = r^\lambda \varphi(\theta)$$

$$s \in H^1(\Omega)$$

- Half-strip \mathcal{B}

$$(z, \theta) = (-\ln r, \theta)$$

$$(r, \theta) = (e^{-z}, \theta)$$



- Equation:

- Modes** in the strip


$$m(z, \theta) = e^{-\lambda z} \varphi(\theta)$$

m is evanescent

$$\Re \lambda > 0$$

Analogy with a waveguide problem

- Bounded sector Ω



- Singularities** in the sector

$$s(r, \theta) = r^\lambda \varphi(\theta)$$

$$= \cancel{r^a} (\cos b \ln r + i \sin b \ln r) \varphi(\theta)$$

$(\Re e \lambda = a, \Im m \lambda = b)$


$$s \in H^1(\Omega)$$

$$s \notin H^1(\Omega)$$

- Half-strip \mathcal{B}

$$(z, \theta) = (-\ln r, \theta)$$

$$(r, \theta) = (e^{-z}, \theta)$$



- Equation

- Modes** in the strip


$$m(z, \theta) = e^{-\lambda z} \varphi(\theta)$$

$$= \cancel{e^{-az}} (\cos bz - i \sin bz) \varphi(\theta)$$

m is **evanescent**
 m is **propagative**

Analogy with a waveguide problem

- Bounded sector Ω



- Equation:

$$\underbrace{-\operatorname{div}(\sigma \nabla u)}_{-r^{-2}(\sigma(r\partial_r)^2 + \partial_\theta \sigma \partial_\theta)u} = f$$

- Singularities** in the sector


$$s(r, \theta) = r^\lambda \varphi(\theta)$$

$$= \cancel{r^a} (\cos b \ln r + i \sin b \ln r) \varphi(\theta)$$


$(\Re \lambda = a, \Im \lambda = b)$

$$s \in H^1(\Omega)$$

$$s \notin H^1(\Omega)$$



- Half-strip \mathcal{B}



- Equation:

$$\underbrace{-\operatorname{div}(\sigma \nabla u)}_{-(\sigma \partial_z^2 + \partial_\theta \sigma \partial_\theta)u} = e^{-2z} f$$

- Modes** in the strip

$$m(z, \theta) = e^{-\lambda z} \varphi(\theta)$$

$$= \cancel{e^{-az}} (\cos bz - i \sin bz) \varphi(\theta)$$

$$\Re \lambda > 0$$


$$\Re \lambda = 0$$

m is **evanescent**

m is **propagative**


- This encourages us to use **modal decomposition** in the half-strip.

Modal analysis in the waveguide




► **Outside the critical interval**. All the modes are exponentially growing or decaying.
→ We look for an exponentially decaying solution. H^1 framework

Modal analysis in the waveguide




► **Outside the critical interval**. All the modes are exponentially growing or decaying.
→ We look for an exponentially decaying solution. H^1 framework




► **Inside the critical interval**. There are exactly two propagative modes.

Modal analysis in the waveguide



► **Outside the critical interval**. All the modes are exponentially growing or decaying.
→ We look for an exponentially decaying solution. H^1 framework



► **Inside the critical interval**. There are exactly two propagative modes.
→ The decomposition on the outgoing modes leads to look for a solution of the form

$$u = \underbrace{c_1 \varphi_1 e^{\lambda_1 z}}_{\text{propagative part}} + \underbrace{u_e}_{\text{evanescent part}}$$

non H^1 framework

The new functional framework

Consider $0 < \beta < 2$, ζ a cut-off function (equal to 1 in $+\infty$) and define

$$W_{-\beta} = \{v \mid e^{\beta z} v \in H_0^1(\mathcal{B})\} \quad \text{space of exponentially decaying functions}$$

The new functional framework

Consider $0 < \beta < 2$, ζ a cut-off function (equal to 1 in $+\infty$) and define

$W_{-\beta} = \{v \mid e^{\beta z} v \in H_0^1(\mathcal{B})\}$ space of exponentially decaying functions

$W_{\beta} = \{v \mid e^{-\beta z} v \in H_0^1(\mathcal{B})\}$ space of exponentially growing functions

The new functional framework

Consider $0 < \beta < 2$, ζ a cut-off function (equal to 1 in $+\infty$) and define

$W_{-\beta} = \{v \mid e^{\beta z} v \in H_0^1(\mathcal{B})\}$	space of exponentially decaying functions
$W^+ = \text{span}(\zeta \varphi_1 e^{\lambda_1 z}) \oplus W_{-\beta}$	propagative part + evanescent part
$W_{\beta} = \{v \mid e^{-\beta z} v \in H_0^1(\mathcal{B})\}$	space of exponentially growing functions

The new functional framework

Consider $0 < \beta < 2$, ζ a cut-off function (equal to 1 in $+\infty$) and define

$$\begin{aligned} \mathring{W}_{-\beta} &= \{v \mid e^{\beta z} v \in H_0^1(\mathcal{B})\} && \text{space of exponentially decaying functions} \\ \mathring{W}^+ &= \text{span}(\zeta \varphi_1 e^{\lambda_1 z}) \oplus \mathring{W}_{-\beta} && \text{propagative part} + \text{evanescent part} \\ \mathring{W}_{\beta} &= \{v \mid e^{-\beta z} v \in H_0^1(\mathcal{B})\} && \text{space of exponentially growing functions} \end{aligned}$$

The new functional framework

Consider $0 < \beta < 2$, ζ a cut-off function (equal to 1 in $+\infty$) and define

$$\begin{aligned} \mathring{W}_{-\beta} &= \{v \mid e^{\beta z} v \in H_0^1(\mathcal{B})\} && \text{space of exponentially decaying functions} \\ \mathring{W}^+ &= \text{span}(\zeta \varphi_1 e^{\lambda_1 z}) \oplus \mathring{W}_{-\beta} && \text{propagative part} + \text{evanescent part} \\ \mathring{W}_{\beta} &= \{v \mid e^{-\beta z} v \in H_0^1(\mathcal{B})\} && \text{space of exponentially growing functions} \end{aligned}$$

THEOREM. Let $\kappa_{\sigma} \in (-1; -1/3)$ and $0 < \beta < 2$. The operator $A^+ : \text{div}(\sigma \nabla \cdot)$ from \mathring{W}^+ to \mathring{W}_{β}^* is an **isomorphism**.

The new functional framework

Consider $0 < \beta < 2$, ζ a cut-off function (equal to 1 in $+\infty$) and define

$$\begin{aligned} \mathring{W}_{-\beta} &= \{v \mid e^{\beta z} v \in H_0^1(\mathcal{B})\} && \text{space of exponentially decaying functions} \\ \mathring{W}^+ &= \text{span}(\zeta \varphi_1 e^{\lambda_1 z}) \oplus \mathring{W}_{-\beta} && \text{propagative part} + \text{evanescent part} \\ \mathring{W}_{\beta} &= \{v \mid e^{-\beta z} v \in H_0^1(\mathcal{B})\} && \text{space of exponentially growing functions} \end{aligned}$$

THEOREM. Let $\kappa_{\sigma} \in (-1; -1/3)$ and $0 < \beta < 2$. The operator $A^+ : \text{div}(\sigma \nabla \cdot)$ from \mathring{W}^+ to \mathring{W}_{β}^* is an **isomorphism**.

IDEAS OF THE PROOF:

- 1 $A_{-\beta} : \text{div}(\sigma \nabla \cdot)$ from $\mathring{W}_{-\beta}$ to \mathring{W}_{β}^* is **injective** but **not surjective**.

The new functional framework

Consider $0 < \beta < 2$, ζ a cut-off function (equal to 1 in $+\infty$) and define

$$\begin{aligned} \mathring{W}_{-\beta} &= \{v \mid e^{\beta z} v \in H_0^1(\mathcal{B})\} && \text{space of exponentially decaying functions} \\ \mathring{W}^+ &= \text{span}(\zeta \varphi_1 e^{\lambda_1 z}) \oplus \mathring{W}_{-\beta} && \text{propagative part} + \text{evanescent part} \\ \mathring{W}_{\beta} &= \{v \mid e^{-\beta z} v \in H_0^1(\mathcal{B})\} && \text{space of exponentially growing functions} \end{aligned}$$

THEOREM. Let $\kappa_{\sigma} \in (-1; -1/3)$ and $0 < \beta < 2$. The operator $A^+ : \text{div}(\sigma \nabla \cdot)$ from \mathring{W}^+ to \mathring{W}_{β}^* is an **isomorphism**.

IDEAS OF THE PROOF:

- 1 $A_{-\beta} : \text{div}(\sigma \nabla \cdot)$ from $\mathring{W}_{-\beta}$ to \mathring{W}_{β}^* is **injective** but **not surjective**.
- 2 $A_{\beta} : \text{div}(\sigma \nabla \cdot)$ from \mathring{W}_{β} to $\mathring{W}_{-\beta}^*$ is **surjective** but **not injective**.

The new functional framework

Consider $0 < \beta < 2$, ζ a cut-off function (equal to 1 in $+\infty$) and define

$$\begin{aligned} \mathcal{W}_{-\beta} &= \{v \mid e^{\beta z} v \in H_0^1(\mathcal{B})\} && \text{space of exponentially decaying functions} \\ \mathcal{W}^+ &= \text{span}(\zeta \varphi_1 e^{\lambda_1 z}) \oplus \mathcal{W}_{-\beta} && \text{propagative part} + \text{evanescent part} \\ \mathcal{W}_{\beta} &= \{v \mid e^{-\beta z} v \in H_0^1(\mathcal{B})\} && \text{space of exponentially growing functions} \end{aligned}$$

THEOREM. Let $\kappa_{\sigma} \in (-1; -1/3)$ and $0 < \beta < 2$. The operator $A^+ : \text{div}(\sigma \nabla \cdot)$ from \mathcal{W}^+ to \mathcal{W}_{β}^* is an **isomorphism**.

IDEAS OF THE PROOF:

- 1 $A_{-\beta} : \text{div}(\sigma \nabla \cdot)$ from $\mathcal{W}_{-\beta}$ to \mathcal{W}_{β}^* is **injective** but **not surjective**.
- 2 $A_{\beta} : \text{div}(\sigma \nabla \cdot)$ from \mathcal{W}_{β} to $\mathcal{W}_{-\beta}^*$ is **surjective** but **not injective**.
- 3 The intermediate operator $A^+ : \mathcal{W}^+ \rightarrow \mathcal{W}_{\beta}^*$ is **injective** (energy integral) and **surjective** (residue theorem).

The new functional framework

Consider $0 < \beta < 2$, ζ a cut-off function (equal to 1 in $+\infty$) and define

$$\begin{aligned} \mathcal{W}_{-\beta} &= \{v \mid e^{\beta z} v \in H_0^1(\mathcal{B})\} && \text{space of exponentially decaying functions} \\ \mathcal{W}^+ &= \text{span}(\zeta \varphi_1 e^{\lambda_1 z}) \oplus \mathcal{W}_{-\beta} && \text{propagative part} + \text{evanescent part} \\ \mathcal{W}_{\beta} &= \{v \mid e^{-\beta z} v \in H_0^1(\mathcal{B})\} && \text{space of exponentially growing functions} \end{aligned}$$

THEOREM. Let $\kappa_{\sigma} \in (-1; -1/3)$ and $0 < \beta < 2$. The operator $A^+ : \text{div}(\sigma \nabla \cdot)$ from \mathcal{W}^+ to \mathcal{W}_{β}^* is an **isomorphism**.

IDEAS OF THE PROOF:

- 1 $A_{-\beta} : \text{div}(\sigma \nabla \cdot)$ from $\mathcal{W}_{-\beta}$ to \mathcal{W}_{β}^* is **injective** but **not surjective**.
- 2 $A_{\beta} : \text{div}(\sigma \nabla \cdot)$ from \mathcal{W}_{β} to $\mathcal{W}_{-\beta}^*$ is **surjective** but **not injective**.
- 3 The intermediate operator $A^+ : \mathcal{W}^+ \rightarrow \mathcal{W}_{\beta}^*$ is **injective** (energy integral) and **surjective** (residue theorem).
- 4 Limiting absorption principle to select the **outgoing mode**.

Naive approximation

- ▶ Let us try a **usual Finite Element Method** (P1 Lagrange Finite Element). We solve the problem

$$\left| \begin{array}{l} \text{Find } u_h \in V_h \text{ s.t.:} \\ \int_{\Omega} \sigma \nabla u_h \cdot \nabla v_h = \int_{\Omega} f v_h, \quad \forall v \in V_h, \end{array} \right.$$

where V_h approximates $H_0^1(\Omega)$ as $h \rightarrow 0$ (h is the **mesh size**).

Naive approximation

- ▶ Let us try a **usual Finite Element Method** (P1 Lagrange Finite Element). We solve the problem

$$\left| \begin{array}{l} \text{Find } u_h \in V_h \text{ s.t.:} \\ \int_{\Omega} \sigma \nabla u_h \cdot \nabla v_h = \int_{\Omega} f v_h, \quad \forall v \in V_h, \end{array} \right.$$

where V_h approximates $H_0^1(\Omega)$ as $h \rightarrow 0$ (h is the **mesh size**).

- ▶ We display u_h as $h \rightarrow 0$.

Naive approximation

- ▶ Let us try a **usual Finite Element Method** (P1 Lagrange Finite Element). We solve the problem

Find $u_h \in V_h$ s.t.:

$$\int_{\Omega} \sigma \nabla u_h = \int_{\Omega} f v_h, \quad \forall v \in V_h,$$

THE SEQUENCE (u_h) DOES NOT CONVERGE AS $h \rightarrow 0$!!!

where V_h approximates $H_0^1(\Omega)$ as $h \rightarrow 0$ (h is the **mesh size**).

- ▶ We display u_h as $h \rightarrow 0$.

(...)

(...)

Contrast $\kappa_{\sigma} = -0.999 \in (-1; -1/3)$.

Remark

► **Outside the critical interval**, the sequence (u_h) converges with the naive approximation.


(...)

(...)

Contrast $\kappa_\sigma = -1.001 \notin (-1; -1/3)$.


How to approximate the solution?

- ▶ We use a **PML** (*Perfectly Matched Layer*) to bound the domain \mathcal{B} + **finite elements** in the truncated strip ($\kappa_\sigma = -0.999 \in (-1; -1/3)$).



How to approximate the solution?

- ▶ We use a **PML** (*Perfectly Matched Layer*) to bound the domain \mathcal{B} + **finite elements** in the truncated strip ($\kappa_\sigma = -0.999 \in (-1; -1/3)$).



PML



Without the PML, the solution in the **truncated strip** of length L **does not converge** when $L \rightarrow \infty$. This is what we observed in our **numerical experiment** for the **rounded corner**.

A black hole phenomenon

- ▶ The same phenomenon occurs for the **Helmholtz equation**.

$$(\mathbf{x}, t) \mapsto \Re e(u(\mathbf{x})e^{-i\omega t}) \quad \text{for } \kappa_\sigma = -1/1.3$$


(...)

(...)


- ▶ Analogous phenomena occur in **cuspidal domains** in the theory of water-waves and in elasticity (**Cardone, Nazarov, Taskinen**).

- 1 Numerical experiments
- 2 Properties of the limit problem
- 3 Asymptotic analysis**

Source term problem




$$(\mathcal{P}^\delta) \quad \left| \begin{array}{l} \text{Find } u^\delta \in H_0^1(\Omega) \text{ s.t.:} \\ -\operatorname{div}(\sigma^\delta \nabla u^\delta) = f \text{ in } \Omega. \end{array} \right.$$




$$(\mathcal{P}) \quad \left| \begin{array}{l} \text{Find } u \in H_0^1(\Omega) \text{ s.t.:} \\ -\operatorname{div}(\sigma \nabla u) = f \text{ in } \Omega. \end{array} \right.$$

- The behaviour of $(u^\delta)_\delta$ depends on the properties of the **limit problem**.

Source term problem



$$(\mathcal{P}^\delta) \quad \left| \begin{array}{l} \text{Find } u^\delta \in H_0^1(\Omega) \text{ s.t.:} \\ -\operatorname{div}(\sigma^\delta \nabla u^\delta) = f \text{ in } \Omega. \end{array} \right.$$




$$(\mathcal{P}) \quad \left| \begin{array}{l} \text{Find } u \in H_0^1(\Omega) \text{ s.t.:} \\ -\operatorname{div}(\sigma \nabla u) = f \text{ in } \Omega. \end{array} \right.$$


- The behaviour of $(u^\delta)_\delta$ depends on the properties of the **limit problem**.

If (\mathcal{P}) **well-posed** (in $H_0^1(\Omega)$), then u^δ is uniquely defined for δ small enough and $(u^\delta)_\delta$ **converges to u** (as for positive materials).

Source term problem



$$(\mathcal{P}^\delta) \quad \left| \begin{array}{l} \text{Find } u^\delta \in H_0^1(\Omega) \text{ s.t.:} \\ -\operatorname{div}(\sigma^\delta \nabla u^\delta) = f \text{ in } \Omega. \end{array} \right.$$



$$(\mathcal{P}) \quad \left| \begin{array}{l} \text{Find } u \in H_0^1(\Omega) \text{ s.t.:} \\ -\operatorname{div}(\sigma \nabla u) = f \text{ in } \Omega. \end{array} \right.$$

- The behaviour of $(u^\delta)_\delta$ depends on the properties of the **limit problem**.

If (\mathcal{P}) **well-posed** (in $H_0^1(\Omega)$), then u^δ is uniquely defined for δ small enough and $(u^\delta)_\delta$ **converges to u** (as for positive materials).

If the limit problem is well-posed only in the **exotic framework**, then (\mathcal{P}^δ) **critically depends** on the value of the **rounding parameter δ** .

Source term problem

IDEA OF THE APPROACH:

① We prove the *a priori estimate* $\|u^\delta\|_{H_0^1(\Omega)} \leq c |\ln \delta|^{1/2} \|f\|_\Omega$ for all δ in some set \mathcal{S} which excludes a discrete set accumulating in zero (♠ hard part of the proof, **Nazarov's** technique).

The diagram shows a horizontal line representing a domain Ω . Above the line, there are several points marked with 'x'. Below the line, there are several points marked with 'x' that are connected by thick red horizontal bars. An arrow points from the text $\ln \mathcal{S} = \{\ln \delta, \delta \in \mathcal{S}\}$ to one of these red bars.

$$\ln \mathcal{S} = \{\ln \delta, \delta \in \mathcal{S}\}$$

$$(\mathcal{P}^\delta) \quad \begin{array}{l} \text{Find } u^\delta \in H_0^1(\Omega) \text{ s.t.:} \\ -\operatorname{div}(\sigma^\delta \nabla u^\delta) = f \text{ in } \Omega. \end{array}$$

$$(\mathcal{P}) \quad \begin{array}{l} \text{Find } u \in H_0^1(\Omega) \text{ s.t.:} \\ -\operatorname{div}(\sigma \nabla u) = f \text{ in } \Omega. \end{array}$$

The behaviour of $(u^\delta)_\delta$ depends on the properties of the limit problem.


If (\mathcal{P}) well-posed (in $H_0^1(\Omega)$), then u^δ is uniquely defined for δ small enough and $(u^\delta)_\delta$ converges to u (as for positive materials).

If the limit problem is well-posed only in the exotic framework, then (\mathcal{P}^δ) critically depends on the value of the rounding parameter δ .

Source term problem

IDEA OF THE APPROACH:

① We prove the *a priori estimate* $\|u^\delta\|_{H_0^1(\Omega)} \leq c |\ln \delta|^{1/2} \|f\|_\Omega$ for all δ in some *set* \mathcal{S} which excludes a *discrete* set accumulating in zero (♠ hard part of the proof, *Nazarov's* technique).



② We provide an *asymptotic expansion* of u^δ , denoted \hat{u}^δ with the error estimate, for some $\beta > 0$,

$$\|u^\delta - \hat{u}^\delta\|_{H_0^1(\Omega)} \leq c \delta^\beta \|f\|_\Omega, \quad \forall \delta \in \mathcal{S}.$$

Source term problem

IDEA OF THE APPROACH:

- 1 We prove the *a priori estimate* $\|u^\delta\|_{H_0^1(\Omega)} \leq c |\ln \delta|^{1/2} \|f\|_\Omega$ for all δ in some set \mathcal{S} which excludes a discrete set accumulating in zero (♠ hard part of the proof, Nazarov's technique).

$\times \text{---} \times \text{---} \times \text{---} \times \text{---} \times \text{---} \times \text{---} \times \text{---} \times \text{---} \times \text{---} \times \rightarrow \ln \delta$

$\ln \mathcal{S} = \{\ln \delta, \delta \in \mathcal{S}\}$

- 2 We provide an *asymptotic expansion* of u^δ , denoted \hat{u}^δ with the error estimate, for some $\beta > 0$,


$$\|u^\delta - \hat{u}^\delta\|_{H_0^1(\Omega)} \leq c \delta^\beta \|f\|_\Omega, \quad \forall \delta \in \mathcal{S}.$$

- 3 The behaviour of $(\hat{u}^\delta)_\delta$ can be explicitly examined as $\delta \rightarrow 0$. The sequence $(\hat{u}^\delta)_\delta$ does not converge, even for the L^2 -norm!

Source term problem

IDEA OF THE APPROACH:

- 1 We prove the *a priori estimate* $\|u^\delta\|_{H_0^1(\Omega)} \leq c |\ln \delta|^{1/2} \|f\|_\Omega$ for all δ in some set \mathcal{S} which excludes a discrete set accumulating in zero (♠ hard part of the proof, **Nazarov's** technique).




- 2 We provide an *asymptotic expansion* of u^δ , denoted \hat{u}^δ with the error estimate, for some $\beta > 0$,

$$\|u^\delta - \hat{u}^\delta\|_{H_0^1(\Omega)} \leq c \delta^\beta \|f\|_\Omega, \quad \forall \delta \in \mathcal{S}.$$


- 3 The behaviour of $(\hat{u}^\delta)_\delta$ can be explicitly examined as $\delta \rightarrow 0$. **The sequence $(\hat{u}^\delta)_\delta$ does not converge, even for the L^2 -norm!**
- 4 Conclusion.

The sequence $(u^\delta)_\delta$ **does not converge**, even for the L^2 -norm!

Source term problem



$$(\mathcal{P}^\delta) \quad \left| \begin{array}{l} \text{Find } u^\delta \in H_0^1(\Omega) \text{ s.t.:} \\ -\operatorname{div}(\sigma^\delta \nabla u^\delta) = f \text{ in } \Omega. \end{array} \right.$$



$$(\mathcal{P}) \quad \left| \begin{array}{l} \text{Find } u \in H_0^1(\Omega) \text{ s.t.:} \\ -\operatorname{div}(\sigma \nabla u) = f \text{ in } \Omega. \end{array} \right.$$

- The behaviour of $(u^\delta)_\delta$ depends on the properties of the **limit problem**.

If (\mathcal{P}) **well-posed** (in $H_0^1(\Omega)$), then u^δ is uniquely defined for δ small enough and $(u^\delta)_\delta$ **converges to u** (as for positive materials).

If the limit problem is well-posed only in the **exotic framework**, then (\mathcal{P}^δ) **critically depends** on the value of the **rounding parameter δ** .

- ▶ In the geometry with a **rounded corner**, we consider the **spectral** problem

$$\left| \begin{array}{l} \text{Find } (\lambda^\delta, u^\delta) \in \mathbb{C} \times (\mathbf{H}_0^1(\Omega) \setminus \{0\}) \text{ s.t.:} \\ -\operatorname{div}(\sigma^\delta \nabla u^\delta) = \lambda^\delta u^\delta \quad \text{in } \Omega. \end{array} \right.$$

- ▶ We define the operator $\mathbf{A}^\delta : D(\mathbf{A}^\delta) \rightarrow L^2(\Omega)$ such that

$$\left| \begin{array}{l} D(\mathbf{A}^\delta) = \{u \in \mathbf{H}_0^1(\Omega) \mid \operatorname{div}(\sigma^\delta \nabla u) \in L^2(\Omega)\} \\ \mathbf{A}^\delta u = \operatorname{div}(\sigma^\delta \nabla u). \end{array} \right.$$

- In the geometry with a **rounded corner**, we consider the **spectral** problem

$$\left| \begin{array}{l} \text{Find } (\lambda^\delta, u^\delta) \in \mathbb{C} \times (H_0^1(\Omega) \setminus \{0\}) \text{ s.t.:} \\ -\operatorname{div}(\sigma^\delta \nabla u^\delta) = \lambda^\delta u^\delta \quad \text{in } \Omega. \end{array} \right.$$

- We define the operator $A^\delta : D(A^\delta) \rightarrow L^2(\Omega)$ such that

$$\left| \begin{array}{l} D(A^\delta) = \{u \in H_0^1(\Omega) \mid \operatorname{div}(\sigma^\delta \nabla u) \in L^2(\Omega)\} \\ A^\delta u = \operatorname{div}(\sigma^\delta \nabla u). \end{array} \right.$$

PROPOSITION. Assume that $\kappa_\sigma \neq -1$. For $\delta > 0$ (in this case the interface is “smooth”), the operator A^δ is **selfadjoint** and has **compact resolvent**. Its spectrum $\mathfrak{S}(A^\delta)$ consists in two sequences of **isolated eigenvalues**:

$$-\infty \xleftarrow{n \rightarrow +\infty} \dots \lambda_{-n}^\delta \leq \dots \leq \lambda_{-1}^\delta < 0 \leq \lambda_1^\delta \leq \lambda_2^\delta \leq \dots \leq \lambda_n^\delta \dots \xrightarrow{n \rightarrow +\infty} +\infty.$$

- In the geometry with a **rounded corner**, we consider the **spectral** problem

$$\left| \begin{array}{l} \text{Find } (\lambda^\delta, u^\delta) \in \mathbb{C} \times (H_0^1(\Omega) \setminus \{0\}) \text{ s.t.:} \\ -\operatorname{div}(\sigma^\delta \nabla u^\delta) = \lambda^\delta u^\delta \quad \text{in } \Omega. \end{array} \right.$$

- We define the operator $A^\delta : D(A^\delta) \rightarrow L^2(\Omega)$ such that

$$\left| \begin{array}{l} D(A^\delta) = \{u \in H_0^1(\Omega) \mid \operatorname{div}(\sigma^\delta \nabla u) \in L^2(\Omega)\} \\ A^\delta u = \operatorname{div}(\sigma^\delta \nabla u). \end{array} \right.$$

PROPOSITION. Assume that $\kappa_\sigma \neq -1$. For $\delta > 0$ (in this case the interface is “smooth”), the operator A^δ is **selfadjoint** and has **compact resolvent**. Its spectrum $\mathfrak{S}(A^\delta)$ consists in two sequences of **isolated eigenvalues**:

$$-\infty \xleftarrow{n \rightarrow +\infty} \dots \lambda_{-n}^\delta \leq \dots \leq \lambda_{-1}^\delta < 0 \leq \lambda_1^\delta \leq \lambda_2^\delta \leq \dots \leq \lambda_n^\delta \dots \xrightarrow{n \rightarrow +\infty} +\infty.$$

? For $n \in \mathbb{Z}^*$, what is the **behaviour** of λ_n^δ when δ tends to zero?

- ▶ In the geometry with a **rounded corner**, we consider the **spectral** problem

$$\left| \begin{array}{l} \text{Find } (\lambda^\delta, u^\delta) \in \mathbb{C} \times (H_0^1(\Omega) \setminus \{0\}) \text{ s.t.:} \\ -\operatorname{div}(\sigma^\delta \nabla u^\delta) = \lambda^\delta u^\delta \quad \text{in } \Omega. \end{array} \right.$$

- ▶ We define the operator $A^\delta : D(A^\delta) \rightarrow L^2(\Omega)$ such that

$$\left| \begin{array}{l} D(A^\delta) = \{u \in H_0^1(\Omega) \mid \operatorname{div}(\sigma^\delta \nabla u) \in L^2(\Omega)\} \\ A^\delta u = \operatorname{div}(\sigma^\delta \nabla u). \end{array} \right.$$

PROPOSITION. Assume that $\kappa_\sigma \neq -1$. For $\delta > 0$ (in this case the interface is “smooth”), the operator A^δ is **selfadjoint** and has **compact resolvent**. Its spectrum $\mathfrak{S}(A^\delta)$ consists in two sequences of **isolated eigenvalues**:

$$-\infty \xleftarrow{n \rightarrow +\infty} \dots \lambda_{-n}^\delta \leq \dots \leq \lambda_{-1}^\delta < 0 \leq \lambda_1^\delta \leq \lambda_2^\delta \leq \dots \leq \lambda_n^\delta \dots \xrightarrow{n \rightarrow +\infty} +\infty.$$



For $n \in \mathbb{Z}^*$, what is the **behaviour** of λ_n^δ when δ tends to zero?

\Rightarrow This depends on the features of the **limit operator** for $\delta = 0$...

- Let $A : D(A) \rightarrow L^2(\Omega)$ denote the **limit operator** ($\delta = 0$) such that

$$\left| \begin{array}{l} D(A) = \{u \in H_0^1(\Omega) \mid \operatorname{div}(\sigma \nabla u) \in L^2(\Omega)\} \\ Au = \operatorname{div}(\sigma \nabla u). \end{array} \right.$$

- For $\delta = 0$, the interface is no longer “smooth” and the properties of A depend on the **values of κ_σ** :

- Let $A : D(A) \rightarrow L^2(\Omega)$ denote the **limit operator** ($\delta = 0$) such that

$$\begin{cases} D(A) = \{u \in H_0^1(\Omega) \mid \operatorname{div}(\sigma \nabla u) \in L^2(\Omega)\} \\ Au = \operatorname{div}(\sigma \nabla u). \end{cases}$$

- For $\delta = 0$, the interface is no longer “smooth” and the properties of A depend on the **values of κ_σ** :

♣ When $\kappa_\sigma \notin I_c$, A is **selfadjoint** and has **compact resolvent**. Its spectrum $\mathfrak{S}(A)$ consists in two sequences of **isolated eigenvalues**:

$$-\infty \xleftarrow{n \rightarrow +\infty} \dots \lambda_{-n} \leq \dots \leq \lambda_{-1} < 0 \leq \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n \dots \xrightarrow{n \rightarrow +\infty} +\infty.$$

In this case, there holds $\mathfrak{S}(A^\delta) \xrightarrow{\delta \rightarrow 0} \mathfrak{S}(A)$.

- Let $A : D(A) \rightarrow L^2(\Omega)$ denote the **limit operator** ($\delta = 0$) such that

$$\begin{cases} D(A) = \{u \in H_0^1(\Omega) \mid \operatorname{div}(\sigma \nabla u) \in L^2(\Omega)\} \\ Au = \operatorname{div}(\sigma \nabla u). \end{cases}$$

- For $\delta = 0$, the interface is no longer “smooth” and the properties of A depend on the **values of κ_σ** :

♣ When $\kappa_\sigma \notin I_c$, A is **selfadjoint** and has **compact resolvent**. Its spectrum $\mathfrak{S}(A)$ consists in two sequences of **isolated eigenvalues**:

$$-\infty \xleftarrow{n \rightarrow +\infty} \dots \lambda_{-n} \leq \dots \leq \lambda_{-1} < 0 \leq \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n \dots \xrightarrow{n \rightarrow +\infty} +\infty.$$

In this case, there holds $\mathfrak{S}(A^\delta) \xrightarrow{\delta \rightarrow 0} \mathfrak{S}(A)$.

♣ When $\kappa_\sigma \in I_c \setminus \{-1\}$, there holds $D(A^*) = D(A) \oplus \operatorname{span}(s_+, s_-)$ where $s_\pm = \zeta r^{\pm i\eta} \varphi(\theta)$ (in particular A is **not selfadjoint**). Moreover, $\mathfrak{S}(A) = \mathbb{C}$.

INSIDE THE CRITICAL INTERVAL:

① The selfadjoint extensions of A are the operators $A(\tau) : D(A(\tau)) \rightarrow L^2(\Omega)$, $\tau \in \mathbb{R}$, such that

$$\left| \begin{array}{l} D(A(\tau)) = D(A) \oplus \text{span}(s_+ + e^{i\tau} s_-) \\ A(\tau)u = \text{div}(\sigma \nabla u). \end{array} \right.$$

► For $\delta = 0$, the operator is "smooth" and the properties of A depend on the values of κ_σ :

♣ When $\kappa_\sigma \in I_\sigma$, A is selfadjoint and has compact resolvent. Its spectrum $\mathfrak{S}(A)$ consists in two sequences of isolated eigenvalues:

$$-\infty \leftarrow_{n \rightarrow +\infty} \dots \lambda_{-n} \leq \dots \leq \lambda_{-1} < 0 \leq \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n \dots \rightarrow_{n \rightarrow +\infty} +\infty.$$

In this case, there holds $\mathfrak{S}(A^\delta) \xrightarrow{\delta \rightarrow 0} \mathfrak{S}(A)$.

♣ When $\kappa_\sigma \in I_\sigma \setminus \{-1\}$, there holds $D(A^*) = D(A) \oplus \text{span}(s_+, s_-)$ where $s_\pm \in W^{1,2}(\Omega)$ are the solutions of $A s_\pm = \pm s_\pm$. Moreover, $\mathfrak{S}(A^*) = \mathfrak{S}(A) \cup \{-1\}$.

INSIDE THE CRITICAL INTERVAL:

- ① The selfadjoint extensions of A are the operators $A(\tau) : D(A(\tau)) \rightarrow L^2(\Omega)$, $\tau \in \mathbb{R}$, such that

$$\begin{cases} D(A(\tau)) = D(A) \oplus \text{span}(s_+ + e^{i\tau} s_-) \\ A(\tau)u = \text{div}(\sigma \nabla u). \end{cases}$$

- ② Using matched asymptotic expansions techniques, we have proved the

THEOREM. Assume that $\kappa_\sigma \in I_c \setminus \{-1\}$. There exist $a \neq 0, b \in \mathbb{R}$, such that $\text{dist}(\mathfrak{S}(A^\delta), \mathfrak{S}(A(a \ln \delta + b))) \rightarrow 0$ on each compact set of \mathbb{R} as $\delta \rightarrow 0$.

In this case, there holds $\mathfrak{S}(A^\delta) \xrightarrow{\delta \rightarrow 0} \mathfrak{S}(A)$.

⊛ When $\kappa_\sigma \in I_c \setminus \{-1\}$, there holds $D(A^*) = D(A) \oplus \text{span}(s_+, s_-)$ where $s_\pm \in H^1(\Omega) \cap H^2(\partial\Omega)$ satisfy $A s_\pm = 0$, $\text{div}(\sigma \nabla s_\pm) = 0$, $\mathfrak{S}(A^*) = \mathfrak{S}(A) \cup \{0\}$.

INSIDE THE CRITICAL INTERVAL:

- ① The **selfadjoint extensions** of A are the operators $A(\tau) : D(A(\tau)) \rightarrow L^2(\Omega)$, $\tau \in \mathbb{R}$, such that

$$\left| \begin{array}{l} D(A(\tau)) = D(A) \oplus \text{span}(s_+ + e^{i\tau} s_-) \\ A(\tau)u = \text{div}(\sigma \nabla u). \end{array} \right.$$

- ② Using **matched asymptotic expansions techniques**, we have proved the

THEOREM. Assume that $\kappa_\sigma \in I_c \setminus \{-1\}$. There exist $a \neq 0, b \in \mathbb{R}$, such that $\text{dist}(\mathfrak{S}(A^\delta), \mathfrak{S}(A(a \ln \delta + b))) \rightarrow 0$ on each compact set of \mathbb{R} as $\delta \rightarrow 0$.

- ③ Conclusion.



The spectrum of A^δ **does not converge** when $\delta \rightarrow 0$. Asymptotically, $\mathfrak{S}(A^\delta)$ is $2\pi/a$ -periodic in $\ln \delta$ -scale.

- Let $A : D(A) \rightarrow L^2(\Omega)$ denote the **limit operator** ($\delta = 0$) such that

$$\left\{ \begin{array}{l} D(A) = \{u \in H_0^1(\Omega) \mid \operatorname{div}(\sigma \nabla u) \in L^2(\Omega)\} \\ Au = \operatorname{div}(\sigma \nabla u). \end{array} \right.$$

- For $\delta = 0$, the interface is no longer “smooth” and the properties of A depend on the **values of κ_σ** :



♣ When $\kappa_\sigma \notin I_c$, A is **selfadjoint** and has **compact resolvent**. Its spectrum $\mathfrak{S}(A)$ consists in two sequences of **isolated eigenvalues**:

$$-\infty \xleftarrow{n \rightarrow +\infty} \dots \lambda_{-n} \leq \dots \leq \lambda_{-1} < 0 \leq \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n \dots \xrightarrow{n \rightarrow +\infty} +\infty.$$

In this case, there holds $\mathfrak{S}(A^\delta) \xrightarrow{\delta \rightarrow 0} \mathfrak{S}(A)$.


♣ When $\kappa_\sigma \in I_c \setminus \{-1\}$, the spectrum of A^δ **does not converge** when $\delta \rightarrow 0$. Asymptotically, $\mathfrak{S}(A^\delta)$ is $2\pi/a$ -**periodic** in $\ln \delta$ -**scale**.

$$\kappa_\sigma = -1.0001 \text{ (outside the critical interval)}$$



- $\mathfrak{S}(A^\delta)$ converges to $\mathfrak{S}(A)$ (A is the limit operator) when $\delta \rightarrow 0$.


$\kappa_\sigma = -0.9999$ (inside the critical interval)



- Asymptotically, $\mathfrak{S}(A^\delta)$ is periodic in $\ln \delta$ -scale as $\delta \rightarrow 0$.


Spectral problem: numerical experiments 4/4

$$\kappa_\sigma = -0.9999 \text{ (inside the critical interval)}$$



- Asymptotically, $\mathfrak{S}(A^\delta)$ is periodic in $\ln \delta$ -scale as $\delta \rightarrow 0$.

$$\kappa_\sigma = -0.9999 \text{ (inside the critical interval)}$$



- Asymptotically, $\mathfrak{S}(A^\delta)$ is periodic in $\ln \delta$ -scale as $\delta \rightarrow 0$.

- 1 Numerical experiments
- 2 Properties of the limit problem
- 3 Asymptotic analysis

Conclusion

Let us remind the initial question:



What is the **behaviour** of $(u^\delta)_\delta$ when δ tends to zero?

Conclusion

Let us remind the initial question:



What is the **behaviour** of $(u^\delta)_\delta$ when δ tends to zero?



This depends on the features of the **limit problem**.

(...)

(...)

$$\kappa_\sigma = -1.0001 \notin I_c$$

$$\kappa_\sigma = -0.9999 \in I_c$$



When $\kappa_\sigma \in I_c$, $(u^\delta)_\delta$ **does not converge**, even for the L^2 -norm!

In this case, it is impossible to **simulate** the fields since it is impossible to **measure** exactly δ . \Rightarrow What happens **physically**?

Thank you for your attention!!!

► ANR project Metamath coordinated by **S. Fliss**.



A.-S. Bonnet-Ben Dhia, L. Chesnel, P. Ciarlet Jr., *T-coercivity for scalar interface problems between dielectrics and metamaterials*, *M2AN*, 46, 1363–1387, 2012.



A.-S. Bonnet-Ben Dhia, L. Chesnel, X. Claeys, *Radiation condition for a non-smooth interface between a dielectric and a metamaterial*, *M3AS*, 23, 2013.




L. Chesnel, X. Claeys, S.A. Nazarov, *A curious instability phenomenon for a rounded corner in presence of a negative material*, *Asymp. Anal.*, in press, 2013.

Spectrum for a small inclusion: setting

- ▶ Let Ω, Ξ be **smooth** domains of \mathbb{R}^3 such that $O \in \Xi, \overline{\Xi} \subset \Omega$.
- ▶ For $\delta \in (0; 1]$, we consider the **spectral** problem

$$\left| \begin{array}{l} \text{Find } (\lambda^\delta, u^\delta) \in \mathbb{C} \times (H_0^1(\Omega) \setminus \{0\}) \text{ s.t.:} \\ -\operatorname{div}(\sigma^\delta \nabla u^\delta) = \lambda^\delta u^\delta \quad \text{in } \Omega, \end{array} \right.$$

$$\text{where } \sigma^\delta = \begin{cases} \sigma_1 > 0 & \text{in } \Omega_1^\delta := \Omega \setminus \overline{\delta \Xi} \\ \sigma_2 < 0 & \text{in } \Omega_2^\delta := \delta \Xi. \end{cases}$$



Spectrum for a small inclusion: setting


- ▶ Let Ω , Ξ be **smooth** domains of \mathbb{R}^3 such that $O \in \Xi$, $\bar{\Xi} \subset \Omega$.
- ▶ For $\delta \in (0; 1]$, we consider the **spectral** problem

$$\left| \begin{array}{l} \text{Find } (\lambda^\delta, u^\delta) \in \mathbb{C} \times (H_0^1(\Omega) \setminus \{0\}) \text{ s.t.:} \\ -\operatorname{div}(\sigma^\delta \nabla u^\delta) = \lambda^\delta u^\delta \quad \text{in } \Omega, \end{array} \right.$$

$$\text{where } \sigma^\delta = \begin{cases} \sigma_1 > 0 & \text{in } \Omega_1^\delta := \Omega \setminus \bar{\delta \Xi} \\ \sigma_2 < 0 & \text{in } \Omega_2^\delta := \delta \Xi. \end{cases}$$

- ▶ We define the operator $A^\delta : D(A^\delta) \rightarrow L^2(\Omega)$ such that

$$\left| \begin{array}{l} D(A^\delta) = \{u \in H_0^1(\Omega) \mid \operatorname{div}(\sigma^\delta \nabla u) \in L^2(\Omega)\} \\ A^\delta u = \operatorname{div}(\sigma^\delta \nabla u). \end{array} \right.$$




Spectrum for a small inclusion: setting

- ▶ Let Ω , Ξ be **smooth** domains of \mathbb{R}^3 such that $O \in \Xi$, $\bar{\Xi} \subset \Omega$.
- ▶ For $\delta \in (0; 1]$, we consider the **spectral** problem

$$\left| \begin{array}{l} \text{Find } (\lambda^\delta, u^\delta) \in \mathbb{C} \times (H_0^1(\Omega) \setminus \{0\}) \text{ s.t.:} \\ -\operatorname{div}(\sigma^\delta \nabla u^\delta) = \lambda^\delta u^\delta \quad \text{in } \Omega, \end{array} \right.$$

$$\text{where } \sigma^\delta = \begin{cases} \sigma_1 > 0 & \text{in } \Omega_1^\delta := \Omega \setminus \bar{\delta \Xi} \\ \sigma_2 < 0 & \text{in } \Omega_2^\delta := \delta \Xi. \end{cases}$$



- ▶ We define the operator $A^\delta : D(A^\delta) \rightarrow L^2(\Omega)$ such that

$$\left| \begin{array}{l} D(A^\delta) = \{u \in H_0^1(\Omega) \mid \operatorname{div}(\sigma^\delta \nabla u) \in L^2(\Omega)\} \\ A^\delta u = \operatorname{div}(\sigma^\delta \nabla u). \end{array} \right.$$

PROPOSITION. Assume that $\kappa_\sigma \neq -1$. For $\delta > 0$, the operator A^δ is **selfadjoint** and has **compact resolvent**. Its spectrum $\mathfrak{S}(A^\delta)$ consists in two sequences of **isolated eigenvalues**:


$$-\infty \xleftarrow{n \rightarrow +\infty} \dots \lambda_{-n}^\delta \leq \dots \leq \lambda_{-1}^\delta < 0 \leq \lambda_1^\delta \leq \lambda_2^\delta \leq \dots \leq \lambda_n^\delta \dots \xrightarrow{n \rightarrow +\infty} +\infty.$$

Spectrum for a small inclusion: setting

- ▶ Let Ω , Ξ be **smooth** domains of \mathbb{R}^3 such that $O \in \Xi$, $\bar{\Xi} \subset \Omega$.
- ▶ For $\delta \in (0; 1]$, we consider the **spectral** problem

$$\left| \begin{array}{l} \text{Find } (\lambda^\delta, u^\delta) \in \mathbb{C} \times (H_0^1(\Omega) \setminus \{0\}) \text{ s.t.:} \\ -\operatorname{div}(\sigma^\delta \nabla u^\delta) = \lambda^\delta u^\delta \quad \text{in } \Omega, \end{array} \right.$$

$$\text{where } \sigma^\delta = \begin{cases} \sigma_1 > 0 & \text{in } \Omega_1^\delta := \Omega \setminus \bar{\delta \Xi} \\ \sigma_2 < 0 & \text{in } \Omega_2^\delta := \delta \Xi. \end{cases}$$



- ▶ We define the operator $A^\delta : D(A^\delta) \rightarrow L^2(\Omega)$ such that

$$\left| \begin{array}{l} D(A^\delta) = \{u \in H_0^1(\Omega) \mid \operatorname{div}(\sigma^\delta \nabla u) \in L^2(\Omega)\} \\ A^\delta u = \operatorname{div}(\sigma^\delta \nabla u). \end{array} \right.$$


- ▶ For all $\delta \in (0; 1]$, A^δ has **negative spectrum**. At the limit $\delta = 0$, the **inclusion of negative material vanishes** and σ is **strictly positive**.

Spectrum for a small inclusion: setting

- ▶ Let Ω , Ξ be **smooth** domains of \mathbb{R}^3 such that $O \in \Xi$, $\bar{\Xi} \subset \Omega$.
- ▶ For $\delta \in (0; 1]$, we consider the **spectral** problem

$$\left| \begin{array}{l} \text{Find } (\lambda^\delta, u^\delta) \in \mathbb{C} \times (H_0^1(\Omega) \setminus \{0\}) \text{ s.t.:} \\ -\operatorname{div}(\sigma^\delta \nabla u^\delta) = \lambda^\delta u^\delta \quad \text{in } \Omega, \end{array} \right.$$

$$\text{where } \sigma^\delta = \begin{cases} \sigma_1 > 0 & \text{in } \Omega_1^\delta := \Omega \setminus \bar{\delta \Xi} \\ \sigma_2 < 0 & \text{in } \Omega_2^\delta := \delta \Xi. \end{cases}$$



- ▶ We define the operator $A^\delta : D(A^\delta) \rightarrow L^2(\Omega)$ such that

$$\left| \begin{array}{l} D(A^\delta) = \{u \in H_0^1(\Omega) \mid \operatorname{div}(\sigma^\delta \nabla u) \in L^2(\Omega)\} \\ A^\delta u = \operatorname{div}(\sigma^\delta \nabla u). \end{array} \right.$$

- ▶ For all $\delta \in (0; 1]$, A^δ has **negative spectrum**. At the limit $\delta = 0$, the **inclusion of negative material vanishes** and σ is **strictly positive**.




What happens to the negative spectrum when δ tends to zero?

Limit operators

► As $\delta \rightarrow 0$, the **small inclusion of negative material disappears**. We introduce the **far field operator \mathbf{A}^0** such that

$$\left| \begin{array}{l} D(\mathbf{A}^0) = \{v \in H_0^1(\Omega) \mid \Delta v \in L^2(\Omega)\} \\ \mathbf{A}^0 v = -\sigma_1 \Delta v. \end{array} \right.$$




There holds $\mathfrak{S}(\mathbf{A}^0) = \{\mu_n\}_{n \geq 1}$ with $0 < \mu_1 < \mu_2 \leq \dots \leq \mu_n \dots \xrightarrow{n \rightarrow +\infty} +\infty$.

Limit operators

► As $\delta \rightarrow 0$, the **small inclusion of negative material disappears**. We introduce the **far field operator A^0** such that


$$\left| \begin{array}{l} D(A^0) = \{v \in H_0^1(\Omega) \mid \Delta v \in L^2(\Omega)\} \\ A^0 v = -\sigma_1 \Delta v. \end{array} \right.$$



There holds $\mathfrak{S}(A^0) = \{\mu_n\}_{n \geq 1}$ with $0 < \mu_1 < \mu_2 \leq \dots \leq \mu_n \dots \xrightarrow{n \rightarrow +\infty} +\infty$.

► Introduce the **rapid coordinate $\xi := \delta^{-1} \mathbf{x}$** and let $\delta \rightarrow 0$. Define the **near field operator B^∞** such that

$$\left| \begin{array}{l} D(B^\infty) := \{w \in H^1(\mathbb{R}^3) \mid \operatorname{div}(\sigma^\infty \nabla w) \in L^2(\mathbb{R}^3)\} \\ B^\infty w = -\operatorname{div}(\sigma^\infty \nabla w). \end{array} \right.$$




$$\sigma^\infty = \sigma_1$$

Limit operators

► As $\delta \rightarrow 0$, the **small inclusion of negative material disappears**. We introduce the **far field operator** \mathbf{A}^0 such that


$$\left| \begin{array}{l} D(\mathbf{A}^0) = \{v \in H_0^1(\Omega) \mid \Delta v \in L^2(\Omega)\} \\ \mathbf{A}^0 v = -\sigma_1 \Delta v. \end{array} \right.$$



There holds $\mathfrak{S}(\mathbf{A}^0) = \{\mu_n\}_{n \geq 1}$ with $0 < \mu_1 < \mu_2 \leq \dots \leq \mu_n \dots \xrightarrow{n \rightarrow +\infty} +\infty$.

► Introduce the **rapid coordinate** $\xi := \delta^{-1} \mathbf{x}$ and let $\delta \rightarrow 0$. Define the **near field operator** \mathbf{B}^∞ such that

$$\left| \begin{array}{l} D(\mathbf{B}^\infty) := \{w \in H^1(\mathbb{R}^3) \mid \operatorname{div}(\sigma^\infty \nabla w) \in L^2(\mathbb{R}^3)\} \\ \mathbf{B}^\infty w = -\operatorname{div}(\sigma^\infty \nabla w). \end{array} \right.$$



$$\sigma^\infty = \sigma_1$$

PROPOSITION. Assume that $\kappa_\sigma \neq -1$. The **continuous spectrum** of \mathbf{B}^∞ is equal to $[0; +\infty)$ while its **discrete spectrum** is a sequence of eigenvalues:

$$\mathfrak{S}(\mathbf{B}^\infty) \setminus \overline{\mathbb{R}_+} = \{\mu_{-n}\}_{n \geq 1} \quad \text{with} \quad 0 > \mu_{-1} \geq \dots \geq \mu_{-n} \dots \xrightarrow{n \rightarrow +\infty} -\infty.$$

Spectrum for a small inclusion: results

Assume that $\kappa_\sigma \neq -1$ and that \mathbf{B}^∞ is injective. For $n \in \mathbb{N}^*$, we denote $\lambda_{\pm n}^\delta$, μ_n^δ , μ_{-n}^δ the eigenvalues of \mathbf{A}^δ , \mathbf{A}^0 , \mathbf{B}^∞ as in the previous slides.

THEOREM. (POSITIVE SPECTRUM) For all $n \in \mathbb{N}^*$, $\varepsilon \in (0; 1)$, there exist constants $C, \delta_0 > 0$ depending on n, ε but independent of δ , such that

$$|\lambda_n^\delta - \mu_n| \leq C \delta^{3/2-\varepsilon}, \quad \forall \delta \in (0; \delta_0].$$

Spectrum for a small inclusion: results

Assume that $\kappa_\sigma \neq -1$ and that \mathbf{B}^∞ is injective. For $n \in \mathbb{N}^*$, we denote $\lambda_{\pm n}^\delta$, μ_n^δ , μ_{-n}^δ the eigenvalues of \mathbf{A}^δ , \mathbf{A}^0 , \mathbf{B}^∞ as in the previous slides.

THEOREM. (POSITIVE SPECTRUM) For all $n \in \mathbb{N}^*$, $\varepsilon \in (0; 1)$, there exist constants $C, \delta_0 > 0$ depending on n, ε but independent of δ , such that

$$|\lambda_n^\delta - \mu_n| \leq C \delta^{3/2-\varepsilon}, \quad \forall \delta \in (0; \delta_0].$$

THEOREM. (NEGATIVE SPECTRUM) For all $n \in \mathbb{N}^*$, there exist constants $C, \gamma, \delta_0 > 0$, depending on n but independent of δ , such that

$$|\lambda_{-n}^\delta - \delta^{-2}\mu_{-n}| \leq C \exp(-\gamma/\delta), \quad \forall \delta \in (0; \delta_0].$$

Spectrum for a small inclusion: results

Assume that $\kappa_\sigma \neq -1$ and that B^∞ is injective. For $n \in \mathbb{N}^*$, we denote $\lambda_{\pm n}^\delta$, μ_n^δ , μ_{-n}^δ the eigenvalues of A^δ , A^0 , B^∞ as in the previous slides.

SCHEMATICALLY, WE HAVE:

$$\mathfrak{S}(A^\delta) \quad \begin{array}{cccc} \lambda_{-2}^\delta & & \lambda_{-1}^\delta & & & & \lambda_1^\delta & & \lambda_2^\delta \\ \cdot \times & \cdots & \times & \cdots & \cdots & | & \times & \cdots & \times \rightarrow \end{array}$$

$\delta \rightarrow 0$

$$\begin{array}{cccc} \delta^{-2}\mu_{-2} & & \delta^{-2}\mu_{-1} & & & & \mu_1 & & \mu_2 \\ \cdot \times & \cdots & \times & \cdots & \cdots & | & \times & \cdots & \times \rightarrow \end{array}$$

$$\delta^{-2}\mathfrak{S}(B^\infty) \cap (-\infty; 0)$$

$$\mathfrak{S}(A^0)$$

Spectrum for a small inclusion: results

Assume that $\kappa_\sigma \neq -1$ and that \mathbf{B}^∞ is injective. For $n \in \mathbb{N}^*$, we denote $\lambda_{\pm n}^\delta$, μ_n^δ , μ_{-n}^δ the eigenvalues of \mathbf{A}^δ , \mathbf{A}^0 , \mathbf{B}^∞ as in the previous slides.

THEOREM. (POSITIVE SPECTRUM) For all $n \in \mathbb{N}^*$, $\varepsilon \in (0; 1)$, there exist constants $C, \delta_0 > 0$ depending on n, ε but independent of δ , such that

$$|\lambda_n^\delta - \mu_n| \leq C \delta^{3/2-\varepsilon}, \quad \forall \delta \in (0; \delta_0].$$

THEOREM. (NEGATIVE SPECTRUM) For all $n \in \mathbb{N}^*$, there exist constants $C, \gamma, \delta_0 > 0$, depending on n but independent of δ , such that

$$|\lambda_{-n}^\delta - \delta^{-2}\mu_{-n}| \leq C \exp(-\gamma/\delta), \quad \forall \delta \in (0; \delta_0].$$

Spectrum for a small inclusion: results

Assume that $\kappa_\sigma \neq -1$ and that \mathbf{B}^∞ is injective. For $n \in \mathbb{N}^*$, we denote $\lambda_{\pm n}^\delta$, μ_n^δ , μ_{-n}^δ the eigenvalues of A^δ , A^0 , B^∞ as in the previous slides.

THEOREM. (POSITIVE SPECTRUM) For all $n \in \mathbb{N}^*$, $\varepsilon \in (0; 1)$, there exist constants $C, \delta_0 > 0$ depending on n, ε but independent of δ , such that

$$|\lambda_n^\delta - \mu_n| \leq C \delta^{3/2-\varepsilon}, \quad \forall \delta \in (0; \delta_0].$$

THEOREM. (NEGATIVE SPECTRUM) For all $n \in \mathbb{N}^*$, there exist constants $C, \gamma, \delta_0 > 0$, depending on n but independent of δ , such that


$$|\lambda_{-n}^\delta - \delta^{-2} \mu_{-n}| \leq C \exp(-\gamma/\delta), \quad \forall \delta \in (0; \delta_0].$$

PROPOSITION. (LOCALIZATION EFFECT) For all $n \in \mathbb{N}^*$, let u_{-n}^δ be an eigenfunction corresponding to the negative eigenvalue λ_{-n}^δ . There exist constants $C, \gamma, \delta_0 > 0$, depending on n but independent of δ , such that


$$\int_{\Omega} (|u_{-n}^\delta|^2 + |\nabla u_{-n}^\delta|^2) e^{\gamma x/\delta} dx \leq C \|u_{-n}^\delta\|_{\Omega}, \quad \forall \delta \in (0; \delta_0].$$

Numerical experiments for the small inclusion

- ▶ We **approximate numerically** the spectrum of A^δ using a **usual P1 Finite Element Method** and we make δ goes to zero.
- ▶ We consider the following **2D geometry**:




Numerical experiments for the small inclusion




- The **positive part** of $\mathfrak{S}(A^\delta)$ converges to $\mathfrak{S}(A^0)$ when $\delta \rightarrow 0$.

Numerical experiments for the small inclusion



- The **negative part** of $\mathfrak{S}(A^\delta)$ is asymptotically equivalent to the **negative part** of $\delta^{-2}\mathfrak{S}(B^\infty)$ when $\delta \rightarrow 0$.

Numerical experiments for the small inclusion




- The **negative part** of $\mathfrak{S}(A^\delta)$ is asymptotically equivalent to the **negative part** of $\delta^{-2}\mathfrak{S}(B^\infty)$ when $\delta \rightarrow 0$.

Localization effect


Eigenfunction associated to the first **negative eigenvalue**

$\delta=0.5$




Eigenfunction associated to the first **positive eigenvalue**


$\delta=0.5$



$\delta=0.05$



$\delta=0.05$



- The **eigenfunctions** corresponding to the **negative eigenvalues** are **localized** around the small inclusion. Here, $\kappa_\sigma = -2.5$.