

Constrained Optimization

Equality Constraint

Objective:

Generalize the necessary condition of $\nabla f(x) = 0$ at the optima of f when f is in \mathcal{C}^1 , i.e. is differentiable and its differential is continuous

Theorem:

Be U an open set of $(E, \|\cdot\|)$, and $f: U \rightarrow \mathbb{R}$, $g: U \rightarrow \mathbb{R}$ in \mathcal{C}^1 .

Let $a \in E$ satisfy

$$\begin{cases} f(a) = \inf \{f(x) \mid x \in \mathbb{R}^n, g(x) = 0\} \\ g(a) = 0 \end{cases}$$

i.e. a is optimum of the problem

If $\nabla g(a) \neq 0$, then there exists a constant $\lambda \in \mathbb{R}$ called *Lagrange multiplier*, such that

$$\underbrace{\nabla f(a) + \lambda \nabla g(a)} = 0 \quad \text{Euler – Lagrange equation}$$

i.e. gradients of f and g in a are colinear

Interpretation of Euler-Lagrange Equation

Intuitive way to retrieve the Euler-Lagrange equation:

- In a local minimum a of a constrained problem, the hypersurfaces (or level sets) $f = f(a)$ and $g = 0$ are necessarily tangent (otherwise we could decrease f by moving along $g = 0$).
- Since the gradients $\nabla f(a)$ and $\nabla g(a)$ are orthogonal to the level sets $f = f(a)$ and $g = 0$, it follows that $\nabla f(a)$ and $\nabla g(a)$ are colinear.

Generalization to More than One Constraint

Theorem

- Assume $f: U \rightarrow \mathbb{R}$ and $g_k: U \rightarrow \mathbb{R}$ ($1 \leq k \leq p$) are \mathcal{C}^1 .
- Let a be such that
$$\begin{cases} f(a) = \min_{\substack{x \in \mathbb{R}^n, \\ g_k(x) = 0, \quad 1 \leq k \leq p}} \{f(x) \mid x \in \mathbb{R}^n, \quad g_k(x) = 0, \quad 1 \leq k \leq p\} \\ g_k(a) = 0 \text{ for all } 1 \leq k \leq p \end{cases}$$
- If $(\nabla g_k(a))_{1 \leq k \leq p}$ are linearly independent, then there exist p real constants $(\lambda_k)_{1 \leq k \leq p}$ such that

$$\nabla f(a) + \sum_{k=1}^p \lambda_k \nabla g_k(a) = 0$$

↑
Lagrange multiplier

again: a does not need to be global but local minimum

The Lagrangian

- Define the Lagrangian on $\mathbb{R}^n \times \mathbb{R}^p$ as

$$\mathcal{L}(x, \{\lambda_k\}) = f(x) + \underbrace{\sum_{k=1}^p \lambda_k g_k(x)}_{\lambda^T g(x)} \in \mathbb{R}$$

- To find optimal solutions, we can solve the optimality system

$$\left\{ \begin{array}{l} \text{Find } (x, \{\lambda_k\}) \in \mathbb{R}^n \times \mathbb{R}^p \text{ such that } \nabla f(x) + \sum_{k=1}^p \lambda_k \nabla g_k(x) = 0 \\ g_k(x) = 0 \text{ for all } 1 \leq k \leq p \end{array} \right.$$

$$\Leftrightarrow \left\{ \begin{array}{l} \text{Find } (x, \{\lambda_k\}) \in \mathbb{R}^n \times \mathbb{R}^p \text{ such that } \nabla_x \mathcal{L}(x, \{\lambda_k\}) = 0 \\ \nabla_{\lambda_k} \mathcal{L}(x, \{\lambda_k\})(x) = 0 \text{ for all } 1 \leq k \leq p \end{array} \right.$$

Inequality Constraint: Definitions

Let $\mathcal{U} = \{x \in \mathbb{R}^n \mid g_k(x) = 0 \text{ (for } k \in E), g_k(x) \leq 0 \text{ (for } k \in I)\}$.

Definition:

The points in \mathbb{R}^n that satisfy the constraints are also called *feasible* points.

Definition:

Let $a \in \mathcal{U}$, we say that the constraint $g_k(x) \leq 0$ (for $k \in I$) is *active* in a if $g_k(a) = 0$.

Inequality Constraint: Karush-Kuhn-Tucker Theorem

Theorem (Karush-Kuhn-Tucker, KKT):

Let U be an open set of $(\mathbb{R}^n, || \cdot ||)$ and $f: U \rightarrow \mathbb{R}$, $g_k: U \rightarrow \mathbb{R}$, all \mathcal{C}^1

Furthermore, let $a \in U$ satisfy

$$\left\{ \begin{array}{l} f(a) = \min_{x \in \mathbb{R}^n} (f(x) \mid g_k(x) = 0 \text{ (for } k \in E), g_k(x) \leq 0 \text{ (for } k \in I)) \\ g_k(a) = 0 \text{ (for } k \in E) \\ g_k(a) \leq 0 \text{ (for } k \in I) \end{array} \right. \quad \text{also works again for } a \text{ being a local minimum}$$

Let I_a^0 be the set of constraints that are active in a . Assume that $(\nabla g_k(a))_{k \in E \cup I_a^0}$ are linearly independent.

Then there exist $(\lambda_k)_{1 \leq k \leq p}$ that satisfy

$$\left\{ \begin{array}{l} \nabla f(a) + \sum_{k=1}^p \lambda_k \nabla g_k(a) = 0 \\ g_k(a) = 0 \text{ (for } k \in E) \\ g_k(a) \leq 0 \text{ (for } k \in I) \\ \lambda_k \geq 0 \text{ (for } k \in I_a^0) \\ \lambda_k g_k(a) = 0 \text{ (for } k \in E \cup I) \end{array} \right.$$

Inequality Constraint: Karush-Kuhn-Tucker Theorem

Theorem (Karush-Kuhn-Tucker, KKT):

Let U be an open set of $(E, || ||)$ and $f: U \rightarrow \mathbb{R}$, $g_k: U \rightarrow \mathbb{R}$, all C^1

Furthermore, let $a \in U$ satisfy

$$\left\{ \begin{array}{l} f(a) = \underset{\text{min}}{\cancel{f}}(f(x) \mid x \in \mathbb{R}^n, g_k(x) = 0 \text{ (for } k \in E), g_k(x) \leq 0 \text{ (for } k \in I)) \\ g_k(a) = 0 \text{ (for } k \in E) \\ g_k(a) \leq 0 \text{ (for } k \in I) \end{array} \right.$$

Let I_a^0 be the set of constraints that are active in a . Assume that $(\nabla g_k(a))_{k \in E \cup I_a^0}$ are linearly independent.

Then there exist $(\lambda_k)_{1 \leq k \leq p}$ that satisfy

$$\left\{ \begin{array}{l} \nabla f(a) + \sum_{k=1}^p \lambda_k \nabla g_k(a) = 0 \\ g_k(a) = 0 \text{ (for } k \in E) \\ g_k(a) \leq 0 \text{ (for } k \in I) \\ \lambda_k \geq 0 \text{ (for } k \in I_a^0) \\ \lambda_k g_k(a) = 0 \text{ (for } k \in E \cup I) \end{array} \right.$$

either active constraint
or $\lambda_k = 0$